

# *HIV Surveillance Report, 2001*



**Minnesota Department of Health  
HIV/AIDS Surveillance System**

## Introduction (I)

- These two introduction slides provide a general context for the data used in this slide set. If you have questions about any of the slides please refer to the *Companion Text to the Minnesota HIV Surveillance Report, 2001* or *HIV Surveillance Technical Notes*.
- This slide set describes new HIV infections (including AIDS at first diagnosis) in Minnesota by person, place, and time.
- The slides rely on data from HIV/AIDS cases diagnosed through 2001 and reported to the Minnesota Department of Health (MDH) HIV/AIDS Surveillance System.
- The data are displayed by year of HIV diagnosis.


## Introduction (II)

- Data analyses exclude persons diagnosed in federal or private correctional facilities, but include state prisoners (number of state prisoners believed to be living with HIV/AIDS = 101).
- Some limitations of surveillance data:
  - ◆ Data do not include HIV-infected persons who have not been tested for HIV
  - ◆ Data do not include persons whose positive test results have not been reported to the MDH
  - ◆ Case numbers for the most recent years may be undercounted due to delays in reporting

# National Context

# United States: State-Specific AIDS Rates

AIDS Rates per 100,000 Population Reported in 1999



SOURCE:  
L178 slide series (slide 12),  
National Center for HIV, STD, and TB Prevention, CDC

# Overview of HIV/AIDS in Minnesota

# Minnesota HIV/AIDS Surveillance: *Cumulative Cases*

- As of December 31, 2001, a cumulative total of 6,661\* persons have been diagnosed and reported with HIV infection in Minnesota
  - ◆ 2,807 persons diagnosed with HIV infection (non-AIDS)
  - ◆ 3,854 persons diagnosed with AIDS
- Of these 6,661 persons, 2,332 are known to be deceased

\* This number includes only persons who reported Minnesota as their state of residence at the time of their HIV and/or AIDS diagnosis.

## Estimated Number of Persons *Living with HIV/AIDS in Minnesota*


- As of December 31, 2001, 4,331\* persons are assumed alive and living in Minnesota with HIV/AIDS
  - ◆ 2,574 living with HIV infection (non-AIDS)
  - ◆ 1,757 living with AIDS
- This number includes 375 persons who were first reported with HIV or AIDS elsewhere and subsequently moved to Minnesota
- This number excludes 418 persons who were first reported with HIV or AIDS in Minnesota and subsequently moved out of the state

\* *This number includes persons who reported Minnesota as their current state of residence, regardless of residence at time of diagnosis.*



# HIV/AIDS in Minnesota:

Number of new cases, prevalent cases, and deaths by year, 1990-2001



\* Deaths among AIDS cases, regardless of cause.

Data Source: Minnesota HIV/AIDS Surveillance System


# **New HIV Infections\* in Minnesota by Person, Place, and Time**

*\* HIV or AIDS at first diagnosis*

*HIV/AIDS in Minnesota: Annual Review*

# Place

# New HIV Infections<sup>†</sup> by County of Residence at Diagnosis, 2001



<sup>†</sup> HIV or AIDS at first diagnosis

Data Source: Minnesota HIV/AIDS Surveillance System


HIV/AIDS in Minnesota: Annual Review

# Map of Metro Area: New HIV Infections<sup>†</sup> by County of Residence at Diagnosis, 2001

Number of  
New Infections



Total number (Metro only) = 254




<sup>†</sup> HIV or AIDS at first diagnosis

Data Source: Minnesota HIV/AIDS Surveillance System

\* Counties in which a state  
correctional facility is located.

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* in Minnesota by Residence at Diagnosis, 2001



Suburban = Seven-county metro area including Anoka, Carver, Dakota, Hennepin (except Minneapolis), Ramsey (except St. Paul), Scott, and Washington counties. Greater MN = All other Minnesota counties, outside the seven-county metro area.


\* HIV or AIDS at first diagnosis

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# Gender and Race/Ethnicity

# New HIV Infections\* by Gender and Year of Diagnosis, 1990-2001




\* HIV or AIDS at first diagnosis

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review



# New HIV Infections\* Among Males by Race/Ethnicity† and Year of Diagnosis, 1990-2001




\* HIV or AIDS at first diagnosis

† "African-born" refers to Blacks who reported an African country of birth; "African American" refers to all other Blacks. Cases with unknown race are excluded.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* Among Males by Race/Ethnicity† and Year of Diagnosis, 1990-2001 (excluding Whites)




\* HIV or AIDS at first diagnosis

† "African-born" refers to Blacks who reported an African country of birth; "African American" refers to all other Blacks. Cases with unknown race are excluded.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* Among Females by Race/Ethnicity and Year of Diagnosis, 1990-2001



\* HIV or AIDS at first diagnosis


† "African-born" refers to Blacks who reported an African country of birth; "African American" refers to all other Blacks. Cases with unknown race are excluded.

Data Source: Minnesota HIV/AIDS Surveillance System


HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* Diagnosed in Year 2001 by Gender and Race/Ethnicity

Males† (n = 209)



Females† (n = 71)



n = Number of persons    Amer Ind = American Indian  
 Afr Amer = African American (Black, not African-born persons)  
 Afr born = African-born (Black, African-born persons)

\* HIV or AIDS at first diagnosis

† 2 males and 1 female had missing race and were excluded.

# Age

## Average Age at HIV Diagnosis Among Males: Three-Year Averages

Race/Ethnicity	Average age in years (Number of cases)		
	1989-1991	1994-1996	1999-2001
White	33 (914)	35 (506)	38 (347)
Black			
African American	33 (212)	34 (186)	35 (134)
African-born	30 (6)	36 (20)	37 (50)
Hispanic	32 (54)	33 (44)	32 (65)
Asian	25 (5)	38 (8)	39 (12)
American Indian	29 (28)	29 (10)	35 (19)

Cases with unknown race or age were excluded.

## Average Age at HIV Diagnosis Among Females: Three-Year Averages

Race/Ethnicity	Average age in years (Number of cases)*		
	1989-1991	1994-1996	1999-2001
White	30 (88)	32 (68)	33 (58)
Black			
African American	29 (55)	30 (79)	32 (83)
African-born	21 (6)	31 (11)	29 (61)
Hispanic	34 (14)	34 (13)	28 (16)
Asian	--	--	32 (8)
American Indian	29 (16)	30 (12)	34 (9)

\* Average age not displayed for subgroups with less than 5 cases.


Cases with unknown race or age were excluded.

# Adolescents & Young Adults (Ages 13-24)\*

*\* Case numbers are too small to present meaningful data separately for adolescents and young adults.*



# New HIV Infections\* Among Adolescents and Young Adults† by Year of Diagnosis, 1990-2001



\* HIV or AIDS at first diagnosis


† Adolescents defined as 13-19 year-olds; Young Adults defined as 20-24 year-olds.

Data Source: Minnesota HIV/AIDS Surveillance System


HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* Among Adolescents and Young Adults† by Gender and Race/Ethnicity, 1999-2001 Combined

Males (n = 63)



Females (n = 51)



\* HIV or AIDS at first diagnosis

† Adolescents defined as 13-19 year-olds; Young Adults defined as 20-24 year-olds.


Data Source: Minnesota HIV/AIDS Surveillance System

n = Number of persons    Amer Ind = American Indian  
 Afr Amer = African American (Black, not African-born persons)  
 Afr born = African-born (Black, African-born persons)


HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* Among Adolescents and Young Adults† by Gender and Exposure Group, 1999-2001 Combined

Males (n = 63)



Females (n = 51)



\* HIV or AIDS at first diagnosis

† Adolescents defined as 13-19 year-olds; Young Adults defined as 20-24 year-olds.

Data Source: Minnesota HIV/AIDS Surveillance System

n = Number of persons

IDU = Injecting drug use


MSM = Men who have sex with men

Heterosex = Heterosexual contact

HIV/AIDS in Minnesota: Annual Review

# Mode of Exposure

# HIV Infections\* Among Males by Mode of Exposure and Year of Diagnosis, 1990-2001




MSM = Men who have sex with men    IDU = Injecting drug use    Heterosexual = Heterosexual contact

\* HIV or AIDS at first diagnosis

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# HIV Infections\* Among Males by Mode of Exposure and Year of Diagnosis, 1990-2001 (excluding MSM category)




MSM = Men who have sex with men    IDU = Injecting drug use    Heterosexual = Heterosexual contact

\* HIV or AIDS at first diagnosis

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# HIV Infections\* Among Females by Mode of Exposure and Year of Diagnosis, 1990-2001



IDU = Injecting drug use    Heterosexual = Heterosexual contact


\* HIV or AIDS at first diagnosis

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 combined

White Males (n = 347)




n = Number of persons    MSM = Men who have sex with men    IDU = Injecting drug use    Heterosex = Heterosexual contact

\* HIV or AIDS at first diagnosis



# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 combined

African American Males† (n = 134)



n = Number of persons    MSM = Men who have sex with men    IDU = Injecting drug use    Heterosex = Heterosexual contact

\* HIV or AIDS at first diagnosis


† Refers to Black, African American (not African-born) males.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

Hispanic Males (n = 65)



n = Number of persons    MSM = Men who have sex with men    IDU = Injecting drug use    Heterosex = Heterosexual contact


\* HIV or AIDS at first diagnosis

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

African-born Males<sup>†</sup> (n = 50)



n = Number of persons    MSM = Men who have sex with men    Heterosex = Heterosexual contact

\* HIV or AIDS at first diagnosis

<sup>†</sup> Refers to Black, African-born males.


Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

American Indian Males (n = 19)

CAUTION: Small number of cases – interpret carefully.




n = Number of persons    MSM = Men who have sex with men    IDU = Injecting drug use    Heterosex = Heterosexual contact

\* HIV or AIDS at first diagnosis

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

Asian Males (n = 12)

CAUTION: Small number of cases – interpret carefully.



n = Number of persons    MSM = Men who have sex with men    Heterosex = Heterosexual contact


\* HIV or AIDS at first diagnosis

Data Source: *Minnesota HIV/AIDS Surveillance System*

*HIV/AIDS in Minnesota: Annual Review*

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

African American Females†  
(n = 83)



n = Number of persons    IDU = Injecting drug use    Heterosex = Heterosexual contact    M w/ HIV = Mother with HIV, HIV risk

\* HIV or AIDS at first diagnosis


† Refers to Black, African American (not African-born) females.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

African-born Females† (n = 61)



n = Number of persons    M w/ HIV = Mother w/ HIV, HIV risk    Heterosex = Heterosexual contact

\* HIV or AIDS at first diagnosis


† Refers to Black, African-born females.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

White Females (n = 58)



n = Number of persons    IDU = Injecting drug use    Heterosex = Heterosexual contact

\* HIV or AIDS at first diagnosis

Data Source: *Minnesota HIV/AIDS Surveillance System*


*HIV/AIDS in Minnesota: Annual Review*



# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

Hispanic Females (n = 16)

CAUTION: Small number of cases – interpret carefully.



n = Number of persons    IDU = Injecting drug use    M w/ HIV = Mother w/ HIV, HIV risk    Heterosex = Heterosexual contact

\* HIV or AIDS at first diagnosis


Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

American Indian Females (n = 9)

CAUTION: Small number of cases – interpret carefully.



n = Number of persons    IDU = Injecting drug use    Heterosex = Heterosexual contact

\* HIV or AIDS at first diagnosis


Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# New HIV Infections\* by Mode of Exposure Diagnosis Years 1999-2001 Combined

Asian Females (n = 8)

CAUTION: Small number of cases – interpret carefully.



n = Number of persons    IDU = Injecting drug use    Heterosex = Heterosexual contact


\* HIV or AIDS at first diagnosis

Data Source: *Minnesota HIV/AIDS Surveillance System*

*HIV/AIDS in Minnesota: Annual Review*

# Emerging Trend

# New HIV Infections\* Among African-Born Persons† by Year of Diagnosis, 1990-2001



\* HIV or AIDS at first diagnosis

† Includes 4 non-Black, African-born individuals.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

## Companion Text for the Slide Set: *Minnesota HIV Surveillance Report, 2001*

### INTRODUCTION

#### Overview

The *Minnesota HIV Surveillance Report, 2001* describes the occurrence of reported HIV infections in Minnesota by person, place, and time through December 31, 2001. Such data provide information about where and among whom HIV transmission is likely occurring. This knowledge can in turn be used to help educate, target prevention efforts, plan for services, and develop policy.

#### Data Source

The data in this report are based on confidential case reports collected through the Minnesota Department of Health (MDH) HIV/AIDS Surveillance System. In Minnesota, laboratory-confirmed infections of human immunodeficiency virus (HIV) are monitored by the MDH through this active and passive surveillance system. State law (Minnesota Rule 4605.7040) requires both physicians and laboratories to report all cases of HIV infection (HIV or AIDS) directly to the MDH (passive surveillance). Additionally, regular contact is maintained with several clinical sites to help ensure completeness of reporting (active surveillance).

Data in this report include cases diagnosed with HIV as of December 31, 2001 and reported to the MDH as of April 2002. All data are displayed by earliest date of HIV diagnosis. Refer to the *HIV Surveillance Technical Notes* for a more detailed description of data inclusions and exclusions.

#### Data Limitations

Factors that impact the completeness and accuracy of the available surveillance data on HIV/AIDS include the level of screening and compliance with case reporting. Thus, any changes in numbers of infections may be due to one of these factors, or due to actual changes in HIV/AIDS occurrence.

The data presented in this report are not adjusted for reporting delays. Thus, the case number presented for the most recent reporting year can be viewed as a minimum and will likely increase in the future as further case reports are received. Changes in past years' totals are updated in every new annual surveillance report.

[Go to > Top of page](#)

### HIV/AIDS in the UNITED STATES

Compared with the rest of the nation, Minnesota is considered to be a low to moderate HIV/AIDS incidence state. In 1999, state-specific AIDS rates ranged from 1.1 per 100,000 persons in North Dakota to 42.3 per 100,000 persons in New York. Minnesota had the 12th lowest AIDS rate (4.0 AIDS cases reported per 100,000 persons). Compared with states in the Midwest region, Minnesota had one of the higher AIDS rates. State-specific HIV rates cannot be compared nationally because some states have not yet instituted HIV case surveillance. The states that have HIV case surveillance are at various stages of implementation.

### HIV/AIDS IN MINNESOTA

#### MDH HIV/AIDS Surveillance: Cumulative cases

AIDS has been tracked in Minnesota since 1982. In 1985, AIDS officially became a reportable disease to state and territorial health departments nationwide. Also in 1985, when the Food and Drug Administration approved the first diagnostic test for HIV, Minnesota became the first state to make HIV infection a reportable condition. As of December 31, 2001, a cumulative total of 6,661 cases of HIV infection have been reported among Minnesota residents. (1) This includes 3,854 AIDS cases and 2,807 HIV, non-AIDS cases. Of these 6,661 HIV/AIDS cases, 2,332 are known to be deceased through correspondence with the reporting source, other health departments, reviews of death certificates and obituaries, active surveillance, and matches with the National Death Index.

### **Overview of HIV/AIDS in Minnesota, 1990-2001**

The annual number of new AIDS cases increased steadily from the beginning of the epidemic to the early 1990s, reaching a peak of 370 cases in 1992. Beginning in 1996, both the number of newly diagnosed AIDS cases and the number of deaths among AIDS cases declined sharply, primarily due to the success of new antiretroviral therapies including protease inhibitors. These treatments do not cure, but can delay progression to AIDS among persons with HIV (non-AIDS) infection and improve survival among those with AIDS. Thus the declines slowed during the late 1990s and the numbers have become relatively stable the past few years. The number of newly diagnosed HIV (non-AIDS) cases has remained fairly constant since the mid 1990s at just under 200 cases per year. Furthermore, the number of prevalent HIV/AIDS cases has continued to increase over time. An estimated 4,331 persons with HIV/AIDS are assumed to be living in Minnesota as of December 31, 2001. (2)

[Go to > Top of page](#)

### **NEW HIV INFECTIONS IN MINNESOTA**

In this report, the term "new HIV infections" refers to HIV-infected Minnesota residents who were diagnosed in a particular calendar year and reported to the MDH. This includes persons whose first diagnosis of HIV infection is AIDS (AIDS at first diagnosis). HIV infection data are displayed by earliest known date of HIV diagnosis.

#### **New HIV Infections by Geography**

Historically, about 90% of new HIV infections diagnosed in Minnesota have occurred in Minneapolis, St. Paul and the surrounding seven-county metropolitan area. This has not changed over time. Although HIV infection is more common in communities with higher population densities and greater poverty, HIV or AIDS has been diagnosed in over 80% of counties in Minnesota.

#### **New HIV Infections by Gender**

Since the beginning of the epidemic, males have accounted for a majority of new HIV infections diagnosed per year. However, the number and the proportion of cases among females have increased over time. In 1990, males accounted for 90% of new HIV infections. In 2001, 75% of new infections occurred among males and 25% among females.

#### **New HIV Infections by Race/Ethnicity (3)**

Trends in the annual number of new HIV infections diagnosed among males differ by racial/ethnic group. New cases among White males drove the epidemic in the 1980s and early 1990s. Although Whites still account for the largest number of new infections among males, this number has generally been decreasing since 1991. A recent exception to this trend occurred between 2000 and 2001 when 130 cases were diagnosed among White males in 2001 compared to only 93 cases in the previous year. Whether this is an anomaly or the start of a trend remains to be seen.

In contrast to the overall large decline in the annual number of cases among White males, the decline among African American males was more gradual. The annual number of cases for African American males peaked in 1992 at 82 and gradually decreased to 34 in 2001.

The numbers of new cases in all other racial/ethnic groups during this same time remained stable or increased. Increases in the annual number of HIV infections diagnosed among Hispanic and African-born males, in particular, have been recorded since the late 1990s. The proportion of new HIV infections diagnosed among men of color as a whole has been increasing over time.

Similarly, trends in the annual number of HIV infections diagnosed among females differ by racial/ethnic group. In the beginning of the epidemic, White women accounted for a majority of newly diagnosed cases among females. Since 1991, the number of new infections among women of color has exceeded the number among White women. From 1990 to 2001, the annual number of new infections diagnosed nearly doubled among African American females (21 cases in 2001) and increased nine-fold among African-born females in just the past 5 years (26 cases in 2001). In fact among females, the only increase in the annual number of HIV infections between 2000 and 2001 occurred among African-born females. The annual number of new infections diagnosed among Hispanic, American Indian, and Asian females continues to be quite small (fewer than 10 cases per year for each of these groups).

The most recent data illustrate that men and women of color are disproportionately affected by HIV/AIDS. Whites make up approximately 88% of the male population in Minnesota and 63% of the new HIV infections diagnosed among men in 2001. Men of color make up approximately 12% of the male population and 37% of the new infections diagnosed among men in 2001. Similarly for females, Whites make up approximately 89% of the female population and 21% of new infections among women in 2001 whereas women of color make up approximately 11% of the female population and 79% of the new infections among women. (4)

Please note that race is not considered a biological reason for disparities in the occurrence of HIV experienced by persons of color. Race, however, can be considered a marker for other personal and social characteristics that put a person at greater risk for HIV exposure. These characteristics may include, but are not limited to, lower socioeconomic status, less education, and greater prevalence of drug use.

[Go to > Top of page](#)

### **Average Age at HIV Diagnosis, Three-year Averages**

In recent years, Hispanic, African American, and American Indian males were slightly younger (approximate age = 34 years) than White, African-born, and Asian males (approximate age = 38 years) at the time of HIV diagnosis. During the past ten years, the average age at HIV diagnosis has been approximately 30 years for females, for all racial/ethnic groups. Age at HIV diagnosis can be used as a proxy for age at HIV *infection*. However, due to differences in testing behavior (e.g. variable lengths of time between HIV infection and diagnosis) across time and between sociodemographic groups, comparisons of average age at diagnosis are difficult to interpret.

### **New HIV Infections among Adolescents and Young Adults (5), 1990-2001**

Many people are infected with HIV for years before they actually seek testing and become aware of their HIV status. This phenomenon especially affects the observed case counts for younger age groups. And as a result, the reported number of HIV infections among youth (5) (with few or no reports of AIDS at first diagnosis) is likely to underestimate the *true* number of new infections occurring in the population more than the reported number of cases in older age groups does.

In 1990, 9% of new HIV infections reported to the MDH were among youth. In 2001 this percentage was 11%. Among young men, the number of new HIV diagnoses peaked in 1992 at 43 cases and then declined through the mid 1990s to a low of 14 cases in 1997. Since 1997 the annual number of cases diagnosed among young men increased steadily to 28 in 2000, but then dropped to 16 cases in 2001.

Unlike young men, the annual number of new HIV infections diagnosed among young women has remained relatively consistent over time. For example, 16 cases of HIV infection were diagnosed among young women in 1992 and 16 cases in 2001. Females accounted for 50% of new HIV infections diagnosed among adolescents and young adults in 2001. In contrast, adult females (25 years of age or older) only accounted for 22% of all adult cases.

Similar to the adult HIV/AIDS epidemic, persons of color account for a disproportionate number of new HIV infections among adolescents and young adults. Among young men, Whites accounted for 36% of new HIV infections diagnosed between 1999 and 2001, African Americans account for 32% and Hispanics 21% of the cases. Among young women, African Americans accounted for 34%, Whites 29%, and African-born 25% of the new infections diagnosed during the same time period.

Men having sex with men (MSM) is the predominant mode of HIV exposure among adolescent and young adult males, accounting for 76% of the new HIV infections diagnosed between 1999 and 2001. Heterosexual contact and the joint risk of MSM and injecting drug use (IDU) each accounted for 2% of the cases. HIV exposure risk was not specified for 22% of the young male cases.

Heterosexual contact accounts for 45% of new HIV infections diagnosed among adolescent and young adult females between 1999 and 2001. IDU accounts for 6% of the cases. The remaining 49% of the young females do not have a risk specified.

Some hypotheses regarding the classification of males and females with unspecified risk are discussed in the next section.

### **New HIV Infections by Mode of Exposure**



Since the beginning, men have driven the HIV/AIDS epidemic in Minnesota and male-to-male sex has been the predominant mode of exposure reported. Though still the majority, both the number and proportion of new HIV infections attributed to MSM have been decreasing since 1991. On a much smaller scale, the numbers of male cases attributed to IDU and MSM/IDU also have been decreasing over the past decade, while the number of cases attributed to heterosexual contact has been increasing.

The number of cases without a specified risk has also been increasing. Throughout the epidemic, heterosexual contact has been the predominant mode of HIV exposure reported among females. IDU is the second most common mode of transmission making up 4% of cases among women in 2001. Unspecified risk has been designated for a growing percentage of cases for the past several years. In 1996, 13% of women diagnosed with HIV infection did not have a specified mode of transmission. This percentage grew to 58% in 2001. A portion of these cases represents women who have not yet been interviewed and, thus, some of these women will later have an identified mode of transmission. According to a study conducted by the Centers for Disease Control and Prevention (CDC) (6), it is likely that at least 80% of women with unspecified risk acquired HIV through heterosexual contact. Heterosexual contact as a mode of HIV transmission is currently only assigned to a female case if she knows that a male sexual partner of hers was HIV-infected or at increased risk for HIV (see *HIV Surveillance Technical Notes* for further details).

The proportion of cases attributable to a certain mode of exposure differs not only by gender, but also by race. Of the new HIV infections diagnosed among males between 1999 and 2001, MSM or MSM/IDU accounted for 81% of cases among White males, 59% of cases among Hispanic males, 50% of cases among African American males, and 4% of cases among African-born males. The latter three also had the highest proportions of cases with unspecified risk (28%, 35%, and 86%, respectively). It is hypothesized that due, in part, to social stigma many of the cases with unspecified risk were unclassified MSM cases. This may not hold as true for African-born cases given that heterosexual contact and contaminated medical equipment have been established modes of HIV exposure in their countries of origin. IDU or MSM/IDU was reported as a risk in 16% of male African American cases diagnosed during 1999-2001, but no more than 5% among other racial/ethnic groups. The number of cases among Asian and American Indian men during the years 1999-2001 were insufficient to make generalizations regarding risk (less than 20 cases in each group).

Heterosexual contact with a partner who has or is at increased risk for HIV infection accounted for 50% of cases among White females during 1999-2001, 46% of cases among African American females, and 18% of cases among African-born females. More than 41% of cases in each of these groups had no specified risk. IDU accounted for 9% of cases among Whites, 6% among African Americans, and 0% among African-born. The number of cases among Hispanic, Asian, and American Indian women during the years 1999-2001 were insufficient to make generalizations regarding risk (16 or less cases in each group).

### **Emerging Trend: New HIV Infections among African-born Persons**

The number of new HIV infections diagnosed among African-born persons in Minnesota has been steadily increasing from 7 cases in 1990 to 46 cases in 2001. During this time there was also a significant increase in African immigration to Minnesota. Among new HIV infections diagnosed in 2001, 16% were among African-born persons. Despite the absence of an accurate estimate of the number of African-born persons living in Minnesota, it is fair to speculate that they make up less than 1% of the total Minnesota population and are, therefore, disproportionately affected by HIV. (7)

(Last Revised: 4/12/2002)

---

(1) This number includes persons who reported Minnesota as their state of residence at the time of their HIV and/or AIDS diagnosis. It also includes persons who may have been diagnosed in a state that does not have HIV reporting and who subsequently moved to Minnesota and were reported here. HIV-infected persons currently residing in Minnesota, but who resided in another HIV-reporting state at the time of diagnosis are excluded.

(2) This number includes persons whose most recently reported state of residence was Minnesota, regardless of residence at time of diagnosis.

(3) Black race was broken down into African-born and African American (Black, not African-born). The numbers exclude 49 persons arriving through the HIV-Positive Refugee Resettlement Program.

(4) Population estimates based on U.S. Census 2000 data.

(5) In this report, adolescents are defined as 13-19 year-olds and young adults as 20-24 year-olds; these two groups are jointly referred to as "youth." Analyses are performed for adolescents and young adults combined because case numbers are too small to present meaningful data separately for each.

(6) MMWR 2001; 50(RR-6):31-40.

(7) Based on U.S. Census 2000 data, the U.S. Census Bureau estimates between 20,424 and 35,188 African-born persons are living in Minnesota out of a total population of 4,919,479. Because there are many reasons African-born persons may not be included in the census count (e.g. difficulties with verbal or written English), even 35,188 is likely an underestimate of the actual size of the African-born population living in Minnesota.

[Go to > Top of page](#)

For questions about the STD and HIV Section, please contact us at (612) 676-5414.

If you have questions or comments about this page, contact [idepcweb@health.state.mn.us](mailto:idepcweb@health.state.mn.us) or call 612-676-5414 (TTY: 612-676-5653) for the MDH [Infectious Disease Epidemiology, Prevention and Control Division](#).

[MDH HOME](#) | [ABOUT US](#) | [LIBRARY](#) | [NORTH STAR](#)

MAIN CATEGORIES: [Health Data & Statistics](#) | [Diseases & Conditions](#) | [Terrorism & Natural Disasters](#) | [Facilities & Professions](#) | [Health Care & Coverage](#) | [Minnesota's Public Health System](#) | [People & Environment](#) | [Policy, Economics & Legislation](#)

[Comments and Questions](#) | [Phone Number, Address and Directions](#) | [Privacy Statement and Disclaimer](#)

Updated Friday, 15-Apr-05 10:25:10

## HIV Surveillance Technical Notes

### Surveillance of HIV/AIDS

The Minnesota Department of Health (MDH) collects case reports of HIV infection and AIDS diagnoses through a passive and active HIV/AIDS surveillance system. Passive surveillance relies on physicians and laboratories to report new cases of HIV infection or AIDS directly to the MDH in compliance with state law (1). Active surveillance conducted by MDH staff involves routine visits and correspondence with select facilities to ensure completeness of reporting and accuracy of the data.

Factors that impact the completeness and accuracy of HIV/AIDS surveillance data include: compliance with case reporting, timeliness of case reporting, test-seeking behaviors of HIV-infected individuals, and the availability and targeting of HIV testing services. Certain events have also impacted trends in HIV/AIDS surveillance data. For example changes over time in the surveillance case definition (most notably the 1993 expansion of the case definition for adults and adolescents (2)) have resulted in artificial jumps in AIDS case counts at the time the new definition went into effect or in the preceding year because changes in case definition allowed for retrospective diagnoses.

### New HIV Infections

New HIV infections refer to persons who are diagnosed with HIV infection and newly reported to the MDH. This includes case-patients that meet the CDC surveillance definition for AIDS at the time they are initially diagnosed with HIV infection (AIDS at first diagnosis). Cases of new HIV infection are displayed by year of earliest HIV diagnosis. The number of new HIV infections in Minnesota includes only persons who were first reported with HIV infection while residents of Minnesota. Persons moving to Minnesota already infected with HIV are excluded if they were previously reported in another state.

### Vital Status of HIV/AIDS Cases

Persons are assumed alive unless the MDH has knowledge of their death. Persons are assumed residing in Minnesota if their most recently reported state of residence was Minnesota and the MDH has not received notice of relocation outside of the state. Vital status information is updated by monthly visits to select reporting facilities, correspondence with other health departments, daily obituary reviews in local newspapers, annual death certificate reviews, and periodic matches with the National Death Index. "AIDS deaths" refers to all deaths among AIDS cases regardless of the cause.

### Place of Residence for HIV/AIDS Cases

Persons are assumed to be residing in Minnesota if their most recently reported state of residence was Minnesota and the MDH has not received notice of relocation outside of the state. Likewise, a person's county or city of residence is assumed to be the most recently reported value unless the MDH is otherwise notified. Residence information is updated through standard case reporting, monthly visits to select reporting facilities and/or correspondence with other state health departments. Persons diagnosed with HIV infection while imprisoned in a state correctional facility are included in the data presented unless otherwise noted (federal and private prisoners are excluded). Residential relocation, including release from state prison, is difficult to track and therefore data presented by *current* residence must be interpreted in this light. Data on residence at *time of diagnosis* are considered more accurate, limited only by the accuracy of self-reported residence location.

### Data Tabulation and Presentation

The data displayed are not adjusted to correct for reporting delays, case definition changes, or other factors.

MDH surveillance reports published before 2000 displayed data by year of report, the data in these documents are displayed by earliest date of HIV diagnosis. The report date is a function of reporting practices and may be months or years after the date of diagnosis and the date of infection. The date of diagnosis is temporally closer to the date of infection. Displaying data by year of diagnosis more closely approximates when infection occurred. Readers should bear in mind that diagnosis date is also an approximation for infection date. Many years may pass between time of infection and diagnosis; the incubation period (3) for HIV is around 10 years. It should also be noted that because of delays in reporting, the annual number of cases reportedly diagnosed in recent years is slightly lower than actual. This discrepancy corrects itself over time. The number of cases diagnosed within a calendar year changes relatively little after two years have passed.

Unless otherwise noted, data analyses exclude persons diagnosed in federal or private correctional facilities (inmates generally are not Minnesota residents before incarceration and do not stay in Minnesota upon their release), infants with unknown or negative HIV status who were born to HIV positive mothers, and HIV-infected refugees who resettled in Minnesota as part of the HIV-Positive Refugee Resettlement Program.

## Mode of Exposure Hierarchy

All state and city HIV/AIDS surveillance systems funded by the Centers for Disease Control and Prevention use a standardized hierarchy of mode of exposure categories. HIV and AIDS cases with more than one reported mode of exposure to HIV are classified in the exposure category listed first in the hierarchy. In this way, each case is counted as having only one mode of exposure. The only exception to this rule is the joint risk of male-to-male sex (MSM) and injection drug use (IDU), which makes up a separate exposure category in the hierarchy. The following is a list of the hierarchy for adolescent/adult HIV/AIDS cases:

- (1) MSM
- (2) IDU
- (3) MSM/IDU
- (4) Hemophilia patient
- (5) Heterosexual contact
- (6) Receipt of blood transfusion or tissue/organ transplant
- (7) Other (e.g. needle stick in a health care setting)
- (8) Risk not specified.

The following is the list of the hierarchy for pediatric HIV/AIDS cases:

- (1) Hemophilia patient
- (2) Mother with HIV or HIV risk
- (3) Receipt of blood transfusion or tissue/organ transplant
- (4) Other
- (5) Risk not specified.

Heterosexual contact is only designated if a male or female can report specific heterosexual contact with a partner who has, or is at increased risk for, HIV infection (e.g. an injection drug user). For females this includes heterosexual contact with a bisexual male (mainly due to the elevated prevalence of HIV infection among men who have sex with men).

"Risk not specified" refers to cases with no reported history of exposure to HIV through any of the routes listed in the hierarchy of exposure categories. These cases include persons who have not yet been interviewed by MDH staff; persons whose exposure history is incomplete because they died, declined to be interviewed, or were lost to follow-up; and persons who were interviewed or for whom follow-up information was available but no exposure was identified/acknowledged.

The growing number of cases with unspecified risk in recent years is, in part, artificial and due to interviews that have not yet been completed. In time, a number of these will be assigned a mode of exposure category. However, part of the observed increase is real. As stated above, a person must have intimate knowledge about his/her partner to meet the criteria for heterosexual mode of exposure. Often cases will not be certain about their partners' HIV status or risk. Additionally, the perception of social stigma presumably decreases the likelihood that a person will acknowledge certain risk behaviors, particularly male-to-male sex or injection drug use. Thus, if the *true* numbers of cases due to heterosexual contact, MSM, and/or IDU increase, a larger number of cases without a specified risk would be expected.

A recent study by the Centers for Disease Control and Prevention used statistical methods to redistribute risk among female HIV/AIDS cases with unspecified risk (4). The results are helpful but are based on national data which are not necessarily applicable to the state or local level. Speculation regarding the distribution of risk behaviors among those with unspecified risk is difficult, especially in men, for whom even a national study is not available.

## Re-distribution of Mode of Exposure

In 2004 the Minnesota Department of Health began estimating mode of exposure for cases with unspecified risk in its annual summary slides. Estimation was done by using the risk distribution for cases reported between 2002 and 2004 with known risk by race and gender and applying it to those with unspecified risk of the same race and gender. There were two exceptions to this method, African-born cases and Asian/Pacific Islander women. For both African-born and Asian/Pacific Islander women a breakdown of 95% heterosexual risk and 5% other risk was used. For African-born males a breakdown of 5% male-to-male sex, 90% heterosexual risk, and 5% other risk was used. These percentages are based on epidemiological literature and/or community experience.

Below is an example of how the process worked for white, African American and African-born females:

Reported Female cases 2002 - 2004					
Race/Risk	Heterosexual n (%†)	IDU n (%†)	Other (5) n (%†)	Unspecified n	Total N
White	25 (86)	4 (14)	0 (0)	14	43
African-American	23 (79)	3 (10)	3 (10)	33	62
African-born	13 (81)	0 (0)	3 (19)	89	105

† Percent of those with know risk.

Female Cases for 2002 - 2004 with Estimated risk:					
Race/Risk	Heterosexual	IDU	Other	Unspecified	Total N
White	$(.86*14) + 25 = 37$	$(.14*14) + 4 = 6$	0	0	43
African-American	$(.79*33) + 23 = 49$	$(.1*33) + 3 = 6$	$(.1*33) + 3 = 6$	0	62
African-born‡	$(.95*89) + 13 = 98$	0	$(.05*89) + 3 = 7$	0	105

‡ Used a distribution of 95% heterosexual and 5% other.

## Definitions Related to Race/Ethnicity

When data are stratified by race, Black race is broken down into African-born and African American (not African-born) based on reported country of birth.

The terms "persons of color" and "non-Whites" refer to all race/ethnicity categories other than White (Black, Hispanic, American Indian, and Asian/Pacific Islander).

## Interstate De-Duplication Project (IDEP)

In 2004, the Minnesota Department of Health (MDH) participated in IDEP. IDEP is a CDC project aimed at eliminating duplicate reports of HIV and AIDS cases among states. Each case of HIV and AIDS is assigned to the state (or states when the diagnosis of HIV and AIDS occurs in two different states) where a person was first diagnosed. The first round looked at cases reported through December 31, 2001. Through this project, MDH identified 164 cases of HIV infection (including AIDS at first report) and 55 AIDS cases whose first diagnosis was not in Minnesota. These cases were previously considered as diagnosed in Minnesota and were counted in the cumulative number of cases diagnosed in Minnesota. As such, the change of "ownership" (where the case was diagnosed) has reduced both cumulative and yearly totals for Minnesota. Additionally, MDH also identified 250 cases that no longer live in Minnesota.

The results of IDEP are particularly noticeable in the total number of persons living with HIV/AIDS in Minnesota, which increased from 4,895 to 5,002, a gain of only 107 cases, instead of 292 (new infections minus deaths plus (difference between people moving in and people moving out)) between 2003 and 2004.

---

- (1) [Minnesota Rule 4605.7040 \(return to text\)](#)
- (2) [MMWR 1992; 41\[no.RR-17\]: 1-19 \(return to text\)](#)
- (3) Incubation period is the time between initial infection with the virus and the development of disease symptoms. [\(return to text\)](#)
- (4) [MMWR 2001; 50\(RR-6\): 31-40 \(return to text\)](#)
- (5) Other includes Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk [\(return to text\)](#)

If you have questions or comments about this page, contact [idepcweb@health.state.mn.us](mailto:idepcweb@health.state.mn.us) or call 612-676-5414 (TTY: 612-676-5653) for the MDH [Infectious Disease Epidemiology, Prevention and Control Division](#).

[MDH HOME](#) | [ABOUT US](#) | [LIBRARY](#) | [NORTH STAR](#)

MAIN CATEGORIES: [Health Data & Statistics](#) | [Diseases & Conditions](#) | [Terrorism & Natural Disasters](#) | [Facilities & Professions](#) | [Health Care & Coverage](#) | [Minnesota's Public Health System](#) | [People & Environment](#) | [Policy, Economics & Legislation](#)

[Comments and Questions](#) | [Phone Number, Address and Directions](#) | [Privacy Statement and Disclaimer](#)

Updated Monday, 11-Apr-05 10:49:41

**Table 1. Number of New Cases and Rates (per 100,000 persons) of  
HIV Infection, HIV (non-AIDS), and AIDS\* -- Minnesota, 1997-2001**

<b>Disease</b>	<b>1997 Cases</b>	<b>1997 Rate</b>	<b>1998 Cases</b>	<b>1998 Rate</b>	<b>1999 Cases</b>	<b>1999 Rate</b>	<b>2000 Cases</b>	<b>2000 Rate</b>	<b>2001 Cases</b>	<b>2001 Rate</b>
HIV Infection	282	5.9	298	6.2	309	6.4	275	5.6	283	5.8
HIV (non-AIDS)	192	4.0	191	4.0	202	4.2	177	3.6	197	4.0
AIDS	198	4.2	198	4.1	162	3.3	162	3.3	124	2.5

\* HIV Infection = New cases of HIV infection (both HIV (non-AIDS) and AIDS at first diagnosis) diagnosed within a given calendar year.  
HIV (non-AIDS) = New cases of HIV infection (excluding AIDS at first diagnosis) diagnosed within a given calendar year. AIDS = All new cases of AIDS diagnosed within a given calendar year.

**Please Note:** The sum of HIV (non-AIDS) cases and AIDS cases will be greater than the number of cases of HIV Infection in a given year. The difference occurs because, unlike the HIV Infection category, the AIDS category also includes persons who progressed from an HIV diagnosis to an AIDS diagnosis (see above definitions).

Numbers and rates exclude federal and private prisoners and refugees in the HIV-Positive Refugee Resettlement Program.

<b>Table 2. Number of Cases and Rates (per 100,000 persons) of HIV Infection by Residence, Age, &amp; Gender* -- Minnesota, 2001</b>			
<b>Group</b>	<b>HIV Infection Cases %</b>		<b>HIV Infection Rate</b>
<b>Residence**</b>			
Minneapolis	127	45%	33.2
St. Paul	41	15%	14.3
Suburban	86	30%	4.4
Greater Minnesota	28	10%	1.2
<i>Total</i>	<i>282</i>	<i>100%</i>	<i>5.7</i>
<b>Age</b>			
<13 yrs	1	0%	0.1
13-19 yrs	6	2%	1.1
20-24 yrs	26	9%	8.1
25-29 yrs	42	15%	13.1
30-34 yrs	56	20%	15.9
35-39 yrs	62	22%	15.0
40-44 yrs	36	13%	8.7
45-49 yrs	23	8%	6.3
50-54 yrs	17	6%	5.6
55+ yrs	14	5%	1.4
<i>Total</i>	<i>283</i>	<i>100%</i>	<i>5.8</i>
<b>Gender</b>			
Male	211	75%	8.7
Female	72	25%	2.9
<i>Total</i>	<i>283</i>	<i>100%</i>	<i>5.8</i>

\* HIV Infection includes all new cases of HIV infection (both HIV (non-AIDS) and AIDS at first diagnosis) among Minnesota residents in 2001.

\*\* Residence information missing for 1 case of HIV infection in 2001.

Suburban = Seven-county metropolitan area except Minneapolis & St. Paul (Anoka, Carver, Dakota, Hennepin (except Minneapolis), Ramsey (except St. Paul), Scott, and Washington counties). Greater Minnesota = Remaining 80 counties outside of the seven-county metropolitan area.

Numbers and rates exclude federal and private prisoners and refugees in the HIV-Positive Refugee Resettlement Program. State prisoners are included (3 new diagnoses in 2001). Rates calculated using U.S. Census 2000 data.



<b>Table 3. Number of Cases and Rates (per 100,000 persons) of HIV Infection by Race/Ethnicity &amp; Mode of Exposure* -- Minnesota, 2001</b>									
	<b>Males</b>			<b>Females</b>			<b>Total</b>		
<b>Group</b>	<b>Cases</b>	<b>%</b>	<b>Rate</b>	<b>Cases</b>	<b>%</b>	<b>Rate</b>	<b>Cases</b>	<b>%</b>	<b>Rate</b>
<b>Race/Ethnicity</b>									
White, non-Hispanic	130	62%	#	15	21%	#	145	51%	3.4
Black**, African-American	34	16%	#	21	29%	#	55	19%	27.1
Black**, African-born	19	9%	X	26	36%	X	45	16%	X
Hispanic	16	8%	#	5	7%	#	21	7%	14.6
American Indian	7	3%	#	2	3%	#	9	3%	11.1
Asian/PI	3	1%	#	2	3%	#	5	2%	3.0
Unknown	2	1%	X	1	1%	X	3	1%	X
<i>Total</i>	211	100%	8.7	72	100%	2.9	283	100%	5.8
<b>Mode of Exposure</b>									
MSM	132	63%	X	--	--	X	132	47%	X
IDU	2	1%	X	4	6%	X	6	2%	X
MSM/IDU	8	4%	X	--	--	X	8	3%	X
Heterosexual	7	3%	X	25	35%	X	32	11%	X
Other	0	0%	X	1	1%	X	1	0%	X
Unspecified	62	29%	X	42	58%	X	104	37%	X
<i>Total</i>	211	100%	8.7	72	100%	2.9	283	100%	5.8

\* HIV infection includes all new cases of HIV infection (both HIV (non-AIDS) and AIDS at first diagnosis) among Minnesota residents in 2001.

\*\* African-born Blacks are reported separately from other Blacks (born in the U.S. or elsewhere).

# U.S. Census 2000 data necessary to calculate race-specific rates (specifically a breakdown of the state population by "Race alone or in Combination with one or more races" by gender) have not yet been released for Minnesota. When these data become available this table will be updated.

X Accurate population estimates not available for calculation of rates.

Numbers exclude federal and private prisoners and refugees in the HIV-Positive Refugee Resettlement Program.

MSM = Men who have sex with men. IDU = Injecting drug use. Heterosexual = For males: heterosexual contact with a female known to be HIV+, an injecting drug user, or a blood product or organ transplant recipient. For females: heterosexual contact with a male known to be HIV+, bisexual, an injecting drug user, or a blood product or organ transplant recipient. Other = Perinatal HIV exposure. Unspecified = Cases who did not acknowledge any of the risks listed above or have not been/refused to be interviewed.

**Table 4. Number of Cases and Rates (per 100,000 persons) of HIV Infection by County of Residence\* -- Minnesota, 2001**

<b>County**</b>	<b>HIV Infection Cases</b>	<b>HIV Infection Rate***</b>
Aitkin	0	-
Anoka	15	5.0
Becker	0	-
Beltrami	0	-
Benton	0	-
Big Stone	0	-
Blue Earth	1	-
Brown	0	-
Carlton	1	-
Carver	0	-
Cass	0	-
Chippewa	0	-
Chisago	2	-
Clay	1	-
Clearwater	0	-
Cook	1	-
Cottonwood	0	-
Crow Wing	0	-
Dakota	9	2.5
Dodge	0	-
Douglas	0	-
Faribault	1	-
Fillmore	0	-
Freeborn	0	-
Goodhue	0	-
Grant	1	-
Hennepin	176	15.8
Houston	0	-
Hubbar	0	-
Isanti	0	-
Itasca	0	-
Jackson	0	-
Kanabec	0	-
Kandiyohi	0	-
Kittson	0	-
Koochiching	0	-
Lac Qui Parle	0	-
Lake	0	-
Lake of theWoods	0	-
Le Sueur	0	-
Lincoln	0	-
Lyon	1	-
McLeod	1	-
Mahnomen	0	-
Marshall	1	-
Martin	0	-
Meeker	0	-
Mille Lacs	0	-
Morrison	0	-
Mower	2	-

Murray	0	-
Nicollet	0	-
Nobles	3	-
Norman	0	-
Olmsted	2	-
Otter Tail	0	-
Pennington	0	-
Pine	0	-
Pipestone	0	-
Polk	0	-
Pope	0	-
Ramsey	46	9.0
Red Lake	0	-
Redwood	0	-
Renville	1	-
Rice	1	-
Rock	0	-
Rosseau	0	-
St. Louis	1	-
Scott	5	5.6
Sherburne	0	-
Sibley	0	-
Stearns	4	-
Steele	0	-
Stevens	1	-
Swift	0	-
Todd	0	-
Traverse	0	-
Wabasha	0	-
Wadena	0	-
Waseca	1	-
Washington	3	-
Watonwan	0	-
Wilkin	0	-
Winona	0	-
Wright	1	-
Yellow Medicine	0	-
<i>State Total**</i>	<i>282</i>	<i>5.8</i>

\* HIV infection includes all new cases of HIV infection (both HIV (non-AIDS) and AIDS at first diagnosis) among Minnesota residents in 2001. County of residence as reported at time of diagnosis.

\*\* Residence information missing for 1 case of HIV infection in 2001; Total rate is based on all cases in the state (n = 283).

\*\*\* Rates not calculated for counties with fewer than 5 cases.

Numbers and rates exclude federal and private prisoners and refugees in the HIV-Positive Refugee Resettlement Program. HIV infection was diagnosed among 3 state prisoners during 2001 (State correctional facilities are located in the following counties: Anoka, Carlton, Chisago, Goodhue, Pine, Rice, Scott, St. Louis, Stearns, and Washington).

Rates calculated using U.S. Census 2000 data.

# *HIV/AIDS Prevalence & Mortality Report, 2001*



**Minnesota Department of Health  
HIV/AIDS Surveillance System**

# Introduction (I)

- These three introduction slides provide a general context for the data used in this slide set. If you have questions about any of the slides please refer to the ***Companion Text to the Minnesota HIV/AIDS Prevalence & Mortality Report, 2001*** or ***HIV/AIDS Prevalence & Mortality Technical Notes***.
- This slide set displays estimates of the number of persons living with HIV/AIDS (**prevalence**) and mortality in Minnesota by person, place, and time.
- The slides rely on data from HIV/AIDS cases diagnosed through 2001 and reported to the Minnesota Department of Health (MDH) HIV/AIDS Surveillance System.

## Introduction (II)

- Data analyses exclude persons diagnosed in federal or private correctional facilities, but include state prisoners (n=101).
- Some limitations of surveillance data:
  - ◆ Data do not include HIV-infected persons who have not been tested for HIV
  - ◆ Data do not include persons whose positive test results have not been reported to the MDH
  - ◆ Case numbers for the most recent years may be undercounted due to delays in reporting


## Introduction (III)

- Persons are assumed to be alive unless the MDH has knowledge of their death.
- Persons whose most recently reported state of residence was Minnesota are assumed to be currently residing in Minnesota unless the MDH has knowledge of their relocation. Our ability to track changes of residence, including within the state, is limited.
- Vital status and current residence are updated through one or more of the following methods:
  - ◆ Standard case reporting
  - ◆ Correspondence with other health departments
  - ◆ Obituary reviews (daily)
  - ◆ Active surveillance (monthly)
  - ◆ Death certificate reviews (annually)
  - ◆ National Death Index match

# National Context



# Estimated Incidence of AIDS, Deaths, and Prevalence by Quarter-Year of Diagnosis/Death, United States, 1985-1999\*




\*Adjusted for reporting delays



# Overview of HIV/AIDS in Minnesota

# HIV/AIDS in Minnesota:

Number of new cases, prevalent cases, and deaths by year, 1990-2001



\* Deaths among AIDS cases, regardless of cause.

# Persons Living with HIV/AIDS in Minnesota


# Estimated Number of Persons Living with HIV/AIDS in Minnesota

- As of December 31, 2001, 4,331\* persons are assumed alive and living in Minnesota with HIV/AIDS
  - ◆ 2,574 living with HIV infection (non-AIDS)
  - ◆ 1,757 living with AIDS
- This number includes 375 persons who were first reported with HIV or AIDS elsewhere and subsequently moved to Minnesota
- This number excludes 418 persons who were first reported with HIV or AIDS in Minnesota and subsequently moved out of the state

*\* This number includes persons who reported Minnesota as their current state of residence, regardless of residence at time of diagnosis.*


# Place

# Living HIV/AIDS Cases by County of Residence, 2001




## Map of Metro Area: Living HIV/AIDS Cases by County of Residence, 2001

Number Living  
with HIV/AIDS




Total number (Metro only) = 3,811



\* Counties in which a state  
correctional facility is located.




# Persons Living with HIV/AIDS in Minnesota by Current Residence, 2001




Suburban = Seven-county metro area including Anoka, Carver, Dakota, Hennepin (except Minneapolis), Ramsey (except St. Paul), Scott, and Washington counties. Greater MN = All other Minnesota counties, outside the seven-county metro area.

# Gender and Race/Ethnicity

# Persons Living with HIV/AIDS in Minnesota by Gender, 2001




# Persons Living with HIV/AIDS in Minnesota by Gender and Race/Ethnicity\*, 2001




\* "African-born" refers to Blacks who reported an African country of birth; "African American" refers to all other Blacks. Persons with unknown race were omitted (n=31).

# Persons Living with HIV/AIDS in Minnesota by Gender and Race/Ethnicity, 2001

Males\* (n = 3,497)



Females\* (n = 803)



n = Number of persons    Amer Ind = American Indian  
 Afr Amer = African American (Black, not African-born persons)  
 Afr born = African-born (Black, African-born persons)


\* 26 males and 5 females had missing race and were excluded.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# Age

# Persons Living with HIV/AIDS in Minnesota by Age Group, 2001




# Mode of Exposure



# Males Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

White Males (n = 2,332)




n = Number of persons  
IDU = Injecting drug use

MSM = Men who have sex with men  
Heterosex = Heterosexual contact

Other = Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk

# Males Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

African American Males<sup>†</sup> (n = 724)



n = Number of persons

IDU = Injecting drug use

<sup>†</sup> Refers to Black, African American (not African-born) males.


MSM = Men who have sex with men

Heterosex = Heterosexual contact

Other = Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk

# Males Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

Hispanic Males (n = 217)




n = Number of persons  
IDU = Injecting drug use

MSM = Men who have sex with men  
Heterosex = Heterosexual contact

Other = Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk

# Males Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

African-born Males<sup>†</sup> (n = 138)



n = Number of persons

MSM = Men who have sex with men


IDU = Injecting drug use

Heterosex = Heterosexual contact

<sup>†</sup> Refers to Black, African-born males.

# Males Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

American Indian Males (n = 55)



n = Number of persons


MSM = Men who have sex with men

IDU = Injecting drug use

Heterosex = Heterosexual contact

# Males Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

Asian Males (n = 31)



n = Number of persons


MSM = Men who have sex with men

IDU = Injecting drug use

Heterosex = Heterosexual contact

# Females Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

African American Females<sup>†</sup> (n = 303)



n = Number of persons

IDU = Injecting drug use


<sup>†</sup> Refers to Black, African American (not African-born) females.

Other = Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk

Heterosex = Heterosexual contact

# Females Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

White Females (n = 281)




n = Number of persons  
IDU = Injecting drug use

Other = Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk  
Heterosex = Heterosexual contact



# Females Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

African-born Females<sup>†</sup> (n = 125)



n = Number of persons

Other = Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk

Heterosex = Heterosexual contact


<sup>†</sup> Refers to Black, African-born females.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

# Females Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

Hispanic Females (n = 44)




n = Number of persons  
IDU = Injecting drug use

Other = Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk  
Heterosex = Heterosexual contact

# Females Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

American Indian Females (n = 32)



n = Number of persons


IDU = Injecting drug use

Heterosex = Heterosexual contact

# Females Living with HIV/AIDS in Minnesota by Mode of Exposure, 2001

Asian Females (n = 18)

CAUTION: Small number of cases – interpret carefully.




n = Number of persons

Other = Hemophilia, transplant, transfusion, mother w/ HIV or HIV risk

Heterosex = Heterosexual contact


# Emerging Trend

# Foreign-Born Persons Living with HIV/AIDS in Minnesota, 1990-2001



# Mortality

## Reported Deaths\* among Male AIDS Cases in Minnesota, 1990-2001




\* Deaths among AIDS cases, regardless of cause.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review



# Reported Deaths\* among Female AIDS Cases in Minnesota, 1990-2001



\* Deaths among AIDS cases, regardless of cause.

Data Source: Minnesota HIV/AIDS Surveillance System

HIV/AIDS in Minnesota: Annual Review

## **Companion Text for the Slide Set: *Minnesota HIV/AIDS Prevalence & Mortality Report, 2001***

### **INTRODUCTION**

The *Minnesota HIV/AIDS Prevalence & Mortality Report, 2001* contains estimates of HIV/AIDS prevalence (the number of persons living with HIV or AIDS) and mortality in Minnesota. These estimates can be used to help educate, plan for HIV/AIDS services and develop policy.

#### **Data Source**

The data in this report are based on confidential case reports collected through the Minnesota Department of Health (MDH) HIV/AIDS Surveillance System. In Minnesota, laboratory-confirmed infections of human immunodeficiency virus (HIV) are monitored by the MDH through this active and passive surveillance system. State law (Minnesota Rule 4605.7040) requires both physicians and laboratories to report all cases of HIV infection (HIV or AIDS) directly to the MDH (passive surveillance). Additionally, regular contact is maintained with several clinical sites to ensure completeness of reporting (active surveillance).

#### **Data Limitations**

The prevalence estimate is calculated by totaling the number of HIV and AIDS cases diagnosed through December 31, 2001 who are not known to be deceased and whose most recently reported state of residence was Minnesota. It bears noting that persons who are HIV-infected but not yet tested are not included in this prevalence estimate. Migration (known HIV-infected persons moving in or out of the state) also affects the estimate. Refer to the *HIV/AIDS Prevalence & Mortality Technical Notes* for a more detailed description of data inclusions and exclusions.

Factors that impact the completeness and accuracy of the available surveillance data on HIV/AIDS include the level of screening and compliance with case reporting. Thus, any changes in numbers of infections may be due to one of these factors, or due to actual changes in HIV/AIDS occurrence.

### **PERSONS LIVING WITH HIV/AIDS IN THE UNITED STATES**

The Centers for Disease Control & Prevention (CDC) estimates that there are 800,000 to 900,000 people currently living with HIV/AIDS in the United States. The number of people specifically living with AIDS in the United States has been increasing in recent years: from approximately 290,400 in 1998 to approximately 339,000 in 2000. (1)

### **PERSONS LIVING WITH HIV/AIDS IN MINNESOTA**

#### **Overview of HIV/AIDS in Minnesota, 1990-2001**

The number of persons assumed to be living with HIV/AIDS in Minnesota has been steadily increasing over time. As of December 31, 2001, an estimated 4,331 persons with HIV/AIDS were residing in Minnesota, a 7% increase from 2000. While the number of new HIV (non-AIDS) cases has remained steady since the mid-1990s at just under 200 cases per year, both the number of newly diagnosed AIDS cases and the number of deaths among AIDS cases have been declining since 1996. The decreases are primarily due to the success of new treatments introduced in 1995 (protease inhibitors) and 1996 (highly active antiretroviral therapy, HAART). These treatments do not cure, but can delay progression to AIDS among persons with HIV (non-AIDS) infection and improve survival among those with AIDS. Thus, the declines slowed during the late 1990s and the numbers have become relatively stable the past few years.

#### **Living HIV/AIDS Cases, 2001**

Among the estimated 4,331 prevalent cases in Minnesota, 2,574 are diagnosed with HIV (non-AIDS) and 1,757 are diagnosed with AIDS. The majority (89%) of prevalent cases reside in the seven-county metropolitan area surrounding the Twin Cities of Minneapolis and St. Paul (Hennepin, Ramsey, Anoka, Dakota, Scott, and Washington counties). Although HIV infection is more common in communities with higher

population densities and greater poverty, HIV or AIDS has been diagnosed in over 80% of counties in Minnesota.

### **Gender & Race/Ethnicity**

Approximately 81% of prevalent HIV/AIDS cases are males. Broken down by race/ethnicity, 66% of male cases are White, 21% African American, 6% Hispanic, 4% African-born, 2% American Indian, and <1% Asian/Pacific Islander. In total, 36% of males living with HIV/AIDS are non-White whereas only 12% of the general male population is Non-White. Among female cases, the distribution is even more skewed toward women of color: 35% White, 38% African American, 16% African-born, 5% Hispanic, 4% American Indian, and 2% Asian/Pacific Islander. Thus, 65% of prevalent female HIV/AIDS cases are non-White whereas only 11% of the general female population in Minnesota is non-White.

Please note that race is not considered a biological reason for disparities related to HIV/AIDS experienced by persons of color. Race, however, can be considered a marker for other personal and social characteristics that put a person at greater risk for HIV exposure. These characteristics may include, but are not limited to, lower socioeconomic status, less education, and less access to health care.

### **Age**

Seventy-six percent (76%) of persons living with HIV/AIDS in 2001 are currently 35 years of age or older. Broken down into five-year age groups, 35-39 year olds make up the largest group (24% of cases), followed by 40-44 year olds (23%) and 45-49 year olds (15%).

### **Mode of Exposure**

The proportions of living cases attributable to particular modes of exposure differ among gender and race groups. While 88% of White males reported male-to-male sex (MSM or MSM/IDU) as a risk factor, only 54% of non-White males reported this mode of exposure. The difference in proportions is partly explained by the relatively large number of non-White males with unspecified risk, particularly among African American, African-born and Asian men. It is hypothesized that due, in part, to social stigma many of the cases with unspecified risk were unclassified MSM cases. This may not hold as true for African-born cases given that heterosexual contact and contaminated medical equipment have been established modes of HIV exposure in their countries of origin. The percent of male cases who identified IDU or MSM/IDU as a risk factor was particularly high for American Indians (28%) and African Americans (24%). The percentages among Hispanic and White males were 18% and 11%, respectively. Presently, no prevalent HIV/AIDS cases among African-born or Asian males identified IDU as a risk factor. Similar to the MSM category, IDU may be underreported due to social stigma.

Across all race/ethnicity groups, females most frequently report heterosexual contact as their mode of HIV exposure. IDU accounts for the second most common mode of exposure among American Indian (34%), African American (20%), White (19%), and Hispanic (16%) females. No cases among African-born females were attributed to IDU. African-born females living with HIV/AIDS had the largest percentage of cases attributed to unspecified risk: 77% compared to approximately 45% among the other female race/ethnicity groups. The number of prevalent HIV/AIDS cases among Asian females was too small ( $n = 18$ ) to make generalizations about risk. See the *HIV/AIDS Prevalence & Mortality Technical Notes* for a detailed discussion of mode of exposure categories.

### **Emerging Trend**

Between 1990 and 2001, the number of foreign-born persons living with HIV/AIDS in Minnesota increased substantially, especially among the African-born population. In 1990, 50 foreign-born persons were reported to be living with HIV/AIDS in Minnesota. By 2001, this number increased eight-fold to 450 persons. This trend illustrates the growing diversity of the infected population in Minnesota and the need for culturally appropriate HIV care services and prevention efforts.

### **HIV/AIDS MORTALITY IN MINNESOTA**

The number of deaths (2) among Minnesota AIDS cases decreased between 1995 and 1997 and remained relatively constant between 1997 and 2001. The largest declines in mortality were observed among White males in the mid 1990s. In recent years, the number of deaths among AIDS cases has been comparable between White and non-White males and between White and non-White females. Only 3 deaths among AIDS cases were reported in 2001 among women and 47 among men.

(Last Revised: 4/12/2002)

---

(1) *HIV/AIDS Surveillance Report* Vol. 13, No. 1: June 2001. Centers for Disease Control & Prevention.

(2) Includes all deaths, regardless of cause.

For questions about the STD and HIV Section, please contact us at (612) 676-5414.

If you have questions or comments about this page, contact [idepcweb@health.state.mn.us](mailto:idepcweb@health.state.mn.us) or call 612-676-5414 (TTY: 612-676-5653) for the MDH [Infectious Disease Epidemiology, Prevention and Control Division](#).

[MDH HOME](#) | [ABOUT US](#) | [LIBRARY](#) | [NORTH STAR](#)

MAIN CATEGORIES: [Health Data & Statistics](#) | [Diseases & Conditions](#) | [Terrorism & Natural Disasters](#) | [Facilities & Professions](#) | [Health Care & Coverage](#) | [Minnesota's Public Health System](#) | [People & Environment](#) | [Policy, Economics & Legislation](#)

[Comments and Questions](#) | [Phone Number, Address and Directions](#) | [Privacy Statement and Disclaimer](#)

Updated Friday, 15-Apr-05 10:28:32

## HIV/AIDS Prevalence and Mortality Technical Notes

### Surveillance of HIV/AIDS

The Minnesota Department of Health (MDH) collects case reports of HIV infection and AIDS diagnoses through a passive and active HIV/AIDS surveillance system. Passive surveillance relies on physicians and laboratories to report new cases of HIV infection or AIDS directly to the MDH in compliance with state law (1). Active surveillance conducted by MDH staff involves routine visits and correspondence with select facilities to ensure completeness of reporting and accuracy of the data.

Factors that impact the completeness and accuracy of HIV/AIDS surveillance data include: compliance with case reporting, timeliness of case reporting, test-seeking behaviors of HIV-infected individuals, and the availability and targeting of HIV testing services. Certain events have also impacted trends in HIV/AIDS surveillance data. For example changes over time in the surveillance case definition (most notably the 1993 expansion of the case definition for adults and adolescents (2)) have resulted in artificial jumps in AIDS case counts at the time the new definition went into effect or in the preceding year because changes in case definition allowed for retrospective diagnoses.

### Vital Status of HIV/AIDS Cases

Persons are assumed alive unless the MDH has knowledge of their death. Persons are assumed residing in Minnesota if their most recently reported state of residence was Minnesota and the MDH has not received notice of relocation outside of the state. Vital status information is updated by monthly visits to select reporting facilities, correspondence with other health departments, daily obituary reviews in local newspapers, annual death certificate reviews, and periodic matches with the National Death Index. "AIDS deaths" refers to all deaths among AIDS cases regardless of the cause.

### Place of Residence for HIV/AIDS Cases

Persons are assumed to be residing in Minnesota if their most recently reported state of residence was Minnesota and the MDH has not received notice of relocation outside of the state. Likewise, a person's county or city of residence is assumed to be the most recently reported value unless the MDH is otherwise notified. Residence information is updated through standard case reporting, monthly visits to select reporting facilities and/or correspondence with other state health departments. Persons diagnosed with HIV infection while imprisoned in a state correctional facility are included in the data presented unless otherwise noted (federal and private prisoners are excluded). Residential relocation, including release from state prison, is difficult to track and therefore data presented by *current* residence must be interpreted in this light. Data on residence at *time of diagnosis* are considered more accurate, limited only by the accuracy of self-reported residence location.

### Data Tabulation and Presentation

Unless otherwise noted, data analyses exclude persons diagnosed in federal or private correctional facilities (inmates generally are not Minnesota residents before incarceration and do not stay in Minnesota upon their release), infants with unknown or negative HIV status who were born to HIV positive mothers. Data include HIV-infected refugees who resettled in Minnesota as part of the HIV-Positive Refugee Resettlement Program.

The HIV/AIDS surveillance system is a live database that is continuously updated to reflect the most current information available. Variables such as current state of residence are over-written when updates are made. Annual archive files were initiated in 2001. Thus, the numbers of HIV/AIDS cases residing in Minnesota in 2000 and 2001 were estimated using the current state of residence variable while the number in previous years (1990-1999) was estimated using state of residence at time of diagnosis, vital status, and date of death variables. The number of HIV/AIDS cases alive in a certain year was calculated by summing cases with an HIV/AIDS diagnosis in that year or prior whose vital status in 2001 was "alive" or whose date of death was

either after the calendar year of interest or missing.

### Mode of Exposure Hierarchy

All state and city HIV/AIDS surveillance systems funded by the Centers for Disease Control and Prevention use a standardized hierarchy of mode of exposure categories. HIV and AIDS cases with more than one reported mode of exposure to HIV are classified in the exposure category listed first in the hierarchy. In this way, each case is counted as having only one mode of exposure. The only exception to this rule is the joint risk of male-to-male sex (MSM) and intravenous drug use (IDU), which makes up a separate exposure category in the hierarchy. The following is a list of the hierarchy for adolescent/adult HIV/AIDS cases:

- (1) MSM
- (2) IDU
- (3) MSM/IDU
- (4) Hemophilia patient
- (5) Heterosexual contact
- (6) Receipt of blood transfusion or tissue/organ transplant
- (7) Other (e.g. needle stick in a health care setting)
- (8) Risk not specified.

The following is the list of the hierarchy for pediatric HIV/AIDS cases:

- (1) Hemophilia patient
- (2) Mother with HIV or HIV risk
- (3) Receipt of blood transfusion or tissue/organ transplant
- (4) Other
- (5) Risk not specified.

Heterosexual contact is only designated if a male or female can report specific heterosexual contact with a partner who has, or is at increased risk for, HIV infection (e.g. an intravenous drug user). For females this includes heterosexual contact with a bisexual male (mainly due to the elevated prevalence of HIV infection among men who have sex with men).

"Risk not specified" refers to cases with no reported history of exposure to HIV through any of the routes listed in the hierarchy of exposure categories. These cases include persons who have not yet been interviewed by MDH staff; persons whose exposure history is incomplete because they died, declined to be interviewed, or were lost to follow-up; and persons who were interviewed or for whom follow-up information was available but no exposure was identified/acknowledged.

The growing number of cases with unspecified risk in recent years is, in part, artificial and due to interviews that have not yet been completed. In time, a number of these will be assigned a mode of exposure category. However, part of the observed increase is real. As stated above, a person must have intimate knowledge about his/her partner to meet the criteria for heterosexual mode of exposure. Often cases will not be certain about their partners' HIV status or risk. Additionally, the perception of social stigma presumably decreases the likelihood that a person will acknowledge certain risk behaviors, particularly male-to-male sex or injecting drug use. Thus, if the *true* numbers of cases due to heterosexual contact, MSM, and/or IDU increase, a larger number of cases without a specified risk would be expected.

A recent study by the Centers for Disease Control and Prevention used statistical methods to redistribute risk among female HIV/AIDS cases with unspecified risk (3). The results are helpful but are based on national data that are not necessarily applicable to the state or local level. Speculation regarding the distribution of risk behaviors among those with unspecified risk is difficult, especially in men, for whom even a national study is not available.

### Re-distribution of Mode of Exposure

In 2004 the Minnesota Department of Health began estimating mode of exposure for cases with unspecified risk in its annual summary slides. Estimation was done by using the risk distribution for living cases with known risk by race and gender and applying it to those with unspecified risk of the same race and gender. There were two exceptions to this method, African-born cases and Asian/Pacific Islander women. For both African-born and Asian/Pacific Islander women a breakdown of 95% heterosexual risk and 5% other risk was

used. For African-born males a breakdown of 5% male-to-male sex, 90% heterosexual risk, and 5% other risk was used. These percentages are based on epidemiological literature and/or community experience.

Below is an example of how the process worked for white, African American and African-born females:

<b>Living Cases Among Females in 2004</b>					
Race/Risk	Heterosexual n (%†)	IDU n (%†)	Other n (%†)	Unspecified n	Total N
White	178 (72)	58 (24)	10 (4)	78	324
African-American	174 (69)	62 (25)	16 (6)	107	359
African-born	43 (83)	0 (0)	9 (17)	223	275

† Percent of those with know risk.

<b>Female Cases with Estimated Risk</b>				
Race/Risk	Heterosexual	IDU	Other	Total N
White	$(.72*78) + 178 = 234$	$(.24*78) + 58 = 77$	$(.04*78) + 10 = 13$	324
African-American	$(.69*107) + 174 = 248$	$(.25*107) + 62 = 88$	$(.06*107) + 16 = 23$	359
African-born‡	$(.95*223) + 43 = 255$	0	$(.05*223) + 9 = 20$	275

‡ Used a distribution of 95% heterosexual and 5% other.

## Definitions Related to Race/Ethnicity

When data are stratified by race, Black race is broken down into African-born and African American (not African-born) based on reported country of birth.

The terms "persons of color" and "non-Whites" refer to all race/ethnicity categories other than White (Black, Hispanic, American Indian, and Asian/Pacific Islander).

## Interstate De-Duplication Project (IDEP)

In 2004, the Minnesota Department of Health (MDH) participated in IDEP. IDEP is a CDC project aimed at eliminating duplicate reports of HIV and AIDS cases among states. Each case of HIV and AIDS is assigned to the state (or states when the diagnosis of HIV and AIDS occurs in two different states) where a person was first diagnosed. The first round looked at cases reported through December 31, 2001. Through this project, MDH identified 164 cases of HIV infection (including AIDS at first report) and 55 AIDS cases whose first diagnosis was not in Minnesota. These cases were previously considered as diagnosed in Minnesota and were counted in the cumulative number of cases diagnosed in Minnesota. As such, the change of "ownership" (where the case was diagnosed) has reduced both cumulative and yearly totals for Minnesota. Additionally, MDH also identified 250 cases that no longer live in Minnesota.

The results of IDEP are particularly noticeable in the total number of persons living with HIV/AIDS in Minnesota, which increased from 4,895 to 5,002, a gain of only 107 cases, instead of 292 (new infections minus deaths plus (difference between people moving in and people moving out)) between 2003 and 2004.

- 
- (1) [Minnesota Rule 4605.7040 \(return to text\)](#)
  - (2) [MMWR 1992; 41\[no.RR-17\]:1-19 \(return to text\)](#)
  - (3) [MMWR 2001; 50\(RR-6\):31-40 \(return to text\)](#)

If you have questions or comments about this page, contact [idepcweb@health.state.mn.us](mailto:idepcweb@health.state.mn.us) or call 612-676-5414 (TTY: 612-676-5653) for the MDH Infectious Disease Epidemiology, Prevention and Control Division.

[MDH HOME](#) | [ABOUT US](#) | [LIBRARY](#) | [NORTH STAR](#)

MAIN CATEGORIES: [Health Data & Statistics](#) | [Diseases & Conditions](#) | [Terrorism & Natural Disasters](#) | [Facilities & Professions](#) | [Health Care & Coverage](#) | [Minnesota's Public Health System](#) | [People & Environment](#) | [Policy, Economics & Legislation](#)

[Comments and Questions](#) | [Phone Number, Address and Directions](#) | [Privacy Statement and Disclaimer](#)

Updated Thursday, 14-Apr-05 14:53:07



<b>Table 1. Number of Persons Living with HIV (non-AIDS) and AIDS by Residence, Age, and Gender* -- Minnesota, 2001</b>						
<b>Group</b>	<b>HIV (non-AIDS)</b>		<b>AIDS</b>		<b>Total</b>	
	<b>Cases</b>	<b>%</b>	<b>Cases</b>	<b>%</b>	<b>Cases</b>	<b>%</b>
<b>Residence**</b>						
Minneapolis	1,251	49%	844	48%	2,095	49%
St. Paul	363	14%	270	15%	633	15%
Suburban	642	25%	441	25%	1,083	25%
Greater Minnesota	279	11%	214	12%	493	11%
<i>Total</i>	2,535	100%	1,769	100%	4,304	100%
<b>Age***</b>						
<13 yrs	23	1%	7	0%	30	1%
13-19 yrs	17	1%	7	0%	24	1%
20-24 yrs	95	4%	18	1%	113	3%
25-29 yrs	182	7%	69	4%	251	6%
30-34 yrs	421	17%	180	10%	601	14%
35-39 yrs	609	24%	432	24%	1,041	24%
40-44 yrs	571	22%	426	24%	997	23%
45-49 yrs	324	13%	312	18%	636	15%
50-54 yrs	184	7%	176	10%	360	8%
55+ yrs	125	5%	145	8%	270	6%
<i>Total</i>	2,551	100%	1,772	100%	4,323	100%
<b>Gender</b>						
Male	2,034	79%	1,489	84%	3,523	81%
Female	525	21%	283	16%	808	19%
<i>Total</i>	2,559	100%	1,772	100%	4,331	100%
<b>State Totals</b>	2,559		1,772		4,331	

\* Cases reported to the MDH, assumed to be alive, and currently residing in Minnesota as of 12/31/01.

\*\* Residence information missing for 24 persons living with HIV and 3 persons living with AIDS.

\*\*\* Age missing for 8 persons living with HIV and 0 persons living with AIDS.

Suburban = Seven-county metropolitan area except Minneapolis & St. Paul (Anoka, Carver, Dakota, Hennepin (except Minneapolis), Ramsey (except St. Paul), Scott, and Washington counties). Greater Minnesota = Remaining 80 counties outside of the seven-county metropolitan area.

Numbers exclude federal and private prisoners, but include 101 state prisoners.

**Table 2. Number of Males & Females Living with HIV (non-AIDS) and AIDS  
by Race/Ethnicity and Mode of Exposure\* - Minnesota, 2001**

Group	Males				Females				Total			
	HIV (non-AIDS)	AIDS	Total		HIV (non-AIDS)	AIDS	Total		HIV (non-AIDS)	AIDS	Grand Total	
			Cases	%			Cases	%			Cases	%
<b>Race/Ethnicity</b>												
White, non-Hispanic	1,335	997	2,332	66%	185	96	281	35%	1,520	1,093	2,613	60%
Black**, African-American	435	287	722	20%	201	101	302	37%	636	388	1,024	24%
Black**, African-born	80	60	140	4%	78	48	126	16%	158	108	266	6%
Hispanic	109	108	217	6%	26	18	44	5%	135	126	261	6%
American Indian	32	23	55	2%	18	14	32	4%	50	37	87	2%
Asian/PI	17	14	31	1%	12	6	18	2%	29	20	49	1%
Unknown	26	0	26	1%	5	0	5	1%	31	0	31	1%
<i>Total</i>	<i>2,034</i>	<i>1,489</i>	<i>3,523</i>	<i>100%</i>	<i>525</i>	<i>283</i>	<i>808</i>	<i>100%</i>	<i>2,559</i>	<i>1,772</i>	<i>4,331</i>	<i>100%</i>
<b>Mode of Exposure</b>												
MSM	1,377	1,056	2,433	69%	--	--	--	--	1,377	1,056	2,433	56%
IDU	152	96	248	7%	69	62	131	16%	221	158	379	9%
MSM/IDU	135	109	244	7%	--	--	--	--	135	109	244	6%
Heterosexual	55	41	96	3%	250	127	377	47%	305	168	473	11%
Other	30	30	60	2%	21	12	33	4%	51	42	93	2%
Unspecified	285	157	442	13%	185	82	267	33%	470	239	709	16%
<i>Total</i>	<i>2,034</i>	<i>1,489</i>	<i>3,523</i>	<i>100%</i>	<i>525</i>	<i>283</i>	<i>808</i>	<i>100%</i>	<i>2,559</i>	<i>1,772</i>	<i>4,331</i>	<i>100%</i>

\* Cases reported to the MDH, assumed to be alive and currently residing in Minnesota as of 12/31/01.

\*\* Black race includes persons born in the U.S., Africa, or elsewhere.

MSM = Men who have sex with men. IDU = Injecting drug use. Heterosexual = For males: heterosexual contact with a female known to be HIV+, an injecting drug user, or a blood product or organ transplant recipient. For females: heterosexual contact with a male known to be HIV+, bisexual, an injecting drug user, or a blood product or organ transplant recipient. Other = Perinatal HIV exposure, receipt of blood product or organ/tissue transplant. Unspecified = Cases who did not acknowledge any of the risks listed above or have not been/refused to be interviewed.

Numbers exclude federal and private prisoners, but include 101 state prisoners.

**Table 3. Number of Persons Living with HIV (non-AIDS) and AIDS  
by County of Residence -- Minnesota, 2001**

<b>County**</b>	<b>HIV (non-AIDS)</b>	<b>AIDS</b>	<b>Total</b>
Aitkin	2	2	4
Anoka	72	61	133
Becker	0	3	3
Beltrami	4	3	7
Benton	3	2	5
Big Stone	0	1	1
Blue Earth	8	5	13
Brown	2	4	6
Carlton	9	4	13
Carver	14	8	22
Cass	1	3	4
Chippewa	0	0	0
Chisago	3	3	6
Clay	10	2	12
Clearwater	0	0	0
Cook	0	2	2
Cottonwood	1	2	3
Crow Wing	3	2	5
Dakota	87	62	149
Dodge	1	3	4
Douglas	1	4	5
Faribault	2	6	8
Fillmore	6	2	8
Freeborn	2	2	4
Goodhue	5	2	7
Grant	2	1	3
Hennepin	1,567	1,069	2,636
Houston	1	1	2
Hubbard	2	1	3
Isanti	4	2	6
Itasca	5	2	7
Jackson	0	0	0
Kanabec	1	2	3
Kandiyohi	4	5	9
Kittson	0	0	0
Koochiching	0	0	0
Lac Qui Parle	0	0	0
Lake	1	0	1
Lake of theWoods	0	0	0
Le Sueur	1	2	3
Lincoln	2	0	2
Lyon	6	1	7
McLeod	1	1	2
Mahnomen	0	1	1
Marshall	1	0	1
Martin	3	1	4
Meeker	3	1	4
Mille Lacs	1	3	4

Morrison	0	4	4
Mower	11	4	15
Murray	0	0	0
Nicollet	2	3	5
Nobles	8	3	11
Norman	1	2	3
Olmsted	30	27	57
Otter Tail	4	2	6
Pennington	1	0	1
Pine	4	4	8
Pipestone	0	1	1
Polk	5	3	8
Pope	1	3	4
Ramsey	439	315	754
Red Lake	0	0	0
Redwood	0	1	1
Renville	2	0	2
Rice	14	12	26
Rock	3	0	3
Rosseau	0	0	0
St. Louis	35	27	62
Scott	20	7	27
Sherburne	5	8	13
Sibley	0	0	0
Stearns	26	15	41
Steele	3	0	3
Stevens	2	1	3
Swift	0	0	0
Todd	1	0	1
Traverse	0	0	0
Wabasha	4	2	6
Wadena	1	1	2
Waseca	3	0	3
Washington	57	33	90
Watonwan	2	2	4
Wilkin	0	0	0
Winona	1	0	1
Wright	9	8	17
Yellow Medicine	0	0	0
<b>State Total**</b>	<b>2,535</b>	<b>1,769</b>	<b>4,304</b>

\* Cases reported to the MDH, assumed to be alive and currently residing in a Minnesota county as of 12/31/01.

\*\* Residence information missing for 24 persons living with HIV and 3 persons living with AIDS.

Numbers exclude federal and private prisoners, but include 101 state prisoners. State correctional facilities are located in the following counties: Anoka, Carlton, Chisago, Goodhue, Pine, Rice, Scott, St. Louis, Stearns, and Washington.

**Table 4. Number of HIV (non-AIDS) Cases, AIDS Cases, and AIDS Deaths\*  
Minnesota, 1990-2001**

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
HIV (non-AIDS)	341	365	361	225	224	219	187	192	191	202	177	197
AIDS	268	337	370	352	334	344	262	198	198	162	162	124
AIDS deaths	165	193	189	206	273	251	183	77	67	66	69	51

\* HIV (non-AIDS) = New cases of HIV infection (excluding AIDS at first diagnosis) diagnosed within a given calendar year. AIDS = All new cases of AIDS diagnosed within a given calendar year. AIDS deaths = Number of deaths known to have occurred among AIDS cases in a given calendar year, regardless of cause.

**Please Note:** These numbers refer to events, not individuals. For example, a person diagnosed as an HIV (non-AIDS) case in 1992 and then diagnosed as an AIDS case in 2000 will be counted twice in Table 4, once for each event. Thus, the numbers of HIV (non-AIDS) and AIDS cases cannot be summed over years to obtain cumulative totals. Please refer to the Minnesota HIV Surveillance Report, 2001 slide set for cumulative totals.

Case numbers exclude federal and private prisoners and refugees in the HIV-Positive Refugee Resettlement Program.

<b>Table 5. Known Mortality among Minnesota AIDS Cases by Year of Diagnosis -- Minnesota, through 2001*</b>				
<b>Year</b>	<b>Cases Diagnosed</b>	<b>Cases Known to be Dead**</b>	<b>Case-Fatality Rate***</b>	<b>Deaths Occurring in this Interval</b>
1982-1989	743	710	96%	368
1990	268	244	91%	165
1991	337	299	89%	193
1992	370	265	72%	189
1993	352	225	64%	206
1994	334	166	50%	273
1995	344	95	28%	251
1996	262	59	23%	183
1997	198	34	17%	77
1998	198	29	15%	67
1999	162	17	10%	66
2000	162	10	6%	69
2001	124	5	4%	51
<b>Cumulative Total</b>	<b>3,854</b>	<b>2,158</b>	<b>56%</b>	<b>2,158</b>

\* CDC 1993 AIDS definition used for all cases.

\*\* Cases known to be dead as of 12/31/2001. Reporting of deaths is incomplete.

\*\*\* Case-fatality rate is calculated by dividing the number of cases known to be dead by those diagnosed in a given interval and multiplying by 100.

Numbers exclude federal and private prisoners.