


A 2005 STUDY OF THE WATER QUALITY OF 172 METROPOLITAN AREA LAKES

June 2006


Randall J. Anhorn



Metropolitan Council

Mears Park Centre, 230 East Fifth Street, Saint Paul, Minnesota 55101
651/602-8743 TDD 651/291-0904
Publication No. 32-04-015

2005 Study of the Water Quality of 172 Metropolitan Area Lakes



By
Randall J. Anhorn

May 2006

EXECUTIVE SUMMARY

To date, the Council's lake monitoring programs (including the staff- and volunteer- monitoring programs) and have provided an important tool for making informed lake management decisions. Data from our regional lake monitoring programs are frequently used to determine possible trends in in-lake water quality, estimate expected ranges in water quality of unmonitored lakes, examine intra-and inter-regional differences, determine potential impairments due to water quality, and investigate the relationships between landscape and water quality.

This report is the latest in a continuing series of reports summarizing results of the Metropolitan Council's (Council) annual lake monitoring program. The Council has collected water quality data on area lakes since 1980. This report contains data from 172 lake sites sampled in 2005, including 12 lake sites on 10 lakes monitored by the Council and 160 lakes monitored by volunteers.

Eighty-six of the 172 lakes monitored in 2005, are listed by the MPCA as impaired waters due to excessive nutrients (phosphorus) affecting the lakes' ability to support their designated recreational uses. Eighty of those lakes were monitored through CAMP, and six were monitored by Council-staff. To learn more about the listing and potential next steps <http://www.pca.state.mn.us/water/tmdl/index.html>.


The objectives of this study were to:

1. Provide lake water quality data to lake, watershed and water resource managers.
2. Advise managers of known or suspected threats to lake water quality.
3. Continue to compile a water quality database on the six area lakes that support a trout fishery.
4. Collect in-lake water quality data on, Alimagnet, Lee, Twin [Burnsville], and Valley lakes to determine the results of in-lake organic carbon amendment (barley straw or cornmeal treatments) on the lakes' algal population and resulting water clarity.
5. Support the Council's satellite lake assessment program by collecting "ground" measurements used to develop empirical models in order to predict water clarity on all the regions lakes. Information about, and results from, the Council's satellite lake assessment program can be found at <http://www.metrocouncil.org/planning/environment/TCWaterClarity2005.pdf>
6. Collect water quality data on Lake McCarron in order to evaluate the "success" of modifications to the Villa Park wetland treatment system and autumn of 2004 in-lake Alum treatment. The lake will again be monitored in 2006 to continue to assess the success of the lake and watershed projects and their effect on the lake's water quality.

The year 2005 marked the thirteenth year that the Council-sponsored volunteer monitoring program, entitled "The Citizen-Assisted Monitoring Program" (CAMP), was used to increase our knowledge of the water quality of area lakes. Once again volunteers measured surface water temperature and transparency, and collected surface water samples that were analyzed for total phosphorus, total Kjeldahl nitrogen, and chlorophyll-a on a biweekly basis from mid-April to mid-October (approximately 14 sampling events).

This year's monitoring program included 17 lakes never before monitored by the Council, and 129 lake sites returning from 2004. The 2005 lakes program included lake data from all of the 26 watersheds/municipalities/counties represented in the 2004 program. Additionally, the 2005 CAMP program added three new enrolling groups to its growing list of monitoring partners.

The greatest percentage of the lakes monitored through CAMP in 2005 received an overall water quality grade of “C” (38.7 percent). When combining the CAMP and Council-staff monitored lake grades 40.1 percent of the lakes received an overall grade of “C”. The water quality of these lakes is considered average as compared to others in the seven-county metropolitan area. When comparing the percentage of above-average lakes (those receiving grades of “A” or “B”) to below-average lakes (those receiving “D” or “F”), more lakes were below average (34 percent to 27 percent). The complete 2005 CAMP lake report card grade tally (for those lakes with sufficient data) assigned “A’s” to 21 lakes (13.5 percent) and “B’s” to 21 lakes (13.5 percent). Sixty lakes acquired “C’s” (38.7 percent), 34 received “D’s” (21.9 percent), and 19 obtained an “F” (12.3 percent).



Of the 129 repeat lakes which a sufficient database from 2004, 24 had a worse overall water quality grade in 2005 (Bald Eagle [Site-1], Barkers, Bavaria, DeMontreville, Forest [West Basin], Golden, Goose [Waconia], Hart, Herber’s Pond, Island, Kismet, Long [Stillwater], Long [Mahtomedi], Louise, MacDonald’s Pond, McDonald, McKusick, Mud, Oak, Olson, Sand, Spring, Schroeder’s Pond, and Westwood lakes), and 14 had better overall water quality grades in 2005 (Big Comfort, Cloverdale, Downs, Eagle (Carver County), George Watch, Hay, McCarrons, Miller, Prior [Lower], Prior [Upper], Staples, Swede, Wilmes, and Windsor lakes), and 91 had the same overall water quality grade for both years. By further breaking down the 91 lakes that had identical overall grades in 2004 and 2005, 44 had similar summertime mean conditions in both 2004 and 2005 (mean TP, CLA and Secchi transparency), 29 had worse means in 2005, and 18 had better means in 2005.

Water quality data from the 129 repeat lakes seem to indicate that the Metro Area lakes experienced slightly worse water quality conditions in 2005 as compared to 2004. This after two years where the lake monitoring program reveal slightly better water quality as that recorded during the previously monitored years (2003 better than 2002, and 2004 better than 2003).

A recently conducted trend analysis by MPCA on lakes with extensive Secchi transparency databases however, revealed that while the majority of statistically assessed lakes showed no trends in water clarity (either negative or improving), more lakes showed an improving trend than a negative trend (MPCA 2005). Of the CAMP 2005 lakes assessed (those with sufficient data), 14 showed an improving trend in water clarity (Big Marine, DeMontreville, Elmo, Halfbreed/Sylvan, Langton (south basin), Little Carnelian, Long (May Township), McKusick, Olson, Sunset, Valentine, Valley, Waconia, and West Boot

lakes) and four showed a negative trend (Farquhar, Lac Lavon, Markgrafs, and Square lakes) (MPCA 2005).

Overall, six of the 12 Council-staff monitored lake sites monitored in 2005 were also monitored in 2004. While four of the six lakes that were monitored through Council programs in 2004 and 2005 received identical overall water quality grades in both years, one lake experienced improved water quality in 2005 (McCarrons) and one had worse (Bald Eagle Site-1). In fact, Lake McCarrons, which had an in-lake alum treatment in the fall of 2004, improved from an overall grade of C in 2004 to A in 2005.

Since 1980, 297 area lakes have been monitored through the Council's Lake Program (including Council-staff monitoring and CAMP). Some of the lakes have multiple monitoring sites [316 sites]. The list of lakes in the Council's monitoring database is shown in Appendix C. The resulting data from the Council's lake monitoring program are permanently stored in the U.S. EPA's national water quality data bank, STORET (stands for STOrage and RETrieval). The majority of the 316 lake sites have been revisited on a rotating schedule throughout the past 26 years to develop a working baseline to help determine possible trends and to aid lake and watershed managers in their decision making. While the Council has done its best to enhance and expand the region's lake water quality database, it is apparent that one of the most economical and efficient method to expand knowledge of our lakes has been with the assistance of volunteers and cooperation and financial support of watershed management organizations, counties, and cities. So while the first 13 years of CAMP have been very successful, our future goal is to continue to expand the coverage of our lake monitoring program in order to better understand and manage the areas water resources.

The Council's lake monitoring program, especially the use of volunteer monitors through CAMP, has played a key role in the Council's recent efforts to use satellite images to assess annual lake water clarity for the region as a whole. The monitoring program provides "ground-based" measurements used to calibrate mathematical models, which in turn are used to interpret the satellite images. The use of satellite technology provides a cost-effective way to extend the analysis of the region's lake water quality from just the lake's involved in our ground-based programs to all the lakes in the region. Over time, the satellite-based information can be used to detect how lake trophic conditions (especially water clarity) have changed over time and space in relation to changes in land-use and land-cover conditions.

Results of the 2005 satellite assessment of the region revealed similar results to that found through the 2004 ground-based monitoring programs, that the region experienced slightly worse lake water quality in 2005 than that recorded in 2004. The complete results of the 2005 satellite analysis can be at <http://www.metrocouncil.org/planning/environment/TCWaterClarity2005.pdf>.

If you have questions pertaining to the lake data or descriptions contained in this report, inquiries about CAMP, or suggestions of lakes the Council should consider monitoring in the future, please contact Randy Anhorn at the Metropolitan Council (651) 602-8743 or randy.anhorn@metc.state.mn.us.

ACKNOWLEDGMENTS

This report represents the coordinated efforts of many individuals. I would like to acknowledge the following people for their technical and supportive contributions to the preparation of this report:

- The various watershed management organizations (WMOs), participating agencies, and volunteers involved in the citizen-assisted monitoring program (CAMP), for without their enthusiastic participation, CAMP would not be successful. A list of involved WMOs, agencies, and volunteer lake monitors is shown in Appendix B. The following of this years volunteers should be given added thanks for their multiple years of service:

13 years of service

Diane and Bob Coderre - Sunset Lake

12 years of service

Washington Co. SWCD- Multiple

11 years of service

Bill Aamadt- Wilmes Lake

Janet and Harvey Bartz- Seidl's Lake

Carver Co. Env. Services- Multiple

Wayne LeBlanc- Lake Peltier

10 years of service

City of Circle Pines- Golden Lake

John Ritter - Lake Alimagnet

Wargo Nature Center- George Watch

9 years of service

Anoka Co. Parks- Multiple

8 years of service

Glen Gramse- Keller Lake

Mona and David Hanson- Pike Lake

Wally Shaver- Lac Lavon Lake

7 years of service

Philip Goodrich- Pike Lake

Hvass Family - Colby Lake

Lakeville- Valley and Lee lakes

John Ryski- Bavaria Lake

Westwood Nature Center- Westwood Lake

6 years of service

Steve Bur- Northwood Lake

Dave Hanson-Sweeney Lake

Renay Leone-Virginia Lake

Dale Wahlstrom- Schmidt Lake

5 years of service

Arnett Family- Crystal Lake

Gene Berwald- Pine Tree Lake

Kevin Bjork- Cloverdale Lake

Tom/Dorothy Goodwin- Orchard Lake

Marvin Groth- Bass Lake (Hennepin Co.)

Madison Groves- Upper Prior Lake

Green Family- Kingsley Lake

Deb Gutzman- Reshanau Lake

Ryan Opdahl- Little Johanna

Wally Potter- Marion Lake

Rice Creek WD- Multiple

Terry Riley- Markgrafs Lake

Mike Shouldrice- Tamarack and Schutz lakes

Sly Family- Downs Lake

Phillip Solseng- Long Lake

Streff Family- South Rice Lake

Bob Videen- Parkers Lake

4 years of service

Kathy Gerlach- Dean Lake

Bonnie Jurand- Klawitter Lake

Nigel Linden- Twin Lake (Burnsville)

Mary Oaster- Earley Lake

Wally/Ardyce Potter- Marion Lake

Serie/Kettlekamp- Long Lake (A.V.)

Tom Sletta- Cates Lake

Diane Stauner- Meadow Lake

3 years of service

Rick Bruneau- Farquhar Lake

Walt Burris- Lower Prior Lake

Conservation League of Edina-Multiple

Bill Feely- Long Lake

Grove Family-McDonald Lake

Kellogg Family- Multiple lakes

Chris Moosbrugger- Mergens Lake

Kitty Francy-Payton- Long Lake

Bill Tisdell- Spring Lake

Ed Voyles- Twin Lake

- The Metropolitan Council Environmental Services - Environmental Planning and Evaluation department (MCES-EPE) for laboratory analysis of the lake samples.
- And, the members of the Metropolitan Council and its environmental staff, Jeffrey Jax, and Craig Skone for support with lake monitoring, data management, and report preparation. Craig Skone deserves additional recognition for developing all the graphics for this report.

CONTENTS

This report is divided into two parts: Part I describes the lake water quality of the 12 lake sites monitored by the Metropolitan Council-staff, while Part II details the water quality of the 160 lakes monitored by volunteers as part of the Citizen-Assisted Monitoring Program.

Executive Summary	i
Acknowledgments	v

PART I - METROPOLITAN COUNCIL 2005 LAKE MONITORING PROGRAM

Introduction	1
Methods	4
Results/Analysis	5
Lake Quality Report Card.....	9
Bald Eagle Lake (Site-1)	11
Bald Eagle Lake (Site-2)	12
Cedar Lake [Scott County]	13
Forest Lake [East Basin]	14
Forest Lake [Middle Basin]	15
McMahon Lake	16
McCarrons Lake	17
O'Dowd Lake	18
Reitz Lake	19
Thole Lake	20
Whaletail Lake (Site-1)	21
Whaletail Lake (Site-2)	22

PART II - CITIZEN-ASSISTED LAKE MONITORING PROGRAM

Acknowledgments	23
Introduction	23
Purpose of Volunteer Program	24
CAMP Methods	24
Obtaining Volunteers	24
Training Volunteers	25
Monitoring Methods	25
Data Handling and Analysis	29
Program Quality Assurance/Quality Control	29
Results and Analysis	33
Alimagnet Lake	34
Armstrong Lake	38
Barker Lake	40
Bass Lake [Hennepin County]	42
Bass Lake [St. Louis Park]	44
Bass Lake [Washington County]	46
Bavaria Lake	48
Benton Lake	52
Benz Lake	54
Big Carnelian Lake	56
Big Comfort Lake	58
Big Marine Lake	60
Birch Lake	62
Bone Lake	64
Brickyard Lake	66
Burandt Lake	68

Campbell Lake	70
Carol Lake	72
Cates Lake	76
Cenaiko Lake	78
Cloverdale Lake	82
Cobblecrest Lake	84
Cobblestone Lake	86
Colby Lake	88
Cornelia Lake	90
Courthouse Lake	92
Crystal Lake [Burnsville]	94
Crystal Lake [Robbinsdale]	96
Dean Lake	100
DeMontreville Lake	102
Downs Lake	104
Eagle Lake (Carver County)	106
Eagle Lake (Maple Grove)	108
Earley Lake	110
East Lake	112
East Boot Lake	114
Edina Lake	116
Edith Lake	118
Elmo Lake	120
Farquhar Lake	122
Fireman's Lake	126
Fish Lake [Scott County]	128
Fish Lake [Washington County]	130
Forest Lake [West Basin]	132
Gaystock Lake	134
George Watch Lake	136
German Lake	138
Goetschel Lake	140
Goggins Lake	142
Golden Lake	144
Goose Lake [Scandia]	146
Goose Lake [Waconia]	148
Hafften Lake	150
Half Breed (Sylvan) Lake	152
Hart Lake	154
Hay Lake	156
Hazeltine Lake	158
Henry Lake	160
Herber's Pond	162
Highland Lake	164
Hydes Lake	168
Island Lake	170
Jane Lake	172
Jellum's Bay [Site-1]	174
Keller Lake [Burnsville]	176
Kingsley Lake	178
Kismet Lake	180
Klawitter Lake	182
La Lake	184
Lac Lavon Lake	186
Langton Lake (North basin)	188
Langton Lake [Site-1]	190

Langton Lake [Site-2]	192
Lee Lake.....	194
Legion Pond Lake	198
Lily Lake	200
Little Carnelian Lake.....	202
Little Johanna Lake.....	204
Long Lake [Apple Valley]	206
Long Lake [Mahtomedi]	208
Long Lake [May Township]	210
Long Lake [Pine Springs]	212
Long Lake [Stillwater]	214
Long Lake [Washington County].....	216
Loon Lake	218
Lotus Lake.....	220
Louise Lake.....	222
MacDonald's Pond.....	224
Maple Marsh Lake	226
Maria Lake	228
Marion Lake	230
Markgrafs Lake	232
McDonald Lake.....	234
McKusick Lake	236
Meadow Lake.....	238
Mergen's Lake	240
Miller Lake.....	242
Mitchell Lake	244
Moody Lake	246
Mud Lake	248
North Twin Lake.....	250
Northwood Lake.....	252
Oak Lake	254
O'Connor Lake.....	256
Olson Lake	258
Orchard Lake.....	260
Pamela Lake	262
Parkers Lake.....	264
Peltier Lake	266
Pike Lake [Ramsey County].....	270
Pike Lake [Scott County] [Site-1]	272
Pine Tree Lake	276
Powers Lake.....	278
Prior Lake [Lower Basin Site-1]	282
Prior Lake [Upper Basin Site-1].....	286
Regional Park Lake	288
Riley Lake	290
Sand Lake.....	292
Schmidt Lake	294
School Lake.....	296
Schroeder's Pond	298
Schutz Lake.....	300
Seidl's Lake.....	304
Shaver Lake.....	308
Shields Lake.....	310
Silver Lake	314
South Rice Lake	316
South School Section Lake	318

South Twin Lake	320
Spring Lake [Scott County].....	322
Square Lake.....	326
Staples Lake	330
St. Croix Lake [Whole Lake]	332
St. Croix Lake [Bayport Pool-Site 2]	334
St. Croix Lake [Troy Beach Pool-Site 3]	336
St. Croix Lake [Troy Beach Pool-Site 5]	338
St. Croix Lake [Black Bass Pool-Site 6]	340
St. Croix Lake [Kinnickinnic Pool-Site 7]	342
St. Joe Lake	344
Sunfish Lake.....	346
Sunnybrook Lake	348
Sunset Lake	350
Swede Lake	352
Sweeney Lake	354
Tamarack Lake.....	358
Terrapin Lake	360
Turtle Lake.....	362
Twin Lake [Burnsville]	364
Twin Lake [Lower] [Robbinsdale].....	366
Twin Lake [Middle] [Crystal]	368
Twin Lake [St. Louis Park].....	370
Valentine Lake	372
Valley Lake	374
Virginia Lake	378
Waconia Lake.....	380
West Boot Lake.....	384
Westwood Lake.....	386
Wilmes Lake	388
Windsor Lake	390
Winkler Lake.....	392
Wood Lake [Burnsville].....	394
Conclusions	398
REFERENCES	402
APPENDICES	
A. 2005 CAMP Lake/Watershed Characteristics	409
B. 2005 Volunteer Lake Monitors	418
C. Lakes Sampled by the Metropolitan Council, 1980-2005	422
FIGURES	
1. 2005 Metropolitan Council Study Lakes.....	3
2. 2005 CAMP Study Lakes.....	26
3. Example of CAMP Sampling Form	28
4. 2005 CAMP-collected TP vs. professionally-collected ‘quality check’ TP	31
5. 2005 CAMP-collected CLA vs. professionally-collected ‘quality check’ CLA	31
6. 2005 CAMP-collected Secchi transparency vs. professionally-collected ‘quality check’ Secchi transparency	31
7. 1993-2005 CAMP-collected TP vs. professionally-collected ‘quality check’ TP.....	32
8. 1993-2005 CAMP-collected CLA vs. professionally-collected ‘quality check’ CLA	32
9. 1993-2005 CAMP-collected Secchi transparency vs. professionally-collected ‘quality check’ Secchi	32

TABLES

1. Trends in Total Phosphorus Concentration	6
2. Trends in Chlorophyll- <i>a</i> Concentration	7
3. Trends in Secchi Disk Transparency	8

Physical/chemical lake data and copies of the volunteer monitoring methods pilot study can be obtained upon request by contacting Randy Anhorn at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

PART I - METROPOLITAN COUNCIL 2005 LAKE MONITORING PROGRAM

INTRODUCTION

The Metropolitan Council-staff sampled 12 lake sites on 10 lakes in 2005 as part of its continuing effort to manage lakes in the Twin Cities Metropolitan Area (TCMA) (Figure 1). This report follows a series of lake studies (Appendix C):

<u>YEAR</u>	<u>NUMBER OF LAKES</u>	<u>REFERENCE</u>
1980	60	Osgood (1981)
1981	30	Osgood (1982a)
1982	7	Osgood (1983)
1983	28	Osgood (1984a)
1984	43	Osgood (1984b)
1985	32	Osgood (1985)
1986/87	10	Osgood (1988a)
1988	6	Osgood (1989a)
1989	20	Osgood (1989b)
1990	21	Osgood (1990)
1991	17	Hartsoe and Osgood (1991)
1993	12 (+ 31 CAMP lakes)	Anhorn (1994)
1994	13 (+ 38 CAMP lakes)	Anhorn (1995)
1995	13 (+ 46 CAMP lakes)	Anhorn (1996)
1996	13 (+ 53 CAMP lakes)	Anhorn (1997)
1997	12 (+ 59 CAMP lakes)	Anhorn (1998)
1998	13 (+ 57 CAMP lakes)	Anhorn (1999)
1999	14 (+ 99 CAMP lakes)	Anhorn (2000)
2000	14 (+110 CAMP lakes)	Anhorn (2001)
2001	12 (+120 CAMP lakes)	Anhorn (2002)
2002	12 (+125 CAMP lakes)	Anhorn (2003)
2003	12 (+128 CAMP lakes)	Anhorn (2004)
2004	13 (+132 CAMP lakes)	Anhorn (2005)
2005	12 (+160 CAMP lakes)	Anhorn (2005)

The long-term goal of the Council's lake studies has been to provide a comprehensive database to enable cities, counties and watershed management organizations (WMOs) to better manage area lakes. The Council believes that without such comprehensive lake data, the foundation of lake and watershed management plans is weakened. While the Council has provided a commendable lake data collection program, the data collection of others, specifically WMOs, is encouraged (Osgood 1989c). Several agencies and cities have taken initiative (for example, Ramsey County, Eagan, Maple Grove, and Minneapolis Park and Rec. Board), but for the most part the WMOs are not collecting adequate data.


To date, the Council's lake monitoring programs have provided an important tool for making informed lake management decisions. Data from our regional lake monitoring programs are frequently used to determine possible trends in in-lake water quality, estimate expected ranges in water quality of unmonitored lakes, examine intra-and inter-regional differences, and investigate the relationships between landscape and water quality. A comprehensive regional lake monitoring program should ensure adequate representation across both space and time. However, due to cost and logistical problems, ground-based monitoring programs usually sacrifice spatial coverage (fewer lakes) in favor of more frequent sampling.

The Council addressed this lack of adequate data collection problem by initiating a citizen-assisted lake monitoring program (CAMP) in 1993. CAMP is funded in part by watershed districts (WDs), WMOs, counties, and cities that are participating in the program. Through this program, citizens collect comprehensive data. To assure that the data collection methods used by citizen volunteers are credible; the Council conducted a pilot study along with its routine monitoring in 1991 (Hartsoe and Osgood 1991). The pilot study and its results are included in the 1993 lake report, and can be obtained by contacting Randy Anhorn at (651) 602-8743 or randy.anhorn@metc.state.mn.us. The methods and results of the CAMP for 2005 are described in Part II of this report.

The Council's lake monitoring program, especially the use of volunteer monitors through CAMP, has played a key role in the Council's recent efforts to use satellite images to assess annual lake water clarity for the region as a whole. The monitoring program provides "ground-based" measurements used to calibrate mathematical models, which in turn are used to interpret the satellite images. The use of satellite technology provides a cost-effective way to extend the analysis of the region's lake water quality from just the lake's involved in our ground-based programs to all the lakes in the region. Over time, the satellite-based information can be used to detect how lake trophic conditions (especially water clarity) have changed over time and space in relation to changes in land-use and land-cover conditions.

Results of the 2005 satellite assessment of the region revealed similar results to that found through the 2004 ground-based monitoring programs, that the region experienced slightly worse lake water quality in 2005 than that recorded in 2004. The complete results of the 2005 satellite analysis can be at <http://www.metrocouncil.org/planning/environment/TCWaterClarity2005.pdf>.

Figure 1.
2005 Study Lakes



METHODS

Twelve sites on 10 lakes were sampled by Council-staff at two-week intervals from mid-April through mid-October, 2005. The lakes were normally visited between 8:00 a.m. and noon on the sampling days. Samples were collected from one station located over the deepest spot near the center of the lakes (the sampling location(s), as well as graphs of the seasonal data are shown on lake information sheets located in alphabetical order at the end of Part I of this report).

A hand-held Global Positioning System (GPS) was used to lock in sampling location coordinates (shown as latitude and longitude on the lake information sheets), and to aid in relocating sampling locations during each ensuing monitoring event. Time, surf and weather conditions, and station depth were recorded upon anchoring at the site. Temperature, dissolved oxygen, pH, specific conductivity, turbidity, chloride, and oxidation reduction potential were measured at one-meter intervals (additional readings are captured at half-meter intervals near the thermocline) using a Yellow Springs, Inc. (model 650 MDS) multiparameter field monitoring system. The YSI was calibrated in the morning, prior to the daily monitoring, and again after the last lake was monitored on that day. Water transparency was measured using a 20 cm black-and-white Secchi disk.

Water was collected from the lakes' surface (0-2 m) using a two-meter PVC pipe that held two liters of water. Two or three such samples were mixed in an 8-liter plastic jug. Subsurface samples (middle and near bottom) are drawn using a 2-liter Van Dorn. All water samples were transported on ice in a dark cooler and processed and preserved within six hours of collection. Water from the surface jug was withdrawn for the following chemical analyses (depending on the lake): total phosphorus (TP), total dissolved phosphorus (TDP), total Kjeldahl nitrogen (TKN), chlorophyll-*a* (CLA), and chloride (Cl). Subsurface water samples were also drawn using a 2-liter Van Dorn. Subsurface samples were taken for TP and Cl analysis on all lakes deeper than 2.5 meters, and for TDP on Bald Eagle (sites 1 and 2), Centerville, and McCarrons lakes.

The routine chemical analyses were performed at the Metropolitan Council Environmental Services - Environmental Planning and Evaluation department (MCES-EPE) laboratory following U.S. EPA approved methods. Surface and subsurface water samples that were analyzed for TDP were filtered through a 0.45 µm membrane filter and analyzed for TP. Water samples tested for phosphorus and TKN were digested with the sulfates of hydrogen, potassium and mercury (H₂SO₄, K₂SO₄ and HgSO₄). Following digestion, phosphorus was analyzed using a modified ascorbic acid reduction method (APHA 1980). Samples tested for TKN were chemically reduced the same way as the total phosphorus samples, then were color-intensified with sodium nitroprusside and assayed for ammonia colorimetrically. TKN and TP from the surface were periodically analyzed in duplicate to determine accuracy, at which time their average values were reported.

Water samples to be analyzed for CLA were filtered onto a 0.45 µm glass-fiber-filter, saturated with magnesium carbonate, and stored frozen in the dark until analyzed (within 30 days). Chlorophyll was extracted from the filters by homogenization in 90 percent aqueous acetone. The optical density of the extract was measured spectrophotometrically at 630, 647, 664 and 750 nm. CLA was calculated from a trichromatic equation that corrects for turbidity (APHA 1980).

RESULTS/ANALYSES

Tables 1, 2, and 3 show summertime average phosphorus concentration in micrograms per liter (µg/l), chlorophyll-*a* concentration in µg/l, and Secchi transparency in meters (m), for the 13 lake sites monitored by the Council-staff. Raw data will be input into the STORET database, or it can be obtained upon request by contacting Randy Anhorn at (651) 602-8743 or randy.anhorn@metc.state.mn.us. Tables 1, 2, and 3, also document summertime means (May through September) for any prior years the lake was monitored by the Council. Seasonal data is graphed for each lake at the end of Part I of this report.

Due to normal seasonal variability, insufficient data collection intensity for each lake, and changing climatological conditions, determining long-range water quality trends in area lakes is generally not statistically reliable. Accurate conclusions are difficult because one year's data may represent only one monitoring date or parameter, water quality may fluctuate greatly from year to year, and/or the lake may only be monitored once every ten years. Therefore, to fully determine if there truly is a change in the water quality of a lake, either additional years of data collection are needed in the future to accurately determine the present condition of the lake, and/or a broader, more complete historical baseline database is needed.

While an extensive database of a lake's present water quality is obtainable; a more extensive historical database is not. In other words, without a complete and accurate historical database, which is rare, it is difficult to determine if a lake's quality has changed because it is not known what its quality used to be. Therefore, an extensive baseline database needs to be constructed now so lake quality trends can be determined in the future. Many of 12 lake sites monitored by Council staff in 2005 have databases that are insufficient in size and quality to determine "statistically significant" long-range trends. Statistical trend analysis on the few lake databases which did contain sufficient data revealed improving water clarity trend on one, Bald Eagle Lake (MPCA 2005).

After comparing the 2005 data to pre-2005 data on the 12 lake sites, a few general comments and observations can be made. A review of each lakes' summertime TP, CLA and Secchi means and water quality grades reveal that all but one (Lake McCarrons), seem to have water quality levels that fall within their normal fluctuation ranges of seasonal water quality.

Overall, six of the 12 Council-staff monitored lake sites monitored in 2005 were also monitored in 2004. While four of the six lakes that were monitored through Council programs in 2004 and 2005 received identical overall water quality grades in both years, one lake experienced improved water quality in 2005 (McCarrons) and one had worse (Bald Eagle Site-1). In fact, Lake McCarrons, which had an in-lake alum treatment in the fall of 2004, improved from an overall grade of C in 2004 to A in 2005.

Table 1
Trends in May - September average surface total phosphorus concentration (µg/l)

Lake	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Bald Eagle (Site-1)	80	63	--	47	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	62	84
Bald Eagle (Site-2)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	65	77
Cedar (Scott Co.)	442	341	--	--	168	--	--	--	--	--	--	--	--	197	--	--	--	--	263	--	--	154	--	--	--	127
Forest (East Basin)	40	--	--	--	36	--	--	--	--	--	--	--	--	--	40	--	--	33	--	--	--	--	26	--	--	35
Forest (Mid. Basin)	--	--	--	--	40	--	--	--	--	--	--	--	--	--	37	--	--	31	--	--	--	--	23	--	--	41
McCarrons	--	--	--	--	20	16	14	26	30	17	18	30	--	15	13	28	16	--	--	19	29	19	28	--	17	7
McMahon	240*	--	--	--	105	--	--	--	--	--	--	--	--	--	--	113	--	--	80	--	--	126	--	--	--	117
O'Dowd	--	--	--	--	57	--	--	--	--	--	--	--	--	--	67	--	--	--	59	--	61	--	78	--	--	60
Reitz	--	--	--	--	--	111	--	--	--	--	--	109	--	149	--	--	--	--	--	49	55	93	107	111	107	95
Thole	--	--	--	--	83	--	--	--	--	--	--	--	--	--	154	--	--	--	91	--	140	--	144	--	--	112
Whaletail (Site-1)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	76	75
Whaletail (Site-2)	70	--	--	--	60	--	--	--	--	--	--	--	--	--	--	--	--	--	32	--	--	45	--	--	52	53

* Sampled only twice in 1980

Table 2
Trends in May - September average surface chlorophyll-a concentration (µg/l)

Lake	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Bald Eagle (Site-1)	65	57	--	41	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	45	48
Bald Eagle (Site-2)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	46	47
Cedar (Scott Co.)	266	27	--	--	73	--	--	--	--	--	72	--	--	24	--	--	--	--	56	--	--	107	--	--	--	39
Forest (East Basin)	49	--	--	--	32	--	--	--	--	--	17	--	--	--	18	--	--	15	--	--	--	--	15	--	--	22
Forest (Mid. Basin)	--	--	--	--	34	--	--	--	--	--	20	--	--	--	18	--	--	12	--	--	--	--	14	--	--	23
McCarrons	--	--	--	--	20	16	14	26	30	17	18	30	--	15	13	28	16	--	--	19	29	19	28	--	17	7
McMahon	85*	--	--	--	73	--	--	--	--	--	--	--	--	--	--	50	--	--	64	--	--	69	--	--	--	90
O'Dowd	--	--	--	--	48	--	--	--	--	--	--	--	--	--	53	--	--	43	--	--	44	--	52	--	--	48
Reitz	--	--	--	--	--	93	--	--	--	--	--	49	--	43	--	--	--	--	--	17	29	56	48	53	39	47
Thole	--	--	--	--	73	--	--	--	--	--	--	--	--	--	72	--	--	44	--	--	64	--	60	--	--	93
Whaletail (Site-1)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	46	44
Whaletail (Site-2)	65	--	--	--	53	--	--	--	--	--	--	--	--	--	--	--	--	--	27	--	--	31	--	--	47	45

* Sampled only twice in 1980

Table 3
Trends in May - September average Secchi disk transparency (m)

Lake	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Bald Eagle (Site-1)	0.8	1.2	--	1.4	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	1.2	1.2
Bald Eagle (Site-2)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	1.2	1.2
Cedar (Scott Co.)	1.4	2.1	--	--	1.4	--	--	--	--	--	0.9	--	--	1.9	--	--	--	--	1.2	--	--	0.7	--	--	--	1.3
Forest (East Basin)	1.3	--	--	--	1.4	--	--	--	--	--	2.0	--	--	1.7	--	--	1.8	--	--	--	--	--	2.0	--	--	1.9
Forest (Mid. Basin)	--	--	--	--	1.3	--	--	--	--	--	1.8	--	--	1.6	--	--	1.7	--	--	--	--	--	2.0	--	--	1.9
McCarrons	--	--	--	--	2.3	2.1	2.8	1.8	1.4	1.8	3.1	1.5	--	2.1	2.3	1.8	1.7	--	--	1.8	2.2	3.1	2.0	--	--	4.0
McMahon	1.5*	--	--	--	1.0	--	--	--	--	--	--	--	--	--	--	1.8	--	--	1.1	--	--	0.9	--	--	--	0.8
O'Dowd	--	--	--	--	1.4	--	--	--	--	--	--	--	--	--	1.2	--	--	1.4	--	--	1.2	--	1.4	--	--	1.4
Reitz	--	--	--	--	--	0.9	--	--	--	--	--	1.3	--	1.1	--	--	--	--	--	1.5	1.4	0.7	1.3	2.8	2.0	1.9
Thole	--	--	--	--	1.0	--	--	--	--	--	--	--	--	--	1.0	--	--	1.7	--	--	1.4	--	0.8	--	--	1.2
Whaletail (Site-1)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	0.6	0.7
Whaletail (Site-2)	0.7	--	--	--	0.8	--	--	--	--	--	--	--	--	--	--	--	--	--	1.0	--	--	1.0	--	--	0.9	0.9

* Sampled only twice in 1980

LAKE QUALITY REPORT CARD

The Metropolitan Council following its 1989 lake survey (Osgood 1989b) developed the lake quality report card. The idea is simply that lake water quality characteristics can be ranked by comparing measured values to those of other Metro Area lakes. In this way, technical information, which in the past had required professional analysis, can more easily be used by a less technical audience to visualize the water quality of their lake relative to other area lakes. The grading curve represents percentile ranges for three water quality indicators - the summertime (May - September) average values for total phosphorus, chlorophyll-a, and Secchi disk. These percentiles use ranked data from 120 lakes sampled from 1980 - 1988:

<u>GRADE</u>	<u>PERCENTILE</u>	<u>TP(μg/l)</u>	<u>CLA(μg/l)</u>	<u>Secchi(m)</u>
A	<10	<23	<10	>3.0
B	10-30	23-32	10-20	2.2-3.0
C	30-70	32-68	20-48	1.2-2.2
D	70-90	68-152	48-77	0.7-1.2
F	>90	>152	>77	<0.7

In 2000, the percentiles determined from the 1980-1988 water quality database of 120 lakes were compared to calculated percentiles from a more current and expanded 1980-1999 water quality database of 230 lakes. It was found that the percentiles from the expanded database were very similar to those determined from the 1980-1988 database. For this reason, and in an attempt to maintain commonality, the original 1980-1988 percentiles are continued to be used for lake quality grading purposes.

The three variables used in the grading system strongly relate to open-water nuisance-aspects of a lake (i.e. algal blooms), which can indicate accelerated aging (cultural eutrophication). For example, lake phosphorus concentration has been related to increased algal abundance, increased frequency of algal blooms, and to the increased abundance of blue-green algae (Osgood 1988b). Chlorophyll-a, which is a pigment in plants (including algae) essential in the photosynthesis process, is used to estimate the algal abundance of a lake. And finally, Secchi transparency relates to the appearance of a lake (generally the fewer algae, the better the transparency of a lake). TKN concentration was not included in the grading process because most lake nuisances in the area are related to the phosphorus concentration of the lake (Osgood 1988b).

These water quality grades however, only characterize the open-water quality of lakes. Other nuisances, such as the abundance of aquatic macrophytes, are not indicated with these grades.

The percentile curve can be used to assign individual TP, CLA and Secchi grades to the monitored lakes. Therefore, a lake having a mean summertime Secchi transparency of 1.7 m would receive a “C” grade, or is considered average compared to other area lakes. Overall lake water quality grades were determined by averaging the individual grades. Grades will generally correspond to descriptive rankings and recreational-use impairments of lakes. Lakes receiving an “A” (<10-percentile) can be deemed exceptional as compared to other area lakes and as having no recreational use impairments. A “B” grade lake is considered to have very good water quality and some recreational use impairment, while lakes receiving a “C” are considered to have average water quality and are recreationally impaired. A “D” grade lake translates to a very poor ranking (severely impaired), and a lake receiving a grade of “F” would mean extremely poor quality compared to other area lakes and indicates no possible recreational use.

The report card for lakes sampled by Metropolitan Council-staff in 2005 is presented below. Grades for CAMP-monitored lakes will be addressed later in this report. The grades are based on all data from past studies, so that the grade represents an overall characterization. Pluses and minuses are assigned to indicate apparent trends, either improvement (+) (e.g., Lake McCarrons) or degradation (-) in the quality of the lake.







2005 LAKE QUALITY REPORT CARD

Bald Eagle (Site-1)	D	McCarrons	A+
Bald Eagle (Site-2)	C	O'Dowd	C
Cedar (Scott Co.)	C	Reitz	C
Forest (East Basin)	C	Thole	D
Forest (Middle Basin)	C	Whaletail (Site-1)	D
McMahon	D	Whaletail (Site-2)	C

Bald Eagle Lake, Site 1 White Bear Twp., Ramsey Co.

Lake ID: 620002
WD: Rice Creek

● Sampling site
Contours in meters



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	B	C		C	C		B		B	C	C	C	B
Chlorophyll a	B	B		B	C		C		C	C	C	C	C
Secchi Depth	B	C		C	C		C	C	C	C	B	C	B
Overall	B	C		C	C		C		C	C	C	C	C

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004 Site 1	2004 Site 2
Total Phosphorus	C	C	C	C	B	B	B	B	B			C	C
Chlorophyll a	B	C	C	C	C	C	C	C	C			C	C
Secchi Depth	B	C	B	C	C	C	C	C	B	B	C	C	B
Overall	B	C	C	C	C	C	C	C	D			C	C







Year	2005 Site 1	2005 Site 2
Total Phosphorus	B	B
Chlorophyll a	B	C
Secchi Depth	C	C
Overall	D	C

Source: Metropolitan Council and STOR ET data

Bald Eagle Lake, Site 2 **White Bear Twp., Ramsey Co.**

Lake ID: 620002
WD: Rice Creek

● Sampling site
Contours in meters



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	B	C	C	C	C	B	B	B	C	C	C	C	B
Chlorophyll a	B	B	C	C	C	C	C	C	C	C	C	C	C
Secchi Depth	B	C	C	C	C	C	C	C	C	C	B	C	B
Overall	B	C	C	C	C	C	C	C	C	C	C	C	C

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004 Site 1	2004 Site 2
Total Phosphorus	C	C	C	C	B	B	B	B	B	B	C	C	C
Chlorophyll a	B	C	C	C	C	C	C	C	C	C	C	C	C
Secchi Depth	B	C	B	C	C	C	C	C	B	B	C	C	B
Overall	B	C	C	C	C	C	C	C	D	D	C	C	C

Year	2005 Site 1	2005 Site 2
Total Phosphorus	B	B
Chlorophyll a	B	C
Secchi Depth	C	C
Overall	D	C








Source: Metropolitan Council and ST0 RET data

Cedar Lake Cedar Lake Twp./Helena Twp., Scott Co.

LAKE ID: 700091
WMD: Scott County

● Sampling site

Contours in meters









Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	F	F			F								
Chlorophyll a	F	D			D						D		
Secchi Depth	C	C	C	C	C	C				F	D	D	D
Overall	F	D			D								

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	F					F			F				D
Chlorophyll a	C					D			F				C
Secchi Depth	C					D			D				C
Overall	D					D			F				C

Source: Metropolitan Council and STORET data









Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C				C		D	C		B		B	
Chlorophyll a	D				C		C			B	B	C	
Secchi Depth	C				C		C	C	C	C	C	C	
Overall	C				C		C			B		C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C			C						B			C
Chlorophyll a	B			B						B			C
Secchi Depth	C			C						C			C
Overall	C			C						B			C

Source: Metropolitan Council and STORET data



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					C		C	C	C	B		C	
Chlorophyll a					C		C		C	B	B	B	
Secchi Depth					C		C	C	C	C	C	C	
Overall					C		C		C	B		C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C		B							A			C
Chlorophyll a	B		B							B			C
Secchi Depth	C		C							C			C
Overall	C		B							B			C






Source: Metropolitan Council and STORET data

Lake McCarrons

Roseville, Ramsey Co.

Lake ID: 620054
WD: Capitol Region

◆ Sampling site
Contours in meters




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					C	C	B	C	C	C		C	
Chlorophyll a					B	B	B	C	C	B	B	C	
Secchi Depth					B	C	B	C	C	C	A	C	
Overall					B	C	B	C	C	C		C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C	C	D	D			C	C	C	C		C	A
Chlorophyll a	B	B	C	B			B	C	B	C		B	A
Secchi Depth	C	B	C	C			C	B	A	C		C	A
Overall	C	B	C	C			C	C	B	C		C	A

Source: Metropolitan Council and STORET data









McMahon Lake

Spring Lake Twp.,
Scott Co.

Lake ID: 700050
WMO: Scott County

● Sampling site

Contours in meters










Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	F				D								
Chlorophyll a	F				D								
Secchi Depth	C				D								
Overall	D				D								

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			D			D			D				D
Chlorophyll a			D			D			D				F
Secchi Depth			C			D			D				D
Overall			D			D			D				D

Source: Metropolitan Council and STORET data

LAKE ID: 700095
WMO: Scott County








Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					C								
Chlorophyll <i>a</i>					C								
Secchi Depth					C								
Overall					C								

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		C			C			C		D			C
Chlorophyll <i>a</i>		D			C			C		D			D
Secchi Depth		C			C			C		C			C
Overall		C			C			C		D			C

Source : Metropolitan Council and STO RET data

Reitz Lake Laketown Twp., Carver Co.

Lake ID: 100052
WMD: Carver County




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus						D						D	
Chlorophyll a						F						D	
Secchi Depth						D						C	
Overall						D						D	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	D						C	C	D	D	D	D	D
Chlorophyll a	C						B	C	D	C	D	C	C
Secchi Depth	D						C	C	F	C	B	C	C
Overall	D						C	C	D	C	C	C	C






Source: Metropolitan Council and STD RET data



Thole Lake Louisville Twp., Scott Co.

LAKE ID: 700120
WD: Scott County

● Sampling site
Contours in meters




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					D								
Chlorophyll a					D								
Secchi Depth					D								
Overall					D								






Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	F				D			D		D			D
Chlorophyll a	D				C			D		D			F
Secchi Depth	D				C			C		D			C
Overall	D				C			D		D			D

Source: Metropolitan Council and STO RET data



Whaletail Lake, Site 1

Minnetrista, Hennepin Co.




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	D					C							
Chlorophyll a	D					D							
Secchi Depth	D					D							
Overall	D					D							

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004 Site 1	2004 Site 2
Total Phosphorus						C			C			D	C
Chlorophyll a						C			C			C	C
Secchi Depth						D			D			F	D
Overall						C			C			D	C

Year	2005 Site 1	2005 Site 2
Total Phosphorus	D	C
Chlorophyll a	C	C
Secchi Depth	D	D
Overall	D	C

Source: Metropolitan Council and STO RET data





Whaletail Lake, Site 2

Minnetrista, Hennepin Co.

Lake ID: 270184-02
WMO: Pioneer-Sarah Creek

◆ Sampling sites

Contours in meters







Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	D					C							
Chlorophyll a	D					D							
Secchi Depth	D					D							
Overall	D					D							

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004 Site 1	2004 Site 2
Total Phosphorus						C			C			D	C
Chlorophyll a						C			C			C	C
Secchi Depth						D			D			F	D
Overall						C			C			D	C

Year	2005 Site 1	2005 Site 2
Total Phosphorus	D	C
Chlorophyll a	C	C
Secchi Depth	D	D
Overall	D	C

Source: Metropolitan Council and STD RET data



PART II - CITIZEN-ASSISTED LAKE MONITORING

ACKNOWLEDGMENTS

The success of the 2005 volunteer lake monitoring program would not have been possible without the greatly appreciated work done by volunteer monitors, and the support of the organizations that enrolled lakes in the program.

The enrolling organizations, which included 16 watershed management organizations/watershed districts (WMO/WD), 12 cities, two counties, one environmental group, one basin planning group, and one lake association were involved in volunteer recruitment, training, and occasional follow up on the progress of their volunteer lake monitors. Without this help, the program would not have been as successful as it was.

However, those deserving the greatest appreciation, are the volunteers themselves. Their help has made this program successful. The list of the volunteers involved in the 2005 Citizen-Assisted Monitoring Program (CAMP) is shown in Appendix B. The Metropolitan Council and local WMO/WDs thank them for the sustained efforts contributed over six months and the quality of their work.

INTRODUCTION

Volunteer monitoring is a growing endeavor around the country. Citizens are finding that good information on the status of local water quality and the causes of water quality degradation is often not available from scientific research projects or government surveys. Therefore, the citizens themselves are collecting this information.

As is the case throughout the United States, the majority of lakes in the Twin Cities Metropolitan Area (TCMA) suffer from this lack of water quality data. Area lakes and watershed managers need a broad, comprehensive water quality database for regulatory and decision-making purposes. Because of the lack of public funding and the large ratio of area lakes to monitoring staff, very little data exist for the majority of the lakes in the area, and local decision-makers are forced to make management decisions lacking adequate information.

CAMP was initiated by the Metropolitan Council in 1993 to help bridge the data gaps for area lakes, provide a more complete and improved Metro database, give local decision makers a better idea of the water quality in the area, and assist them in decision making on water quality issues. The Council's goal for CAMP is to provide a means to gather as much information on area lakes, as is economically possible.

Previous volunteer programs conducted throughout the United States have shown that with proper equipment and instructions, volunteers can be trained to produce credible water quality data. Because most of the volunteers live near the lakes they are monitoring, they are very interested in determining any trends and/or changes in local water quality (Nichols 1992).

Not only does volunteer involvement in the lake monitoring process substantially reduce the cost of obtaining data, but it enhances the grass-root understanding of how lakes work and how certain lake conditions relate to the surrounding watershed.

PURPOSE OF THE VOLUNTEER PROGRAM

The main purpose of CAMP is to provide lake and watershed managers with water quality data that will not only support them in properly managing the resources, but also provide much needed historic baseline data to help document water quality impacts. As noted earlier, an additional function of the monitoring program is the volunteer's increased awareness of their lake's condition and workings throughout the summer, which may foster grass-roots initiatives to protect lakes and promote support for lake management.

CAMP involved the collection of in-lake samples by volunteers. Monitoring procedures and sample handling methods were determined through a pilot study during the summer of 1991. The pilot study was designed to evaluate the validity of data collected using several possible citizen monitoring and sample handling methods by comparing them to routine methods (Hartsoe and Osgood 1991). The pilot study and results are presented in Appendix D of the Council's 1993 lake monitoring report (Anhorn 1994) and can be obtained by contacting Randy Anhorn at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Volunteers collected surface water samples that were analyzed for total phosphorus (TP), total Kjeldahl nitrogen (TKN), and chlorophyll-a (CLA) [a select few of the lakes collected samples to be analyzed for chloride as well]. In addition, they measured surface water temperature, water transparency, and recorded user perceptions (some monitors also recorded dissolved oxygen). Most lakes were visited biweekly from April through October (fourteen sampling dates) and were sampled at the lake's deepest open-water location. Quite a few of the lakes, however, were not monitored each of the desired 14 sampling weeks. The reasons for the missed sampling dates varied. However, the majority of the lakes, even with the missed sampling dates, were sampled adequately and often enough to provide an annual overview of the water quality of each lake. Samples were submitted to Council-staff and then forwarded to the MCES-EPE laboratory.

CAMP METHODS

OBTAINING VOLUNTEERS

Active recruitment for lakes and interested volunteers for the 2005 volunteer monitoring program began in the winter months of 2004. Letters and registration forms were sent to various WMOs, counties, and cities to determine their interest in enrolling lakes within their jurisdiction in the program. The organizations were then encouraged to obtain volunteers for each lake they enrolled in the program. If there were problems finding willing volunteers the Council assisted in the search; however, the belief was that the supervising organization would benefit in the long run by having direct contact in recruiting its volunteers. This contact would hopefully open a two-way communication line between concerned citizens and the WMOs.

The year 2005 marked the thirteenth year of the Council's volunteer program. Sixteen watershed management organizations/watershed districts (WMO/WD), 12 cities, two counties, one environmental group, one basin planning group, and one lake association participated in CAMP in 2005, enrolling a total of 160 lakes. This year's volunteer-monitoring program included 17 lake sites never before monitored by the Council and 129 lake sites which were also monitored in 2004. A map indicating the 2005 CAMP lakes and their affiliated enrolling entity is shown in Figure 2, while a list of the volunteer monitors for each lake is provided in Appendix B.

TRAINING VOLUNTEERS

Volunteer training was conducted by Council-staff at various locations throughout the seven- county metropolitan area. Volunteer training was scheduled between late-February and early-April 2004. At each training session, volunteers were given a handbook describing the program, outlining basics in the biology and ecology of lake systems, and containing detailed written instructions for the lake monitoring and data form completion procedures.

At each training session, volunteers received the necessary equipment for the lake monitoring. This equipment was purchased by the enrolling agency through the Council and loaned to the volunteers. At the end of the year's monitoring season, equipment was returned to the enrolling agency to be used in future years. Each lake's volunteer received:

- Chlorophyll hand pump, flask, and filters
- LCD thermometer
- Map of lake with sampling site(s)
- Sampling observation forms
- Sample jug
- Sample vials and labels
- Secchi disk
- Aluminum foil
- Tweezers (forceps)


During the training session, volunteers were given a brief description of the inner working of a lake system as described in their handbook, instructed on proper lake monitoring procedures, and shown how each piece of sampling equipment worked. After this discussion, the volunteers received a package containing the equipment, and the proper use of each piece of equipment was again described and practiced. Finally, each volunteer was asked to sign a waiver of liability stating that they were not an employee of either the Council or the agency enrolling the lake in the program (i.e. the watershed management organization), and that they would use proper safety equipment and observe boat operating methods specified by the State of Minnesota.

MONITORING METHODS

Volunteers were instructed to monitor their designated lake site(s) on a biweekly basis from mid-April to mid-October. Thus, there were 14 possible sampling periods. The methods they used were determined through a pilot study in 1991 that tested simplified methods for using volunteers to obtain credible water quality data (Anhorn 1994). The monitoring methods are detailed in the following paragraphs.

First, during pre-arranged sampling weeks, volunteers located and anchored their boat at pre-determined monitoring locations (the deep open-water area of the lake). Once at the monitoring location, an observation form for lake and meteorological conditions was completed. The form, shown in Figure 3, provided space to mention natural and cultural observations which may have influenced what was happening in the lake (i.e., heavy rains two days before monitoring), and an area to relate general perceptions of the lake's condition and suitability for recreation.

2005 CAMP Lakes



- | | | | |
|-------------------|-------------------|--------------------------|----------------------|
| 1. Goetschel Pond | 7. North Twin | 13. South School Section | 19. Schroeder's Pond |
| 2. McDonald | 8. Carol | 14. Jellum's Bay | 20. Staples |
| 3. Klawitter | 9. Silver | 15. Fish | 21. Turtle |
| 4. DeMontreville | 10. Herber's Pond | 16. MacDonald Pond | 22. Mud |
| 5. Long | 11. Loon | 17. East Boat | |
| 6. South Twin | 12. Louise | 18. West Boat | |

Next, the volunteers took a water transparency reading by lowering a Secchi disk on the shaded side of the boat to the point at which it disappeared. The point where the disk reappears is the Secchi transparency depth that was recorded on the observation form. The next lake monitoring step involved the collection of the surface water sample.

Collecting a surface water sample. A surface water sample was collected in a clean one-gallon plastic milk jug. To begin, the volunteer pre-rinsed the jug three times with lake water. After rinsing, the jug was filled by submersing it upside down to forearm depth and turning it upright while still submersed. After filling the sample jug, volunteers tested and prepared it for the following parameters:

- **Temperature.** Surface water temperature was measured from the volunteer's sampling jug using a LCD thermometer that is readable to 0.1°C. The temperature was measured immediately following sample collection. Special care was taken to keep the sample out of direct sunlight in order to minimize temperature change.
- **Total Phosphorus (TP) and Total Kjeldahl Nitrogen (TKN).** Two samples, one each for TP and TKN, were decanted from the volunteer's jug in the field into their respective triple pre-rinsed, pre-labeled (including lake name, date, time, and parameter) 50-milliliter (ml) vials. These samples were then placed in the cooler, taken home, and stored in the freezer until they were picked up and delivered to the laboratory for analysis.
- **Chlorophyll-a (CLA).** CLA samples from the volunteer's jug were filtered in the field (*out of direct sunlight*) onto a 0.45 micrometer (μm) glass-fiber filter using a field filtration apparatus and a hand pump. Water from the sampling jug was measured and poured into the pump reservoir using a graduated cylinder. The pump reservoir holds approximately 250 ml. By squeezing the handle of the pump, the sample water was forced through the filter and the suspended planktonic algae became attached to the filter. The filtered water was then dumped back into the lake. If possible, this was repeated until a total of 1000 ml of sample water was allowed to pass through the filter. However, if the water sample was too green and the filter became clogged without allowing more water to pass through, the amount of water that did pass through the filter was calculated and recorded on the observation form. The filters were then removed from the filter holder with tweezers, and placed in a petri dish. The sample container was then labeled using the same methods used on the TP and TN sample vials (except the amount of water pumped through the filter was also included on the label), wrapped in tin foil, and frozen until pick-up and delivery to the lab.

The frozen samples were picked up within approximately 30-60 days by Council-staff and delivered to the MCES-EPE's laboratory for chemical analysis. Results from the 1991 pilot study reveal that the volunteer monitoring and handling methods chosen for use in the CAMP program yield results comparable to routine methods used by the Council (Hartsoe and Osgood 1991).

In addition, a few WMO/WDs had their volunteer(s) record dissolved oxygen (DO) and temperature profiles, as well as collect surface chloride and subsurface TP and CLA samples. Chloride samples were prepared in the field identical to the TP/TKN samples. The WMO/WDs provided their volunteers with supplementary equipment and training to use this equipment, as well as paying for the additional cost of laboratory analysis for the TP samples. The additional profiles, and subsurface samples were picked up by the Council along with the routine samples. Profiles obtained by the volunteers were then mailed to the WMO/WD, and the samples were delivered to the lab for analysis.

Figure 3. Example of Sampling Form

Lake Name and ID #: _____

Site #: _____

Sampling Date: _____

Time: _____

Name(s) of Volunteer(s):

SECCHI DISK DEPTH: _____ meters

SURFACE TEMPERATURE: _____ °C

VOLUME OF FILTERED LAKE WATER (CLA) _____ ml

GENERAL OBSERVATIONS

(Circle)

* Water Color

Clear Yellow
Green Gray
Brown Blue-Green
Comment:

* Odor of Water

None Rotten Egg-like
Fishy Septic-like
Musty _____
Comment:

* Wind Conditions

Calm Strong
Breezy
Direction: _____

* Water Surface

Calm Moderate Waves
Ripple Whitecaps
Small Waves
Comment:

* Cloud Cover

0% 75%
25% 100%
50%

* Lake Level

Above Normal
Normal
Below Normal
Staff Gage Reading _____

* Amount of Aquatic Plants

None Moderate
Minimal Substantial
Slight

* Air Temperature (F)

< 40 81-90
41-60 > 90
61-80

* Unusual Conditions in the
past week (storms, high
winds, temp. extremes):

* Physical Condition

Crystal Clear(1)
Some Algae Present(2)
Definite Algae Present(3)
High Algal Color(4)
Severe Bloom (Odor, Scum)(5)

* Suitability For Recreation

Beautiful(1)
Minor Aesthetic Problem(2)
Swimming..Slightly Impaired(3)
No Swim..Boating OK(4)
No Aesthetics Possible(5)

DATA HANDLING AND ANALYSIS

Once each lake's sampling forms and lab analyses were delivered to the Council, the data were entered into a data management and statistical analysis program called Statistical Analysis System (SAS). This data handling system served three purposes:

1. Check-in of forms and tracking of volunteer participation;
2. Entry of nutrient, Secchi, and user perception data into a database for statistical, graphical, and tabular outputs; and
3. Entry into the U.S. Environmental Protection Agency's (U.S. EPA) national water quality data bank called STORET.

If there were questions concerning the data and/or lake observations, the volunteer was called by the Council-staff. The Council maintained contact with most volunteers throughout the season by telephone or in person during sample pick-up. Statistical analyses were performed, and tables and plots of the data were prepared.

PROGRAM QUALITY ASSURANCE/QUALITY CONTROL

The quality assurance/quality control (QA/QC) objective for CAMP is to prevent erroneous data from being produced and used. If by chance errors did occur, they were identified and corrected. Additionally, all suspect data were excluded in lake databases or conclusions.

The MCES-EPE's laboratory follows its own internal QA/QC program. The MCES-EPE lab uses an extensive internal and external check and balance system to ensure credible data. Documentation of the lab's QA/QC procedures can be obtained through the MCES-EPE.

To ensure that CAMP volunteer monitors were using proper sampling techniques and producing credible data, two QA/QC methods were used. Either Council-staff accompanied a volunteer on a sampling event to oversee their collection and preparation procedures, or staff monitored a CAMP lake site during the same week (although not necessarily the same day) that volunteers were to sample the lake site. The first method was used to simply observe the monitor's methods to determine if there were any problems that needed to be addressed. This procedure was usually undertaken when Council-staff was in a volunteer's area on a known sampling day, or when it seemed necessary.

The most common quality check method, however, involved monitoring of the lake by the Council during a scheduled monitoring week. For these sampling events, Council-staff used the same type of equipment and same methods as the volunteers. The Council-collected QA/QC samples were then treated just as the volunteer samples were so that the nutrient concentrations and Secchi transparencies of both sampling events could be compared to determine if any procedural problems existed. If there seemed to be discrepancies, Council-staff would accompany the volunteer on their next sampling event to observe their methods and, if necessary, re-train them. Data determined to be erroneous were thrown out of the database.

During the 2005-monitoring season, 40 percent of the CAMP lake sites monitored more than three times throughout the summer were monitored by Council staff during scheduled monitoring weeks to determine the credibility of the volunteer data. Many of the lakes that were 'checked' by Council-staff in 2005 were

monitored by volunteers (and ‘checked’) as part of past CAMP monitoring years. Council-sampled QA/QC measurements are presented along with volunteer samples in each lake’s descriptive section. A regression analysis was performed on the QA/QC dataset to determine if a statistically significant difference was found between the volunteer and professionally collected data. The resulting statistical analysis of the quality check data revealed excellent agreement between volunteer and professionally-collected samples.

Regression analysis. The 2005 and 1993-2005 QA/QC volunteer- and professional-collected TP, CLA and Secchi data were plotted on a scatterplot graph (Figures 4-9). A linear regression (shown on the graph as a solid line) was run on the resulting data. If the professional- (y) and volunteer-collected (x) data were identical, the data points would fall along the dashed line shown on the following graphs ($x=y$).

The graphs show that while the majority of the data points do not fall exactly on the $x=y$ -line, they do, for the most part, fit the $x=y$ -line well. The graphs also show that while the regression-lines for each parameter are nearly identical to the $x=y$ -lines when the tested parameters are low, the regression-line begins to fall away from the $x=y$ -line as the parameter levels increase. Because of the close fit of the regression-line to the $x=y$ -line and because of the strong linear relationships of each parameters data (shown as a large R^2), it is determined that there is no statistically significant difference found between samples collected by volunteers and those collected by Council staff.

Figure 4. 2005 CAMP-collected TP vs. professionally-collected "quality check" TP




Figure 5. 2005 CAMP-collected CLA vs. professionally-collected "quality check" CLA








Figure 6. 2005 CAMP-collected Secchi transparency vs. professionally-collected "quality check" Secchi transparency





RESULTS AND ANALYSIS

The water quality of the CAMP lakes will be discussed on a lake-by-lake basis in the following pages. *The Handbook for the Citizen-Assisted Lake Monitoring Program* (Anhorn 93), handed out at the volunteer training sessions, overviews the basic inner workings of lakes.

The results and subsequent analysis of the water quality of each lake includes a written section describing the lake's current condition as determined through the 2005 CAMP monitoring and a separate lake information sheet. Each information sheet includes current 2005 water quality data, shown in both tabular and graphic form, and all 1980-to-the-present lake water quality grades (the methodology and percentile ranges of the grading system were discussed in Part I of this report). To determine any water quality trends (i.e., whether the lake quality is improving, degrading, staying the same, or has no trend) each lake's 1980-to-the-present database was used.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Alimagnet Lake (19-0021) *City of Apple Valley*

Approximately half of Lake Alimagnet's 109-acre surface area is located within the City of Apple Valley, the other half in the City of Burnsville (Dakota County). The lake's shoreline is 3.2 miles. The lake has maximum and mean depths of 3.0 and 1.5 m (10 and five feet), respectively. Because the lake is relatively shallow, it does not develop and maintain a thermocline (a density gradient owed to changing water temperatures throughout the water column), and the entire lake is considered littoral, (the shallow [0-15 feet] area dominated by aquatic plants). The approximate lake volume is 545 acre-feet (ac-ft). The lake has a 1,094-acre watershed and a watershed-to-lake area ratio of 10:1 (Blue Water Science 2005). The greater the ratio, the greater the potential stress on the lake from surface runoff.

There are 12 inlets into the lake. A 1990 Clean Water Partnership Diagnostic-Feasibility Study on the lake estimated land use for the watershed at: 29 percent single-family residential, eight percent multi-family residential, three percent commercial/industrial, 19 percent wooded, 10 percent open waters/wetlands, and 31 percent open/undeveloped (Montgomery Watson 1990). Land use percentages have no doubt continued to shift from open/undeveloped to urban uses (single-family residential, multi-family residential, and commercial/industrial) since that study.

The lake, which has been monitored through CAMP since 1995, was sampled 10 times between early-May and early-October, 2005.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	120.0	76.0	177.0	D
CLA (µg/l)	51.5	6.7	100.0	D
Secchi (m)	0.6	0.3	0.8	F
TKN (mg/l)	1.96	1.10	3.90	
Overall Grade				D

The 2005 overall grade is similar to that of 1990, 1996, and 1999-2004 and worse than those recorded in 1995, and 1997-1998. The lake's historic overall water quality grades indicate that the lake fluctuates between a C and D. Most recently the lake's overall grade has consistently been D (1999-2005). The lake's water quality was at its best in 1995, 1997, and 1998 (overall grade of C). The lake's 2005 summertime TP, CLA, and Secchi means were similar to those recorded in 1999-2004 (which represent some of the lake's worst water quality).

In an attempt to reduce algal blooms and improve the lake's water quality, crushed cornmeal was used in 2005 as an in-lake organic carbon amendment. A recent study on Valley Lake-Lakeville, Minnesota (discussed later in Valley Lake section of this report), has suggested that carbon from the decaying barley straw inhibits algal populations via microbial competition for phosphorus (McComas and Anhorn 2004). The use of the cornmeal, however, did not result in the anticipated improvements in Alimagnet's water quality (McComas 2005). The lack of improvement was determined by comparing the 2005 CLA and Secchi transparency means to those of recent "non-crushed cornmeal" years. One potential factor limiting the beneficial effects of the organic carbon amendment could be the result of recent partial winterkills creating an unbalanced fish population (bluegill and bullhead population densities greater than the norm). It is speculated that the lake's large bluegill and bullhead population, could adversely be impacting the water quality and negating the benefits of the carbon amendment (McComas 2005).

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The summertime mean physical condition was 3.6 on a 1-to-5 scale shown on the lake information sheet (between 3- "definite algae present" and 4-"high algal color"). The mean suitability for recreation ranking, also on a 1-to-5 scale, was 3.0 (3- "swimming slightly impaired").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Alimagnet Lake **Apple Valley/Burnsville, Dakota Co.**




Lake ID: 190021
WMO: Vermillion River
Volunteer: John Ritter

◆ Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 thru 5	1 thru 5
5/8/05	14.6				42	119		0.8	3	3
5/18/05	15.2				29	76		0.8		
5/23/05	15.7				33	78		0.8	4	3
6/10/05	24.08				100	78		0.8	4	3
6/30/05	23				12.3	17.4		0.4		
7/5/05	23				6.7	140		0.3	4	3
8/7/05	27.4				38	102		0.5	3	3
8/28/05	24.1				59	136		0.5	3	3
9/16/05	20.7				82	177		0.3	4	3
10/9/05	13.4				130	154		0.3	4	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	F	D									F		
Chlorophyll a											D		
Secchi Depth	F	F	D	D	C	D	F	F	F	F	D	C	D
Overall											D		

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			D	D	C	D	F	D	D	D	D	D	D
Chlorophyll a			B	C	C	C	D	D	C	C	C	D	D
Secchi Depth	C	C	C	D	C	C	D	F	D	F	F	F	F
Overall			C	D	C	C	D	D	D	D	D	D	D

Source: Metropolitan Council and STORET data



Armstrong Lake (82-0116-02) South Washington Watershed District

Armstrong Lake has been annually monitored through CAMP since 1998. There is very little physical information available on the lake or the lake's watershed. Located partially within the cities of Lake Elmo and Oakdale (Washington County), the 39-acre lake has a mean and maximum depth of 1.0 m (3.2 feet) and 1.5 m (roughly 5 feet), respectively. Because of the shallowness of the lake, its entire area is considered littoral (the shallow [0-15 foot depth] area dominated by aquatic vegetation), and it never maintains a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column) through the summer months. The lake's surface area and mean depth translate to a volume of roughly 128 ac-ft. There is no public access to the lake.

Armstrong Lake was monitored 14 times between mid-April and mid-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	52.6	25.0	81.0	C
CLA (µg/l)	8.2	3.9	.0	A
Secchi (m)	1.0	0.9	1.4	D
TKN (mg/l)	0.88	0.82	2.00	
Overall Grade				C

The lake's 2005 overall water quality grade was identical to that of 2000, 2002-2004, and better than those recorded in 1998-1999 and 2001 (D). While the lake's 2004 and 2005 parameter means recorded are the best recorded to date. The main reason for the lake's improvement was the reduction in mean chlorophyll concentration as compared to previous years.

By comparing the lake's historic database TP (nutrient), CLA (algal biomass estimator), and Secchi (water clarity) grades, it is apparent that the TP and Secchi grades are quite a bit worse than the CLA grade. In a most cases, the three should be fairly comparable. One possible explanation for the lake's recent findings may be that the majority of the lake's TP comes from either in-lake suspended sediments (re-suspension), or the intrusion of sediment-laden runoff to the lake, which in turn lessens the clarity of the water and inhibits algal growth.

Statistical analysis of the lake's water quality database failed to produce any statistically significant long-term trends. To better understand the lake's current water quality condition, and which direction it may be heading, continued monitoring is suggested. In the short-term, however, the lake's quality seems best described by a high D/low C grade.


Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The mean physical condition ranking was 2.6 (ranking between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 4.2 (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

If you detect any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Armstrong Lake

Lake Elmo/Oakdale,
Washington Co.,
Washington Co.

LAKE ID: 820116
WCD: South Washington
Volunteer: Washington Co.
SWCD







2005 Data

Date	Surf Temp	Bot Temp	Surf DO	Bot DO	Chl a	Surf TP	Bot TP	Secchi	PC	RS
4/1/05	13.5	13.5	4.78	4.57	1.5	58	58	0.9	2	4
4/22/05	15.1	15.1	5.32	5.04	9.9	51	51	1.4	3	4
5/9/05	17.8	17.6	5.69	5.32	9.6	62	62	0.9	3	4
5/23/05	16.2	15.9	5.3	6.04	6.9	31	31	1.4	2	4
6/7/05	26.3	22.7	5.91	6.34	6	25	25	1.1	2	4
6/20/05	24.2	23.5	6.46	2.64	4.4	62	62	0.9	2	4
7/5/05	21.3	20.5	2.62	0.16	3.9	72	72	0.9	2	4
7/18/05	25.1	25.1	4.5	2.78	9.9	81	81	0.9	3	5
8/3/05	26.2	24.1	5	0.46	11	40	40	1.1	3	4
8/17/05	24.1	23.9	8.84	7.42	10	44	44	0.9	3	4
8/29/05	24	20.5	4.7	2.04	8.2	63	63	0.9	3	5
9/12/05	22.6	22.6	4.79	4.53	9.1	41	41	1.2	3	4
9/30/05	14.6	14.4	8.48	8.08	11	58	58	1.2	3	4
10/13/05	14.1	14	7.49	7.2	11	93	93	1.1	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													
Year	1983	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Source: Metropolitan Council and STORET data



Barker Lake (82-0076) *Carnelian - Marine Watershed District*

Barker Lake is a 45-acre lake located within May Township (Washington County). The mean and maximum depth of the lake is 4.4 m (14 feet) and 9.0 m (roughly 29 feet), respectively. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). Additionally, the surface area and mean depth of the lake result in a calculated volume of 648 ac-ft.). The lake has an 823-acre watershed and a rather large watershed-to-lake area ratio of 19:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This marks the sixth year in which Barker Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake revealed a limited amount of data (1997-2004) collected over the past twenty years.

The lake's Secchi transparency was monitored seven times from late-April to early-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

Water samples to be analyzed for TP, TKN and chlorophyll were not collected for the lake in 2005. Because Secchi transparency was the only data collected there are no nutrient or chlorophyll concentration means to compare to previous years. The lake's 2005 summertime (May through September) mean Secchi transparency was 1.16 m (minimum of 0.90 m and a maximum of 2.00 m). This translates to a grade of D for water clarity (worse than the water clarity grades recorded in 1998-2004).

Statistical analysis on the lake's water quality database did not detect any long-term trends. To better understand the lake's current water quality and in which direction it may be heading, continued monitoring is suggested.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.1 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 3.6 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Barker Lake




Hugo, May Twp., Washington Co.

LAKE ID: 820076

WD: Carleton-Marine


Volunteer: Washington Co.

SWCD



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/21/05	14.9	6.4	4.97	0.57				2.0	2	3
5/19/05	13.3	9.4	5.21	1.74				1.5	3	4
6/14/05	24.2	9.7	7.28	0.57				1.4	2	3
7/13/05	30.6	19.1	9.55	0.05				1.2	4	4
8/12/05	24.1	10.9	5.65	0.45				0.8	4	4
9/9/05	21.4	10.4	8.87	0.49				0.9	4	4
10/6/05	15.7	10.9	6.13	0.49				0.9	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				C	D	D	C	D					
Chlorophyll a				C	C	D	B	C					
Secchi Depth				D	C	C	C	C	C	C	C	C	D
Overall				C	C	D	C	C					

Source: Metropolitan Council and STORET data



Bass Lake (27-0098) Shingle Creek Watershed Management Commission

Bass Lake is located within the City of Plymouth (Hennepin County). The lake covers an area of 194 acres and has a maximum and mean depth of 9.4 m (roughly 31 feet) and 2.9 m (9.5 feet). About 82 percent of the lake's area is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation. The approximate volume of the lake is 1,640 acre-feet (ac-ft) and its approximate residence time (the amount of time required to completely replace the lake's current volume of water with an equal volume of "new" water) is 0.7 years. The lake's watershed of 3,100 acres translates to a rather large watershed-to-lake size ratio of 16:1. The larger the ratio the greater the potential stress put on the lake from surface runoff. 1990 land use estimates indicate that approximately 23.1 % of the watershed is single family residential, 1.2 % is commercial/retail, 0.4 % is industrial/manufacturing, 13.0 % is public waters/wetlands, and 62.3 % is available for potential growth (Montgomery Watson 1994).

Additionally, the lake is considered a "Priority Lake" by the Metropolitan Council, due to its multi-recreational uses. Primary management concerns in the past have revolved around the lake's sizable aquatic macrophyte population and periods of low oxygen levels.

Bass Lake, which was also monitored through CAMP in 1994, 1997, 1999, 2001, and 2003, was monitored 15 times from mid-April to late-October, 2005.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	65.9	16.0	98.0	C
CLA (µg/l)	48.5	4.3	120.0	D
Secchi (m)	1.4	0.5	3.4	C
TKN (mg/l)	1.96	0.73	3.20	
Overall Grade				C

The water quality database for Bass Lake contains six years of CAMP data collection (1994, 1997, 1999, 2001, 2003, and 2005). The lake has received identical overall grades in all six years of CAMP monitoring, 2003 represents the lakes worst monitored water quality year. The best-monitored water quality for the lake was recorded in 1997 (TP= 43.8 µg/l, CLA = 21.0 µg/l, and Secchi = 1.8 m)

While the limited nature of the lake's water quality database makes any statistically significant long-term trend detection impossible, on the short-term the lake seems to consistently have water quality that is representative of a lake grade of C. The last two years of data (2001, 2003, and 2005), however, have shown a slight decrease in water quality over that in recorded the 1990's. This is especially shown in the increase in summer mean total phosphorus and chlorophyll-a concentrations.

The summertime mean physical condition was ranked 3.2 on a 1-to-5 scale shown on the lake information sheet (between 3- "definite algae present" and 4- "high algal color"). The mean suitability for recreation ranking, also on a 1-to-5 scale, was 3.0 (3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Bass Lake Plymouth, Hennepin Co.


LAKE ID: 270098

WMO: Shingle Creek

Volunteer: Marvin Groth

● Sampling site

Contours in meters



2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/21/05	16				52	33		2.7	1	1
4/29/05	11.6				4.5	29		2.7	1	1
5/10/05	15.8				4.4	26		3.4	1	1
5/26/05	16.5				6	24		2.7	1	1
6/2/05	23.2				4.3	16		3.1	2	2
6/15/05	23.4				7.4	30		2.0	2	2
7/1/05	22.3				68	92		1.1	4	3
7/8/05	27				120	98		0.5	5	4
7/19/05	26.2				74	87		0.8	4	4
8/4/05	27.3				79	92		0.6	4	4
8/15/05	25.6				96	89		0.6	4	4
8/30/05	23.7				61	71		0.6	4	4
9/14/05	22.2				58	78		1.1	4	4
9/26/05	19.7				44	88		0.9	3	3
10/26/05	11				33	125		1.4	2	2

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C			C		C		C		C			
Chlorophyll a	C			C		C		C		C			D
Secchi Depth	C			C	D	C		C		C			C
Overall	C			C		C		C		C			C

Source: Metropolitan Council and STORET data



Bass Lake (27-0015) City of St. Louis Park

Bass Lake is a small shallow lake 95-acre lake located within City of St. Louis Park (Hennepin County). There is very little known morphological data available for the lake.

This marks the second year in which Bass Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data revealed only the 2002 CAMP data. Thus, 2002 and 2005 are the only years of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 12 times between late-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	147.1	78.0	199.0	D
CLA (µg/l)	42.2	2.2	200.0	C
Secchi (m)	0.67	0.50	1.00	F
TKN (mg/l)	1.26	0.67	1.90	
<i>Overall Grade</i>				D

Similar to that recorded in 2002, the lake's TP (nutrient), CLA (algal biomass estimator), and Secchi (water clarity) grades do not correspond well to one another. It is apparent that the TP and Secchi grades are quite a bit worse than the CLA grade. In a most cases, the three should be fairly comparable. Possible explanations for the lake's 2002 and 2005 findings may be that the lake the majority of the lake's TP comes from in-lake suspended sediments (re-suspension), or the intrusion of sediment-laden runoff to the lake, which in turn lessens the clarity of the water and inhibits algal growth.

As mentioned earlier, there are no water quality data available for Bass Lake other than the 2002 and 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.0 for physical condition (2- "some algae present"), and 2.7 for recreational suitability (between 2- "minor aesthetic prblem" and 3- "swimming slightly impaired"). The amount of emergent vegetation, however, makes recreating on the lake very difficult.



If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Bass Lake

St. Louis Park, Hennepin Co.




LAKE ID: 270015
 W/D: Minnehaha Creek
 Volunteer:
 Jason Westrum

● Sampling site
 Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/27/05	9.6				30	198		0.2	1	
5/15/05	13.1				5.6	78		1.0	1	
5/27/05	17.2				16	145		0.6	2	2
6/11/05	21.9				2.2	196		0.9	1	1
6/27/05	24.4				19	126		0.5	1	5
7/10/05	26.7				88	193		0.6	2	
7/24/05	29.7					178		0.7	3	
8/2/05	32.4				200	199		0.6	3	
9/4/05	20.5				5.4	82		0.6	2	
9/18/05	25.1				31	127		0.6	3	
10/6/05	14.2				2.3	87		0.8	1	
10/16/05	15.3				7.7	185		0.7	2	




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus										F			D
Chlorophyll a										C			C
Secchi Depth										F			F
Overall										D			D

Source: Metropolitan Council and STORET data



Bass Lake (82-0035) Carnelian - Marine Watershed District

Bass Lake is an 81-acre lake located within May Township (Washington County). The maximum depth of the lake is 4.3 m (roughly 14 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This was the sixth year that Bass Lake was monitored through CAMP. A search through the STORET nationwide water quality database provided a moderate amount of historic data including Secchi data from 1991-2003 and nutrient and CLA data in 1991-1992, 1996-2001, and 2003-2004.

The lake was monitored seven times between mid-April and early-October, 2005. The resulting data and graphs appear on the next page. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	34.6	28.0	43.0	C
CLA (µg/l)	10.8	4.2	17.0	B
Secchi (m)	2.3	2.0	3.4	B
TKN (mg/l)	0.84	0.64	1.10	
Overall Grade				B

The 2005 grade of B is similar to that recorded in 1992 and 2004, and better than the C's recorded in 1991, 1997-2001, and 2003. The 2005 summer means were slightly worse than those recoded in 2004 (which are the lake's best recorded water quality to date).


Statistical analysis on the lake's water quality database did not detect any long-term trends. The lake's water quality seems to be well represented by an overall grade of C+/B-. To better understand the lake's water quality and where it may be heading, more data are needed.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-5-scale. The user perception rankings are shown on the lake's associated information sheet on the following page. The mean summertime physical condition was ranked 2.6 on a (between 2- "some algae present" and 3- "definite algae present"). The mean suitability for recreation ranking, also on a 1-to-5 scale, was 3.2 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Bass Lake

May Twp., Washington Co.



2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
C	C	mg/L	mg/L	ug/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/20/05	16.1	7.7	5.09	1.08	5	36		2.4	2	2
5/19/05	13.8	11.9	5.41	3.25	10	28		2.1	2	3
6/15/05	23.9	17.1	7.09	3.5	4.5	35		3.4	2	3
7/14/05	29.2	20.2	8.55	0.09	8.5	28		2.1	3	3
8/11/05	25.4	22	5.51	0.38	17	43		2.0	3	4
9/8/05	22.8	20.5	8.34	0.79	14	39		2.0	3	3
10/6/05	15.8	16	6.42	6.1	16	34		1.8	2	3

Lake Water Quality Grades Based on Summer Time Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus												C	B
Chlorophyll a												B	B
Secchi Depth												C	C
Overall												C	B

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				C	C	C	C	C	C		C	B	C
Chlorophyll a				C	C	B	B	B	B		B	A	B
Secchi Depth	C	C	C	C	C	C	C	C	C	B	C	B	B
Overall					C	C	C	C	C		C	B	B

Source: Metropolitan Council and STORET data



Bavaria Lake (10-0019) City of Chaska

Bavaria Lake, located in the City of Chaska (Carver County), the 200-acre lake has a mean and maximum depth of 5.6 m (18.4 feet) and 18.3 m (60 feet), respectively. Roughly 65 percent of the lake is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation. Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has been reported on the lake.


The lake's surface area and mean depth translates to an approximate lake volume of 3,674 ac-ft. The lake has a 711-acre immediate watershed, which translates to a watershed-to-lake area ratio of 3.5:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff). A 1999 water quality report on water resources in Carver County estimates land use for the watershed at: 17.5 percent residential, 52.7 percent agricultural, 29.7 percent commercial/industrial, and 0.2 percent open/undeveloped (Carver County Planning 1999). A public access is located on the lake's western edge and because of its multi-recreational uses, it is considered a "Priority Lake" in the Metropolitan Area.

While 2005 was the tenth year that Bavaria has been involved in CAMP, the lake has been monitored by Council staff in the past and has recently been involved in the MPCA's volunteer Secchi transparency program (included in the lake's report card grading system on the following page). Additionally, Lake Bavaria was included within the MPCA's Lake Assessment Program (LAP) in 2001. Through this program additional data, besides in-lake data through CAMP, was collected to help complete a more comprehensive study on the lake.

Lake Bavaria was monitored 18 times between mid-April and mid-October, 2005.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP ($\mu\text{g/l}$)	49.7	30.0	119.0	B
CLA ($\mu\text{g/l}$)	22.9	4.4	50.0	B
Secchi (m)	1.5	0.8	2.0	C
TKN (mg/l)	1.44	0.91	1.90	
Overall Grade				B



Available data for Bavaria Lake reveal that the lake water quality remained constant through the 1980's (C's) and improved through the mid-1990s (overall grades of B in 1994 and 1996, and A in 1997-1998), before falling back to overall grades of B in 1999-2005. The lake's water quality report card shown on the information sheet indicates that the lake has received an overall grade of C during the 1980's, A and B grades throughout the 1990's and early-2000's.

The lake's summer mean graph and report card grades clearly depict that the lake's water quality has recently (mid-1990s to present) started to degrade


Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake's associated information sheet on the following page. The mean physical condition ranking was 2.2 (2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking for the lake was 2.0 (2- "minor aesthetics problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Bavaria Chaska/Laketown Twp., Carver Co.

LAKE ID: 100019
WMO: Carver County
Volunteer: John Ryski



2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/18/05	13.2				49	105		0.9	1	1
5/8/05	12				24	70		1.5	1	1
5/24/05	15				33	119		2.0	2	2
5/25/05	17				34	70		1.8		
5/31/05	18				50	69		0.8	3	3
6/5/05	19					47		1.0	3	3
6/16/05	22				26	34		1.3	3	
6/24/05	25					37		2.0	3	
7/6/05	23				35	37		1.5	2	2
7/24/05	25				19	32		1.5	2	2
8/2/05	25				44	44		1.3	2	2
8/14/05	23				20	37		1.5	2	1
8/21/05	23				18	30		1.5	2	2
8/29/05	21				14	48		1.8	2	2
9/11/05	20				7.5	33		1.5	2	2
9/25/05	18				15	39		1.5	2	2
10/9/05	13				9	66		1.5	1	
10/16/05	13				6.3	74		1.8	1	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus			C			C	C						
Chlorophyll a			C			C	C						
Secchi Depth			C			C	C						
Overall			C			C	C						

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B		C	A	B	B	C	B	B	C	B	C	
Chlorophyll a	A		A	A	A	B	B	B	B	A	B	C	
Secchi Depth	B	B	C	A	A	B	B	B	C	B	C	C	
Overall	B		B	A	A	B	B	B	B	B	B	C	

Source: Metropolitan Council and STORET data



Benton Lake (10-0069) Carver County Environmental Services

Benton Lake is a 115-acre lake located within Benton Township (Carver County). The maximum depth of the lake is 2.0 m (roughly 6.5 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake has a 322-acre immediate watershed, which translates to a watershed-to-lake area ratio of 2.8:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff). A 1999 water quality report on water resources in Carver County estimates land use for the watershed at: 19 percent residential, 55 percent agricultural, 16 percent commercial/industrial, and 10 percent open/undeveloped (Carver County Planning 1999).

This was the fifth year that Benton Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake provided only three years of prior data (collected through CAMP in 1999-2001 and 2003). The lake was monitored 14 times between mid-April and mid-October, 2005. During each monitoring event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as its perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	231.5	115.0	357.0	F
CLA (µg/l)	109.2	37.0	240.0	F
Secchi (m)	0.3	0.1	0.6	F
TKN (mg/l)	4.39	2.00	7.30	
Overall Grade				F

Similar to that recorded from 1999-2001 and 2003, the resulting overall grade for the lakes 2005 water quality was F.

As mentioned earlier, there are no water quality data available for Benton Lake other than the 1999-2001, 2003 and 2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term, however, the lakes water quality is well represented by an overall grade of F. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake's associated information sheet on the following page. The average user perception rankings, on a 1-to-5 scale, were 3.8 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.0 for recreational suitability (4- "no swimming - boating ok").



If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Benton

Cologne, Carver Co.




Lake ID: 100069
WMO: Carver County
Volunteer: Carver Co.
Env. Services

● Sampling site
Contours in meters



2005 Data

Date	Surf Temp	Bot Temp	Surf DO	Bot DO	CLA	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	kg/L	kg/L	kg/L	M	1 to 5	1 to 5
4/15/05	14.5		8.77		130	283		0.4	3	4
4/29/05	8.8		13.1		53	174		0.5	3	4
5/9/05	17.7		9.1		49	135		0.6	3	4
5/25/05	16.7		6.1		70	307		0.4	4	4
6/7/05	24.5	10.72			50	196		0.4	3	4
6/16/05	25.9				37	174		0.4	4	4
7/6/05	25		11.9		130	115		0.3	4	4
7/21/05	27.2				240	161		0.2	4	4
8/3/05	27.8				120	291		0.1	4	4
8/17/05	23.7	13.82			170	256		0.1	4	4
8/30/05	24.1	6.06			96	176		0.1	4	4
9/16/05	20.7	9.2			140	342		0.2	4	4
9/25/05	17.6				99	345		0.2	4	4
10/12/05	12.7		11.24		46	299		0.4	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							F	F	F		F		F
Chlorophyll a							F	F	F		F		F
Secchi Depth				C			F	F	F		F		F
Overall							F	F	F		F		F

Source: Metropolitan Council and STORET data



Benz Lake (82-0120) Browns Creek WMO

Benz Lake is a 36-acre lake located in Grant Township (Washington County) with a maximum depth of approximately 2.7 m (9 feet). Because of the shallowness of the lake, its entire surface area is considered littoral (the shallow [0-15 foot depth] area dominated by aquatic vegetation).

The year 2005 marks the second year that Benz Lake has been involved in CAMP (1998 being the first). A search through the STORET nationwide water quality database provided no additional data other than Secchi transparency data collected in 1998.

On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	186.3	120.0	260.0	F
CLA (µg/l)	94.1	23.0	140.0	F
Secchi (m)	0.6	0.3	0.9	F
TKN (mg/l)	3.09	2.20	4.50	
Overall Grade				F

As mentioned earlier, there is a very limited amount of water quality data available for Benz Lake. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.




Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.7 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.0 for recreational suitability (4- "no swimming - boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Benz Lake


Grant, Washington Co.

Lake ID: 820120
WMO: Browns Creek
Volunteer: Wash. Co.
SWCD



2005 Data

Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	Chl ug/L	Surf. TP ug/L	Bot. TP ug/L	Secchi M	PC 1 to 5	RS 1 to 5
4/11/05	12.8	12.6	7.4	7.37	54	177		0.9	3	4
4/28/05	11.2	11.2	6.34	6.09	22	120		1.2	3	3
5/11/05	11.2	16	6.34	5.86	37	161		0.9	3	4
5/26/05	16.8	16.6	3.93	3.6	23	159		0.9	3	4
6/8/05	23.4	22.8	5.02	3.42	62	260		0.8	4	4
6/23/05	26.6	24.5	9.29	0.42	84	142		0.8	3	3
7/5/05	24	23.5	8.02	4.13	89	170		0.5	5	5
7/18/05	27.9	27.9	5.92	5.35	120	211		0.5	3	4
8/5/05	25.8	25.5	3.88	2.04	140	186		0.3	4	4
8/19/05	23.4	23.3	5.87	3.7	140	190		0.5	4	4
8/31/05	24.8	22.4	11.15	6.23	120	212		0.5	4	4
9/15/05	21.5	21.3	5.26	4.89	100	138		0.5	4	4
9/30/05	17.3	16.5	10.13	8.87	120	260		0.5	4	4
10/14/05	14.3	13.8	9.18	8.03	94	197		0.5	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													F
Chlorophyll a													F
Secchi Depth													F
Overall													F

Source: Metropolitan Council and STORET data



Big Carnelian Lake (82-0049) *Carnelian - Marine Watershed District*

Big Carnelian Lake, located within May Township (Washington County), has a public access on its southwestern side, and is considered a “Priority Lake” due to its multi-recreational uses. The lake covers an area of 455 acres and has a maximum and mean depth of 20 m (roughly 66 feet) and 9.8 m (32 feet). Roughly 28 percent of the lake’s area is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation. The approximate volume of the lake is 14,560 acre-feet (ac-ft). The lake’s watershed of 1,900 acres translates to a rather small watershed-to-lake size ratio of 4:1. The larger the ratio the greater the potential stress put on the lake from surface.

Big Carnelian Lake was monitored 14 times between mid-April and mid-October, 2005. The data and related graphs are presented on the information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	20.0	16.0	24.0	A
CLA (µg/l)	7.4	4.3	19.0	A
Secchi (m)	4.3	2.6	6.1	A
TKN (mg/l)	0.63	0.47	0.83	
Overall Grade				A

The lake received overall grades of A in 1980, 1989, 1991, 1994, 1996-1998, 2000-2002, and 2004-2005, and a grade of B in 1984, 1999, and 2003.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake’s physical and recreational conditions on a 1-to-5 scale. The resulting user perception rankings are shown on the information sheet. The mean physical condition ranking was 1.9 (between 1- “crystal clear” and 2- “some algae present”), while the mean recreational suitability ranking was 1.5 (between 1- crystal clear” and 2- “minor aesthetic problem”).

No statistically significant long-term trend is evident from the lake’s water quality database, in the short-term however, the lake’s quality seems well represented by an overall grade of A.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Big Carnelian Lake May Twp., Washington Co.


LAKE ID: 820049

WD: Carnelian-Marine

Washington Co. SWCD

◆ Sampling site

Contours in meters



2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/13/05	7.4	4.2	6.13	0.18	3	18	4.6	2	1	1
4/25/05	10.7	4.9	6.49	0.12	9.5	17	5.6	1	1	1
5/10/05	13.1	5.2	7.29	0.27	8.4	24	4.7	1	1	1
5/20/05	12.8	5.5	6.15	0.31	6.6	20	4.3	2	1	1
6/6/05	20	6.1	5.36	0.34	4.4	17	6.1	2	2	2
6/21/05	23.7	6.7	8.05	0.66	4.4	17	5.3	2	1	1
7/6/05	24.6	6.9	7.87	0.63	4.6	21	5.8	2	1	1
7/19/05	26.6	6.9	6.86	0.05	19	23	3.8	2	2	2
8/1/05	25.6	7	7.17	0.8	4.3	20	4.7	2	1	1
8/15/05	24.7	7	7.12	1	4.8	16	3.7	2	2	2
8/29/05	23.3	7.1	6.5	1.85	7.4	17	2.6	2	2	2
9/13/05	22.9	7.3	8.03	0.73	8.6	24	2.7	2	2	2
9/29/05	18.7	7.3	7.27	0.88	9	21	3.4	2	1	1
10/10/05	15.8	7.9	7.97	0.68	8.1	31	3.4	2	1	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	A			B						A		A	
Chlorophyll a	A			B						A		A	
Secchi Depth	A			B						A		B	B
Overall	A			B						A		A	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	A		A	A	A	A	A	A	A	B	A	A	A
Chlorophyll a	A		A	A	A	B	A	A	A	A	A	A	A
Secchi Depth	B	B	B	B	A	A	B	A	A	A	B	A	A
Overall	A		A	A	A	B	A	A	A	A	B	A	A

Source: Metropolitan Council and STORET data



Big Comfort Lake (13-0053) *Comfort Lake-Forest Lake Watershed District*

Big Comfort Lake is located just north east of the City of Forest Lake, in Isanti County. This year marked the sixth year that the 219-acre lake has been enrolled in CAMP (1998 [it was, however, only monitored a two times in October] and 2000-2002). The lake has a maximum depth of 14.3 m (47 feet). Roughly 41 percent of the lake's area is considered littoral, the shallow (0-15 foot) depth area dominated by aquatic vegetation.

An indepth lake assessment was undertaken on the lake by the MPCA in 1994.

Big Comfort Lake was monitored 13 times between late-April and late-October, 2005. The data and related graphs are presented on the information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	32.0	18.0	46.0	B
CLA (µg/l)	14.6	8.9	26.0	B
Secchi (m)	2.0	1.2	3.1	C
TKN (mg/l)	1.06	0.58	1.30	
Overall Grade				B

The lakes 2005 overall grade is similar to that recorded in 2001 and better than those recorded in 2000 and 2002-2004.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The resulting user perception rankings are shown on the information sheet. The mean physical condition ranking was 2.8 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.6 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

Statistical analysis on the lake's water quality database did not detect any long-term trends. In the short-term however, the lake seems well represented by an overall grade of C+. To better understand the lake's current water quality and in which direction it may be heading, continued monitoring is suggested.




The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Big Comfort Lake Wyoming Twp., Chisago Co.


Lake ID: 130053
WMO: Comfort Lake - Forest Lake
Volunteer: Washington Co. SWCD

● Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	m	1 to 5	1 to 5
4/14/05	10.1	6.6	8.03	3.4	40	54		1.1	2	2
4/26/05	11	7.3	6.76	0.12	25	42		1.7	3	2
5/13/05	12.2	8.9	5.9	0.23	8.9	46		3.0	2	2
5/24/05	15.8	9.2	4.95	0.05	11	45		2.0	3	2
6/21/05	26.8	11.5	7.42	0.51	9.1	27		2.1	2	2
7/7/05	23.6	11.5	9.63	0.57	22	37		1.8	3	3
7/22/05	27.1	11.5	7.74	0.56	26	39		1.2	3	3
8/2/05	26.4	11.7	6.77	0.53	13	23		1.7	3	2
8/16/05	24.4	11.7	6.65	0.43	13	18		2.0	3	3
9/1/05	22.4	11.7	7.24	0.61	16	22		1.8	3	3
9/14/05	21.7	12	6.67	0.45	16	31		2.3	3	2
9/30/05	17.2	11.9	5.01	0.62	12	30		2.1	3	3
10/11/05	14.6	11.8	5.59	0.36	15	38		1.8	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth								B	B	B			
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	D							C	B	C	C	C	B
Chlorophyll a	B							C	B	C	C	B	B
Secchi Depth	C	C		C	C			C	C	C	C	C	C
Overall	C							C	B	C	C	C	B

Source: Minnesota Pollution Control and STORET data



Big Marine Lake (82-0052) *Carnelian - Marine Watershed District*

Big Marine Lake, located within New Scandia Township (Washington County), has two public accesses, and is considered a “Priority Lake” due to its multi-recreational uses. The lake covers an area of 1,706 acres and has a maximum and mean depth of 15.2 m (roughly 50 feet) and 7.6 m (25 feet). Roughly 67 percent of the lake’s area is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation. The approximate volume of the lake is 42,527 acre-feet (ac-ft). The lake’s watershed of 2,659 acres translates to a small watershed-to-lake size ratio of 1.5:1. The larger the ratio the greater the potential stress put on the lake from surface runoff.

Big Marine Lake was monitored 14 times between mid-April and mid-October, 2005. The data and related graphs are presented on the information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	18.1	12.0	24.0	A
CLA (µg/l)	8.4	3.6	29.0	A
Secchi (m)	3.7	2.4	5.2	A
TKN (mg/l)	0.64	0.45	1.20	
Overall Grade				A

The lake received overall grades of A in 1989, 1994, 1996-1998, 2000-2001, and 2003-2005 and a grades of B in 1980, 1981, 1984, 1991, 1999, and 2002.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake’s physical and recreational conditions on a 1-to-5 scale. The resulting user perception rankings are shown on the information sheet. The mean physical condition ranking was 2.2 (between 2- “some algae present” and 3- “definite algae present”), while the mean recreational suitability ranking was 1.8 (between 1- “beautiful” and 2- “minor aesthetic problem”).

While no statistically significant long-term trend is evident from the lake’s whole water quality database (including TP, CLA and Secchi data), a recent MPCA conducted trend analysis using just the lake’s Secchi transparency data, revealed a statistically significant improvement in recent water clarity. In the short-term, the lake’s quality seems well represented by an overall grade of B+/A.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.


If you notice any errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Big Marine Lake May Twp. and New Scandia Twp., Washington Co.

LAKE ID: 820052

WD: Carleton-Marine

Volunteer: Washington Co. SWCD



Birch Lake (13-0042) *Comfort Lake-Forest Lake Watershed District*

Birch Lake is a 65-acre lake located in southern Chisago County. There is very little other known morphological data available for the lake.

This marks the first year in which Birch Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake came up empty. Therefore, 2005 is the only known year of water quality data available. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	117.0	44.0	299.0	D
CLA (µg/l)	60.6	11.0	290.0	D
Secchi (m)	1.3	0.8	2.0	C
TKN (mg/l)	1.29	0.75	2.10	
<i>Overall Grade</i>				D

Throughout the monitoring period, the volunteers ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The resulting user perception rankings are shown on the information sheet. The mean physical condition ranking was 2.5 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 3.9 (roughly equal to 4- "no swimminmg - boating ok").

Because of the limitedness of the lake's water quality database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Birch Lake

Chisago Lake Twp. and Wyoming Twp.,
Chisago Co.

LAKE ID: 130042

WMO: Comfort Lake-Forest Lake
Volunteer: Washington Co. SWCD



● Sampling site
Contours in meters

2005 Data





Date	Surf Temp	Bot Temp	Surf DO	Bot DO	CLA	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	kg/L	kg/L	kg/L	M	1 to 5	1 to 5
4/18/05	13.9	9.7	4.02	0.32	15	46		2.1	2	4
4/26/05	10.5	10.5	5.6	5.33	7.9	56		2.7	2	4
5/16/05	11.5	11	6.71	5.64	13	64		1.8	2	4
5/24/05	17.7	14.8	4.81	2.08	11	44		2.0	2	4
6/6/05	23.9	16	4.02	0.23	12	88		1.2	2	4
6/22/05	26.2	16.8	9.97	0.42	18	100		1.5	2	3
7/7/05	23.2	18	7.86	0.45	50	108		1.2	2	4
7/18/05	24.5	18.4	4.15	0.08	40	50		1.1	2	4
8/2/05	26.5	19	5.56	0.43	290	299		0.8	3	4
8/16/05	23.1	18.5	3.56	0.31	95	132		1.1	3	4
8/30/05	21.7	20.3	5.8	0.43	70	116		1.1	3	5
9/14/05	20.3	19.1	3.85	0.7	38	167		1.4	3	4
9/29/05	14.8	14.5	6.43	4.47	30	79		1.4	3	3
10/11/05	12.1	10.7	7.12	2.57	15	63		1.8	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													D
Chlorophyll a													D
Secchi Depth													C
Overall													D

Source: Metropolitan Council and STORM data



Bone Lake (82-0054) *Comfort Lake-Forest Lake Watershed District*

Bone Lake was previously monitored as a part of CAMP in 1993, 1995, 1997-1999, and 2001-2004. In 2005, the lake was monitored 14 times between mid-April and mid-October. Results are presented on the information sheet on the following page.

The 212-acre lake is located within New Scandia Township (Washington County). It receives flow through three inlets. The lake has a public access on its northwestern side and has a maximum and mean depth of 9.8 m and 3.7 m (32 and 12 feet), respectively. The approximate lake volume of Bone Lake, which has been stocked with walleye by the MDNR in the 1990's, is 2,820 ac-ft. The lake's 5,177-acre watershed translates to a rather large watershed-to-lake size ratio of 24:1. The greater the ratio, the greater the potential stress on the lake from surface runoff. Roughly 59 percent of the lake is considered littoral zone, that is, the area of aquatic plant dominance. The lake is considered a Metropolitan Council "Priority Lake" due to its multi-recreational uses.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	57.0	26.0	189.0	C
CLA (µg/l)	19.9	7.0	53.0	B
Secchi (m)	2.0	1.1	3.7	C
TKN (mg/l)	1.24	1.10	1.50	
Overall Grade				C

Based on the lake water quality grade, shown on the facing information page, the lake's quality throughout the mid-1980's, 1990's, and early-to-mid 2000's seems to be consistently represented by an overall grade of C.

Throughout the summer, the volunteer(s) ranked the lake's perceived physical and recreational conditions on a 1-to-5 scale (see lake information sheet). The mean rankings were 3.4 for physical condition (between 3- "definite algae present" and 4- high algal color"), and 3.2 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of any erroneous lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Bone Lake New Scandia Twp, Washington Co.

Lake ID: 820054
WMO: Comfort Lake Forest Lake
Volunteer: Washington Co.
SWCD

● Sampling site
Contours in meters



2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	mg/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/1/05	10.4	8.8	6.42	0.17	11	61		2.1	2	2
4/26/05	11.5	11.4	5.75	5.43	11	52		2.0	3	3
5/16/05	12.2	11.8	6.84	6.05	7	58		3.7	4	3
5/24/05	16.9	14.2	5.21	4.75	7.4	43		3.5	4	4
6/9/05	23.2	14	5.34	0.25	18	189		2.0	3	3
6/22/05	25.7	13.9	11.11	0.46	11	55		2.1	2	2
7/7/05	24.1	14.3	9.28	0.48	12	36		1.8	3	3
7/22/05	28	16	7.26	0.46	9.2	26		1.7	3	3
8/2/05	28.1	15.6	7.73	0.38	22	38		1.2	4	4
8/16/05	25.2	15.9	8.54	0.3	27	40		1.1	4	4
8/30/05	23	19.5	8.3	0.45	27	36		1.4	4	3
9/1/05	22	21	7.28	1	53	49		1.4	3	3
10/3/05	18.3	17.4	8.23	0.75	44	48		1.2	3	3
10/11/05	15	14.7	8.14	0.5	27	52		1.7	3	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					D			C	C	C		D	
Chlorophyll a					C			B	C	C		C	
Secchi Depth					C		D	C	D	C	C	C	
Overall					C			C	C	C		C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C				C	C	C		C	C	D	C	C
Chlorophyll a	C				B	B	C		C	C	C	C	B
Secchi Depth	C	D	C		C	C	D		C	D	C	C	C
Overall	C				C	C	C		C	C	C	C	C

Source: Metropolitan Council and STORET data



Brickyard Lake (10-0225) Carver County Environmental Services

Brickyard Lake is a 17-acre lake located near the City of Chaska (Carver County). The maximum depth of the lake is 13.1 m (roughly 43 feet). Thirty-five percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance).

This was the fourth year that Brickyard Lake has been involved in CAMP (2002 being the first). The lake was monitored 14 times between mid-April and mid-October, 2005. During each monitoring event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as its perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	16.1	8.0	35.0	A
CLA (µg/l)	3.1	1.0	7.7	A
Secchi (m)	4.7	2.4	6.9	A
TKN (mg/l)	0.26	1.20	0.48	
Overall Grade				A

To the best of our knowledge, there are no water quality data available for Brickyard Lake other than the 2002-2005 CAMP data. Therefore it is not possible to determine any long-term or trends. In the short-term however, the lake's water quality is well represented by an overall grade of A. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 1.1 for physical condition (between 1- "crystal clear" and 2- "some algae problem"), and 1.0 for recreational suitability (1- "beautiful").



The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Brickyard Lake Chaska, Carver Co.




Lake ID: 100225
WMO: Carver County
Volunteer: Carver Co.
Env. Services

◆ Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/14/05	13.6		11.2		2.8	24		4.1	1	1
4/29/05	11.3		11.4		2.8	16		4.4	1	1
5/10/05	16.4		11.5		3.8	35		3.3	1	1
5/23/05	17.2		9.3		2.9	16		4	1	1
6/8/05	22.1		7.8		1.7	16		3.2	2	1
6/22/05	25.4		8.64		2.8	14		5.7	1	1
7/5/05	24.5		8.82		1	8		4.7	1	1
7/20/05	27.1		7.27		1.1	10		4.9	1	1
8/2/05						12		6.5	1	1
8/16/05	24.6		7.88		2.5	12		6.9	1	1
8/29/05	24.3		8.34		1.7	18		5.9	1	1
9/13/05	23.3		9.24		5	17		2.4	1	1
9/26/05	20.8		16.9		2.3	19		4.7	1	1
10/11/05	15.3		7.93		4.5	59		2.4	1	1




Lake Water Quality Grades Based on Summertime Averages

Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus										A	A	A	A
Chlorophyll a										A	A	A	A
Secchi Depth										A	A	A	A
Overall										A	A	A	A

Source: Metropolitan Council and STORET data



Burandt Lake (10-0084) Carver County Environmental Services

Burandt Lake is a 116-acre lake located near the City of Waconia (Carver County). The mean and maximum depths of the lake are 3.0 m (roughly 10 feet) and 7.3 m (roughly 24 feet). Roughly 57 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance).

This was the fifth year that Burandt Lake has been involved in CAMP (1999-2001 and 2004 being the others). The lake was monitored 22 times between mid-April and early-October, 2005. During each monitoring event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as its perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	49.5	24.0	91.0	C
CLA (µg/l)	20.9	2.9	61.0	C
Secchi (m)	1.5	0.6	3.6	C
TKN (mg/l)	1.16	0.47	1.90	
Overall Grade				C

To the best of our knowledge, there are no water quality data available for Burandt Lake other than the 1999-2001 and 2004-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term however, the lake's water quality is well represented by an overall grade of C. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.7 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 2.6 for recreational suitability (between 2- "minor aesthetic problems" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.




If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Burandt Lake

Waconia Twp, Carver Co.


Lake ID: 100084
WMO: Carver Co.
Volunteer: Don Westerman

● Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	12.8		10.9		18	50		1.5	2	1
4/30/05	11				21	41		1.1	2	2
5/5/05	14.9		12.1		14	47	81		1	2
5/15/05	13				18	32		1.3	2	2
5/25/05	19				18	33		1.8		
5/29/05	21				19	30		2	3	3
6/5/05		9		16	24	37	3.1	2	2	2
6/12/05	25			2.9	30		3.6	2	2	2
6/20/05	24.7	8.33		8.8	38	49	2.6	2	3	3
6/29/05	25			6	36		2.6	2	2	2
7/6/05	23	8.77		11	26	55		3	2	2
7/15/05	31			9.9	72		1.1	2	2	2
7/18/05	26.9	8.55		18	26	245	1.3	3	3	3
7/29/05	27			7.9	58		1.1	2	2	2
8/1/05	25.5			21	43	92	1.4	3	3	3
8/14/05	25			11	56		0.7	4	3	3
8/17/05	24.4	9.79		51	48	191	0.7	3	3	3
8/29/05	22.6	12.6		45	80	94	0.6	3	3	3
9/4/05	24			28	90		0.7	3	2	2
9/14/05	22.5	7.41		61	81	83	0.9	3	3	3
9/18/05	22			27	91		0.6	5	4	4
10/3/05	20			26	102		0.6	3	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	C	C			C	C
Chlorophyll a							C	C	C			C	C
Secchi Depth							D	D	D			C	C
Overall							D	C	C			C	C

Source: Metropolitan Council and STORET data



Campbell Lake (10-0127) Carver County Environmental Services

Campbell Lake is located within Hollywood Township (Carver County). The maximum depth of the 72-acre lake is 2.0 m (roughly six-and-a-half feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This was the fourth year in which Campbell Lake has been involved in CAMP (the others being 1999 [where it was only monitored twice (no grade determined)], 2000, and 2002). Other than the mentioned CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty. Therefore, 1999-2000, 2002 and 2005 are the only years of available data.

The lake was monitored 14 times from mid-April to mid-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.




2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	327.8	172.0	516.0	F
CLA (µg/l)	166.7	55.0	270.0	F
Secchi (m)	0.3	0.2	0.6	F
TKN (mg/l)	3.93	0.46	6.60	
Overall Grade				F

As mentioned earlier, there is no water quality data available for Campbell Lake other than that collected in 1999 (just two data points), 2000, 2002, and now 2005. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, more data are needed.


The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.3 for physical condition (between 3- "definite algal presence" and 4- "high algal color"), and 4.2 for recreational suitability (between 4- "no swimming - boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	13.8		12.5		45	405		0.6	3	4
4/28/05	9.7		11.4		30	194		0.5	2	4
5/9/05	17.2		9.5		55	198		0.6	3	4
5/24/05	20.9		16.3		74	172		0.5	3	4
6/7/05	24.1		12.78		75	314		0.4	1	4
6/21/05	24.6		15		270	261		0.2	4	5
7/6/05	23.9		6.05		210	299		0.2	4	5
7/19/05	24.9		8.76		180	291		0.2	4	4
8/4/05	25.2				220	516		0.3	3	4
8/15/05	24.6		11.72		210	426		0.2	4	4
8/29/05					190	231		0.2	4	4
9/16/05	20				160	473		0.3	3	4
9/27/05	17.7		10.4		190	425		0.2	3	4
10/11/05	11.7		13.45		150	349		0.3	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								D	F				F
Chlorophyll a								F	C				F
Secchi Depth								F	D				F
Overall								F	D				F

Source: Metropolitan Council and STORET data



Carol Lake (82-0017) Carnelian - Marine Watershed District

Carol Lake is located within Stillwater Township (Washington County). The lake covers an area of 63 acres and has a maximum and mean depth of 1.8 m (roughly 6 feet) and 0.9 m (3 feet). Because of the shallowness of the lake, the entire lake is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation, and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The approximate volume of the lake is 186 acre-feet (ac-ft). The lake's watershed of 375 acres translates to a watershed-to-lake size ratio of 6:1. The larger the ratio the greater the potential stress put on the lake from surface runoff.

This was the sixth year that Carol Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake revealed a fair amount of historic data (1996-2004).

The lake was monitored seven times from mid-April to early-October, 2005. The collected data and resulting graphs showing TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	47.2	24.0	103.0	C
CLA (µg/l)	11.5	4.7	27.0	B
Secchi (m)	1.0	0.3	1.4	D
TKN (mg/l)	0.78	0.59	1.20	
Overall Grade				C

Although no "statistically significant" trend can be determined from the lake's water quality database, the 2003-2005 overall grades are the lake's worst to date. The lake had received overall grades of B in the earlier years of monitoring (1996-2001). In fact, the lake's Secchi transparency grade has steadily fallen from B's in 1996-1999, to C's in 2000-2001, to D's in 2002-2003. This decrease in the lake's short-term water quality should cause some concern and a watchful eye should be kept on the lake's future quality. To better understand the lake's overall water quality and where it may truly be heading, more data are needed.

As mentioned in past reports, the lake's overall grade may be skewed due to the shallowness of the lake. When looking at the lake's 2000 and 2001 mean TP and CLA readings, it seems that the associated Secchi readings could have been limited by the shallowness of the lake rather than excessive nutrients and algal growth. So, while the lake only received an overall grade of B, the actual water quality may have been more representative of an A. This, however, does not explain the drop in mean clarity from grades of B in the late-1990's, to C in 2000-2001, and D in 2002-2005.

Additionally, the difference between the TP, CLA and Secchi grades in recent years (see report grade on the lake's information page), may indicate that suspended sediments may play a large role in the inner workings of the lake. This scenario can be fairly typical for shallow lakes where wind action and storm sewer inflow either increase the influx of sediments to the system or cause the re-suspension of existing bottom sediments. That is, the suspended sediments influence the lake's phosphorus make-up (a larger portion of the in-lake phosphorus in particulate form rather than a soluble form more readily available for algal uptake), reduce water clarity, and could actually be limiting the amount of light available for algal growth, thus keeping the CLA concentrations down (resulting in a better than expected grade). The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.0 for physical condition (3-

“definite algae present”), and 4.4 for recreational suitability (between 4- “no swimming – boating ok” and 5- “no aesthetics possible”).

If you notice any errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

**Carol Lake
Stillwater Twp.,
Washington Co.**

LAKE ID: 820017

WD: Carnelian-Marine

Volunteer: Washington Co.

SWCD





• Sampling site

Contours in meters



2005 Data

Date	Surf Temp	Bot Temp	Surf DO	Bot DO	CLA	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	kg/L	kg/L	kg/L	M	1 to 5	1 to 5
4/20/05	17.7	16.9	4.46	3.72	11	42		1.5	3	4
5/17/05	13.3		6.6	6.7	7.7	103		1.4	2	4
6/16/05	22.2	21.8	4.09	3.18	6.2	36		1.1	2	4
7/14/05	27.6	26.6	3.48	0.15	12	36		0.9	4	5
8/12/05	24.9	23.6	4.96	0.77	20	38		0.3	4	5
9/6/05	25.4		10	2.54	4.7	24		1.2	3	4
10/4/05	20.5		7.99	5.23	11	22		1.4	2	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				B	A	A	A	A	B		C	C	C
Chlorophyll a				B	C	C	C	A	A		B	B	B
Secchi Depth				B	B	B	B	C	C	D	D	D	D
Overall				B	B	B	B	B	B		C	C	C

Source: Metropolitan Council and STORET data

Cates Lake (70-0018) Prior Lake – Spring Lake Watershed District

Cates Lake is a 27-acre lake located in the City of Savage (Scott County). The maximum depth of the lake is 4.0 m (roughly 13 feet). Because of the shallowness of the lake, its entire area is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation), and the lake does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake has no public access.

This was the fourth year that Cates Lake has been involved in CAMP (2002 being the first). The lake was monitored 13 times between mid-May and mid-October, 2005. During each monitoring event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as its perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	22.0	15.0	40.0	A
CLA (µg/l)	4.0	2.5	5.8	A
Secchi (m)	1.9	1.8	2.4	C
TKN (mg/l)	0.77	0.59	1.10	
<i>Overall Grade</i>				B

To the best of our knowledge, there are no water quality data available for Cates Lake other than the 2002-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term however, the lake's water quality is well represented by an overall grade of B. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

During each monitoring event, the volunteers' opinion of the lake condition was ranked on a 1-to-5 scale as shown on the lake information sheet. The average score for physical condition was 3.0 (between 2- "some algae present" and 3- "definite algae present"), and 4.0 for recreational suitability (4 - "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Cates Lake

Prior Lake, Scott Co.


LAKEID: 700018

WD: Prior Lake- Spring Lake

Volunteer: Tom Sletta





• Sampling site

Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/20/05	19.6				5.8			1.8	3	4
5/29/05	19.6					28		1.8	3	4
6/6/05	22.4				4.2	40		2	2	4
6/20/05	26				2.5	23		1.9	3	4
6/28/05	28				3.8	20		1.8	3	4
7/4/05	24.3				2.8	17		2	3	4
7/19/05	25.4				4.9	18		1.9	4	4
8/1/05	25.4				3.7	19		2.1	3	4
8/15/05	24.4				3.9	23		1.8	3	4
8/27/05	24.3				3.2	22		2	3	4
9/16/05	22.9				4.9	17		1.9	3	4
9/26/05	19.6				4.3	15		1.8	3	4
10/19/05	13.4				4.5	15		2.3	2	4



Lake Water Quality Grades Based on Summertime Averages

Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus										A	B	B	A
Chlorophyll a										A	A	A	A
Secchi Depth										C	C	C	C
Overall										B	B	B	B

Source: Metropolitan Council and STORET data

Cenaiko Lake (2-0654) Anoka County Parks

This was the eighth year in which Cenaiko Lake, located within Coon Rapids Dam Regional Park in the City of Coon Rapids in Anoka County, has been monitored through CAMP. Except for the eight years of CAMP data, a search through the STORET nationwide water quality database for historic data on the lake came up empty.

The lake is maintained by groundwater and has a very small watershed that is completely publicly owned (MDNR 1996). No boats, canoes, or floatables are allowed on the 29-acre man-made lake that is one of only six lakes in the seven-county metropolitan area that are stocked with trout (brook and rainbows). The only fishing access to the lake is two fishing docks and the lake's shoreline. The lake, which is 0.6 miles in circumference, has a maximum depth of 9.1 m (30 ft). Only 12 percent of the lake is considered littoral zone (the 0-15 foot depth zone of the lake dominated by aquatic vegetation). Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has been reported on the lake.

Cenaiko Lake was monitored 14 times between mid-April and mid-October, 2005. The data and resulting graphs showing seasonal variability in TP and CLA concentrations, Secchi transparency, and user perceptions are presented on the information sheet following these written comments.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	11.5	10.0	14.0	A
CLA (µg/l)	3.6	1.2	13.0	A
Secchi (m)	2.96	2.30	4.40	B
TKN (mg/l)	0.49	0.31	0.79	
<i>Overall Grade</i>				A

No statistically significant trends are evident from the lake's water quality database. The lake seems well represented by an overall grade of B+/A. To better understand the quality of the lake and what direction it may be heading, continued monitoring is recommended.

At each monitoring event, the volunteers' opinion of the lake condition was ranked on a 1-to-5 scale as shown on the lake information sheet. The average score for physical condition was 2.0 (2- "some algae present"), and 1.3 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problems").

Cenaiko Lake was one of eight lakes in Minnesota and one in Wisconsin that were a part of a research project supported by the MDNR and conducted by researchers at the University of Minnesota that has examined the possibilities of an aquatic weevil *Euryhchiopsis lecontei* as a biological control agent for EWM (U.S.EPA 1997). The following is an excerpt from a U.S.EPA document detailing research in weevils as a biological control:

Of the nine sites, the most pronounced weevil infestation was found in Cenaiko Lake in Anoka County, Minnesota. Weevils caused severe damage to the EWM plants in Cenaiko Lake, most likely resulting in the plants' decreased abundance. EWM biomass (wet weight) at Cenaiko declined from 974 g/m² in July 1996, to 239 g/m² in September 1996 (Newman et al. 1996). Researchers estimate that the biomass in June 1996 (before sampling) was close to 2,000 g/m² (Newman et al. 1996). In July 1996, EWM was approximately 50 percent of the total plant biomass in the lake; by September 1996, this value had decreased to 14 percent.

Monitoring of Cenaiko Lake did not begin until June 1996 when a dense population of weevils was discovered during reconnaissance studies for introduction sites (Newman et al. 1996). Cenaiko Lake was then added to the list of regular sampling sites. Plant samples collected at Cenaiko Lake, as well as at other sites, were processed for invertebrates, plant biomass, and stem damage.

Because monitoring is still ongoing, sampling and data are limited for this study. However, the preliminary results indicate the weevils in Cenaiko Lake may be responsible for the natural decline of EWM.

Since that report however, the lake's biological make-up has changed slightly. The lake's Sunfish population has dramatically increased, which has resulted in a reduced aquatic weevil population (the Sunfish feed on the weevils). The reduction in the aquatic weevil population has resulted in an increase in abundance of EWM within the lake.


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Cenaiko Coon Rapids, Anoka Co.





Lake ID: 20654
WD: Coon Creek
Volunteer: Anoka Co. Parks

● Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/11/05	13.1				6.1	23		1.1	2	
4/26/05	11				1.7	14		2.6	2	
5/12/05					2.7	14		2.6	2	
5/27/05	19.1				3.5	11		2.3	2	
6/9/05					2.8	11		3.8	2	
6/20/05	27				1.5	12		3.2	2	2
7/6/05	26.3				1.8	10		4.4	2	1
7/19/05	28.4				1.2	11		2.5	2	1
8/3/05	24.5				2.6	13		3.2	2	1
8/15/05					1.7	11		2.5	2	1
8/30/05	22.4				5.3	11		2.6	2	2
9/15/05	21.8				13	11		2.5	2	1
10/5/05	18.9				20	17		2	2	1
10/12/05	14.4				56	17		2.4	1	1



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus					A	A	A	A	A	A	A	A	A
Chlorophyll a					A	A	A	A	A	A	A	A	A
Secchi Depth					C	A	A	B	C	A	A	B	B
Overall					B	A	A	A	B	A	A	A	A

Source: Metropolitan Council and STORET data

Cloverdale Lake (82-0009) Valley Branch Watershed District

Cloverdale Lake is a 45-acre landlocked lake located within Baytown Township (Washington County). The mean and maximum depth of the lake is 3.0 m (roughly 10 feet) and 8.5 m (almost 30 feet), respectively. Roughly 86 percent of the lake's area is considered littoral (the 0-15 foot depth area of aquatic vegetation dominance). The lake's size and mean depth results in an approximate lake volume of 450 ac-ft.

The lake's surface area and watershed size (819 acres) translates to an 18:1 watershed-to-lake size ratio. Generally the larger the ratio, the greater the potential stress on the lake from surface runoff.

This was the fifth year that Cloverdale Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake came up empty. Thus, 2001-2005 CAMP data are the only years of available nutrient data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 12 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	28.8	23.0	35.0	B
CLA (µg/l)	6.3	2.3	12.0	A
Secchi (m)	3.0	1.6	4.3	A
TKN (mg/l)	0.84	0.48	1.30	
Overall Grade				A

The lake's 2005 overall lake quality grade of A (better than the B's recorded in 2002-2004, and C in 2001) is the best recorded to date.

As mentioned earlier, there are no nutrient data available for Cloverdale Lake other than the 2001-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

The perceived physical and recreational conditions (ranked on a 1-to-5 scale) are shown on the lake's information sheet on the next page. The average user perception rankings, were 1.8 for physical condition (between 1- "crystal clear" and 2- "some algae present"), and 1.6 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Cloverdale Lake Lake Elmo, Washington Co.

● Sampling site
Contours in meters

Lake ID: 820009
WVD: Lower St. Croix Valley
Volunteer: Kevin Bjork



Bathymetry
Unknown

2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/18/05	14.6				1.7	24		4.8	1	1
5/12/05	18.2				4.7	30		4.3	1	1
5/30/05	21.7				4.4	34		3.8	2	1
6/6/05	24.5				3.4	35		3.1	2	2
6/21/05	28.3				2.3	23		3.4	2	1
7/6/05	26.8				8.6	28		3.1	1	1
7/30/05	27.6				4.8	26		2.4	2	2
8/14/05					6.5	26		2.3	2	2
9/7/05					12	27		3.1	2	2
9/18/05					7.6	30		1.6	2	2
10/2/05					13	41		2.3	2	2
10/11/05					3.4	54		3.2	1	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus									C	C	C	C	B
Chlorophyll a									B	B	B	B	A
Secchi Depth									C	B	B	A	A
Overall									C	B	B	B	A

Source: Metropolitan Council and STORET data



Cobblecrest (27-0053) City of St. Louis Park

Cobblecrest Lake is a small shallow lake located within City of St. Louis Park (Hennepin County). There is very little known morphological data available for the lake.

This marks the second year in which Cobblecrest Lake has been involved in CAMP (2002 and 2004 being the others). Other than for the mentioned CAMP data, a search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Thus, 2002 and 2004-2005 are the only complete, year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 16 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	185.1	126.0	264.0	F
CLA (µg/l)	144.2	44.0	280.0	F
Secchi (m)	0.3	0.1	0.5	F
TKN (mg/l)	3.32	2.1	5.40	
<i>Overall Grade</i>				F

As mentioned earlier, there are no water quality data available for Cobblecrest Lake other than the 2002 and 2004-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 1.3 for physical condition (between 1- "crystal clear" and 2- "some algae present"), and 4.0 for recreational suitability (4- "no swimming – boating ok").


If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Cobblecrest Lake St. Louis Park, Hennepin Co.

Lake ID: 270053
WID: Minnehaha Creek

Volunteers:
Jim and
Graham Kellogg

◆ Sampling site
Contours in meters



2005 Data





Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	CLA ug/L	Surf. TP ug/L	Bot. TP ug/L	Secchi M	PC 1 to 5	RS 1 to 5
4/5/05	11.8					194		0.4	4	2
4/17/05	15.2				72	200		0.3	1	4
5/1/05	9.1				75			0.5	1	4
5/15/05	13.1				93	215		0.3	1	4
5/17/05	15.4				102	212		0.4		
5/25/05						125				
5/30/05	17.3				160	264		0.1	1	4
6/12/05	21.8				280	215		0.2	1	4
6/26/05	28.1				170	199		0.2	1	4
7/10/05	30.2				44	125		0.5	1	4
7/26/05	23.4				150	176		0.3	1	
8/1/05	30				130	170		0.3	2	4
8/15/05	24.9				170	180		0.2	2	4
9/16/05	22				170	153		0.3	2	4
10/2/05	20.7				200	161		0.3	2	4
10/16/05	13.9				60	117		0.4	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus										C	D	F	F
Chlorophyll a										C	F	F	F
Secchi Depth										C	F	F	F
Overall										C	F	F	F

Source: Metropolitan Council and STORET data



Cobblestone Lake (19-0456) City of Apple Valley

Birch Lake is a small lake located in Apple Valley (Dakota County). There is very little morphological data available for the lake.

This marks the first year in which Cobblestone Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake came up empty. Therefore, 2005 is the only complete year of water quality data available. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP ($\mu\text{g/l}$)	93.7	66.0	137.0	D
CLA ($\mu\text{g/l}$)	60.5	33.0	91.0	D
Secchi (m)	0.5	0.4	0.8	F
TKN (mg/l)	1.78	1.30	2.10	
<i>Overall Grade</i>				A


Throughout the monitoring period, the volunteers ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The resulting user perception rankings are shown on the information sheet. The mean physical condition ranking was 3.0 (3- "definite algae present"), while the mean recreational suitability ranking was 3.3 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

Because of the limitedness of the lake's water quality database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Cobblestone Lake Apple Valley, Dakota Co.





Lake ID: 190456
WMO: Vermillion River
Volunteer: City of Apple Valley



● Sampling site
Contours in meters

2005 Data

Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	CLA µg/L	Surf. TP µg/L	Bot. TP µg/L	Secchi M	PC 1 to 5	RS 1 to 5
4/14/05	11.8				62	124		0.4	4	2
4/28/05	10.5				44	105		0.4	3	2
5/10/05	14.7				63	137		0.4	2	2
5/23/05	15				60	129		0.4		2
6/9/05	23				50	98		0.5		2
6/24/05	25.4				90	87		0.4	3	4
7/6/05	24.5				91	94		0.4	3	4
7/21/05	27.8				68	81		0.4		4
8/4/05	25.8				59	74		0.4	3	4
8/16/05	26				69	99		0.5	3	4
9/2/05	22.6				33	90		0.6		4
9/14/05	22				45	76		0.7	3	4
9/27/05	19.7				37	66		0.8	4	2
10/14/05	13.7				68	106		0.5	2	4



Lake Water Quality Grades Based on Summer Time Averages

Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													D
Chlorophyll a													D
Secchi Depth													F
Overall													D

Source: Metropolitan Council and STORET data

Colby Lake (82-0094) City of Woodbury

Colby Lake is located in the City of Woodbury in Washington County. Colby Lake's database now includes 12 years of CAMP collected data (1994-2005). Analysis on the lake's water quality database reveals no statistically significant trend in its water quality (either improving or degrading). The lake's water quality seems well represented by an overall water quality grade of D/F.

Information from the City of Woodbury revealed that the lake has a surface area of 71 acres and a maximum depth of just 3.4 m (11 feet). The lake's large 8,088-acre contributing watershed results in a large 114:1 watershed-to-lake size ratio. The larger the ratio the greater the potential for stress on the lake from surface runoff. Because of the shallowness of the lake, its entire area is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation), and the lake does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake has no public access.

As part of the city's involvement in CAMP in 2005, the lake was monitored 6 times between late-April and late-October. During each sampling event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	124.3	46.0	195.0	D
CLA (µg/l)	151.7	16.0	340.0	F
Secchi (m)	0.7	0.4	1.3	D
TKN (mg/l)	1.73	1.20	2.10	
Overall Grade				D

The lake's 2005 overall grade (D) was similar to that of 1994, 1997, 1999-2000, 2002, and 2004, and better than that of 1995, 1996, 1998, 2001, and 2003 (F's).

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the following page. The mean physical condition ranking was 2.3 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 4.0 (4- "no swimming - boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.


Colby Lake Woodbury, Washington Co.

Lake ID: 820094

WCD: South Washington




Volunteers: The Hvas Family

• Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 thru 5	1 thru 5
4/24/05	15.7				17	202		1.9	1	4
5/30/05	21.3				16	46		1.3	1	4
8/1/05	26				340	195		0.5	3	4
9/1/05	25.3				99	132		0.4	3	4
10/11/05	16.8				80	153		0.5	3	4
10/23/05	10.1				190	120		0.4	1	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	D	D	F	F	F	F	D	D	F	F	D	D	D
Chlorophyll a	D	F	F	C	F	F	D	F	C	D	C	F	F
Secchi Depth	F	F	F	F	F	D	D	D	F	F	F	D	D
Overall	D	F	F	D	F	D	D	F	D	F	D	D	D

Source: Metropolitan Council and STORET data



Cornelia Lake (27-0028-01) Conservation League of Edina

Lake Cornelia is a small shallow lake located within Edina (Hennepin County). There is very little known morphological data available for the lake.

This marks the second year in which Lake Cornelia has been involved in CAMP (2003 being the first). In fact, a search through the STORET nationwide water quality database for historic data on the lake produced only the mentioned CAMP collected data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored seven times between mid-May and late-September, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	155.8	112.0	236.0	F
CLA (µg/l)	61.9	25.0	98.0	D
Secchi (m)	0.4	0.3	0.7	F
TKN (mg/l)	2.05	1.20	4.70	
<i>Overall Grade</i>				F

While the lake's 2005 mean TP and CLA concentrations are better than those recorded in 2003, the 2005 Secchi mean was worse.

Because of the limitedness of the Lake Cornelia water quality database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.




Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.0 for physical condition (between 3- "definite algae present"), and 4.0 for recreational suitability (between 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Lake Cornelia Edina, Hennepin County


Lake ID: 270028-01
WD: Nine Mile Creek
Volunteer: Conservation League of Edina

● Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/13/05	12.3				43	119		0.7	3	4
5/25/05	19				37	115		0.5		
5/26/05	20				42	120		0.5	3	4
6/10/05	23.6				41	112		0.5	2	4
6/24/05	28				63	196		0.4	3	4
7/8/05	27				96	176		0.3	3	4
7/22/05	29				25	236		0.3	2	4
8/5/05	29				62	141		0.4	3	4
8/19/05	25				97	161		0.4	4	4
9/9/05	24.3				85	151		0.4	4	4
9/23/05	20				63	146		0.4	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus											F	F	F
Chlorophyll a											F	D	D
Secchi Depth											F	F	F
Overall											F	F	F

Source: Metropolitan Council and STORET data



Courthouse Lake (10-0005) Carver County Environmental Services

Courthouse Lake, located in the City of Chaska (Carver County) is a unique resource in the Twin Cities Metropolitan Area. The lake is only one of six lakes in the seven-county metropolitan area stocked with trout (rainbows). Very little lake data (or physical information) are available for Courthouse Lake. The 10-acre lake (0.6 miles in circumference) has a maximum depth of 17.4 m (57 feet) and only three percent of the lake is considered littoral zone (the 0-15 foot depth zone of the lake dominated by aquatic vegetation). The lake's level is maintained by groundwater. It has a very small watershed that is completely publicly owned (MDNR 1996).

The only data available for Courthouse Lake are a result of CAMP monitoring from 1996-2005.

Courthouse Lake was monitored biweekly from mid-April to mid-October, 2005, for a total of 14 monitoring events. The data collected by volunteers showed seasonal variability in TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability). Results are presented on the lake's information sheet.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	11.8	5.0	23.0	A
CLA (µg/l)	2.0	1.0	3.5	A
Secchi (m)	4.9	3.0	7.1	A
TKN (mg/l)	0.48	0.30	0.65	
Overall Grade				A

The lake's 2005 overall grade was similar to that of 1996, 1998-2001, and 2003-2004, and better than 1997 and 2002 (overall grades of B). When comparing the lake's historical summer means, it is apparent that 2005 was the lake's best overall water quality year and 1997 was the worst.

Analysis on the lake's water quality database reveals no statistically significant trend in its water quality (either improving or degrading). The lake's water quality seems well represented by an overall water quality grade of A/B+.

The average user perception rankings, on a 1-to-5 scale, were 1.1 for physical condition (between 1- "crystal clear" and 2- "some algae present"), and 1.0 for recreational suitability (1- "beautiful").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you detect any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Courthouse Lake





Chaska, Carver Co.

Lake ID: 100005
WQ: Lower Minnesota River
Volunteer: Carver Co.
Env. Services



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	kg/L	kg/L	kg/L	M	1 to 5	1 to 5
4/14/05	13.5		12.4		8.8	32		1.6	2	1
4/29/05	10.7		13.3		1.8	22.5		5.2	1	1
5/10/05	15.8		13		3.5	22		6	1	1
5/23/05	19.1		8.5		1.8	23		3	1	1
6/6/05	21.9		9.32		1	12		4.7	2	1
6/22/05	24.2		9.03		1	8		7.1	1	1
7/5/05	25.2		8.26		1	6		6.7	1	1
7/20/05	27.3		7.89		1	8		5.6	1	1
8/2/05					1.9	7.5		4.2	1	1
8/17/05	25.8		8.54		3.3	11		4.3	1	1
8/29/05	25.6		8.21			10		3.7	1	1
9/13/05	23.8		7.53		2.5	12		4	1	1
9/27/05	21.6		7.2		2.6	15		4.6	1	1
10/11/05	16.1		9.77		5.5	18		3.5	1	1



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				A	A	A	A	A	A	B	A	A	A
Chlorophyll a				A	A	A	A	A	A	A	A	A	A
Secchi Depth				A	C	A	B	A	A	B	A	B	A
Overall				A	B	A	A	A	A	B	A	A	A

Source: Metropolitan Council and STORM data

Crystal Lake [Burnsville] (19-0027) Black Dog Watershed Management Commission

Crystal Lake is located mainly in the City of Burnsville (Dakota County) covers an area of 292 acres, with 5.3 miles of shoreline. The maximum and mean depths of the lake are 11.3 m (37 feet) and 3.1 m (10 feet), respectively. The lake's surface area and mean depth translate to an approximate lake volume of 2,920 acre-feet. The lake's watershed covers approximately 2,001 acres of which roughly two-thirds is urban/developed. The watershed and lake surface areas translate to a moderate watershed-to-lake size ratio of 7:1 (the smaller the ratio the less stress on the lake from surface runoff).

Roughly 72 percent of the lake's area is considered littoral (the 0-15 foot depth area of aquatic vegetation dominance).

Because of its multi-recreational uses, the lake is considered a "Priority Lake" in the Metropolitan Area. The lake, managed by the MDNR as a panfish lake and stocked with tiger muskellunge, has a public access and fishing pier on its north side and a public swimming beach on its eastern shore. One problem that may possibly hinder future recreational activity on the lake, however, is Eurasian Water Milfoil (*Myriophyllum spicatum*), which has been reported in the lake.

This was the seventh year that Crystal Lake has been involved in CAMP (1999-2005). The lake was monitored during each of the five years prior to 1999 by Council staff. A search of the STORET nationwide water quality database for data on the lake revealed an extensive database since the 1980's, with nutrient data available in 1980, 1983, 1989, and 1994-2005. Additionally, Secchi transparency data are available for all years between 1980 and 1999 except 1993.

The lake was monitored 16 times between mid-April and mid-October, 2005. Results are presented on graphs and data tables on the following page. During each monitoring event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as its perceived physical condition and recreational suitability.

2005 summer (May-September) data summary




<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	37.3	21.0	58.0	C
CLA (µg/l)	22.4	5.6	50.0	C
Secchi (m)	1.6	0.9	2.7	C
TKN (mg/l)	0.85	0.46	1.60	
Overall Grade				C

The 2005 grade is similar to those recorded from 1994-2000, and 2002-2004, and worse than 1983, 1989, and 2001. Analysis on the lake's water quality database reveals no statistically significant trend in its water quality (either improving or degrading). The lake's water quality seems well represented by an overall water quality grade of C/B-.

Throughout the monitoring period, the volunteer's opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The average user perception rankings, were 2.5 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 2.2 for recreational suitability (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1-5	1-5
4/17/05	13.9				7.9	22		1.9	1	1
4/30/05	11.2					26		2.3	2	1
5/15/05	12.1				5.6	22		2.7	3	2
5/18/05	15.2				6.2	23		2.5		
5/25/05	16.2				7.6	21		2.4	3	3
6/11/05	24				8.5	40		2.2	3	2
6/25/05	25.8				31	98		1.3	2	2
7/4/05	24				42	52		0.9	3	2
7/23/05	27.5				23	40		1.1	2	2
8/6/05	26.1				24	35		1	3	3
8/20/05	24.6				28	40		1.5	2	2
9/3/05	21.8				50	40		1	2	2
9/17/05	21.3				25	39		1.6	2	2
10/1/05	18.3				30	55		1.3	3	2
10/15/05	14.7				25	75		1.7	3	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C	C		C						B			
Chlorophyll a	C			B				C		B			
Secchi Depth	C	C	C	B	C	B	B	C	C	B	C	B	B
Overall	C			B						B			

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		C	C	C	C	C	C	C	B	C	C	C	C
Chlorophyll a		B	C	C	C	C	B	C	B	B	C	B	C
Secchi Depth		C	C	C	C	C	C	C	C	C	C	C	C
Overall		C	C	C	C	C	C	C	B	C	C	C	C

Source: Metropolitan Council and STORET data



Crystal Lake [Robbinsdale] (27-0034) *Shingle Creek Watershed Management Commission*

Crystal Lake is a 76-acre lake located in the City of Robbinsdale (Hennepin County). The lake was first enrolled in CAMP in 1994. The lake has a maximum and mean depth of 10.4 and 3.7 m (34 and 12 feet), and an approximate volume of 917 ac-ft (Beduhn 1993). Sixty-eight percent of the lake's surface area is considered littoral zone (the 0-15 foot depth area of aquatic vegetation dominance). The lake's fishing pier and earthen public access are located on the southeastern end of the lake.

The lake, managed by the MNDNR as a warm-water gamefish lake, has been stocked with muskellunge and northern pike in the past. Because of the numerous historical winter fishkills, an aeration system was installed in 1973 to maintain the lake's oxygen levels. While the aerator stopped the winterkill problem, it may have created another. The aeration system may actually allow phosphorus in the sediment to circulate throughout the lake's water column by disrupting the thermocline. In fact, recent summerkills have been reported because the lake's high TP concentrations lead to heavy algal blooms which in turn choke the lake's oxygen load.

The lake's watershed is mostly urban/developed with some undeveloped/park areas. The urban areas consist of single family residential homes around portions of the lake, as well as multi-family and commercial/industrial areas within the watershed. The majority of the undeveloped areas are associated with parkland on the shores of the lake. Eighty-nine percent of the land use within the lake's 1,272 acre watershed is urban, while the remaining 11 percent is undeveloped (Beduhn 1993). The resulting watershed-to-lake size ratio is 10.4:1.

Crystal Lake, which was monitored through CAMP in 1994, 1997, and 2001, was monitored seven times from mid-June to early-October in 2005. On each sampling date the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	113.5	80.0	136.0	D
CLA (µg/l)	41.2	17.0	110.0	C
Secchi (m)	0.9	0.5	1.7	D
TKN (mg/l)	2.48	0.80	3.20	
Overall Grade				D

Results are presented on graphs and data tables on the following page.

Graphs showing the volunteer perceptions of the lake's physical condition and recreational suitability seem to be correlated to the other graphs for this lake. Generally, the greater the lake's clarity, the better the lake's perceived physical condition and recreational suitability. The summertime mean recorded physical condition was 2.7 on a 1-to-5 ranking scale (between 2- "some algae present" and 3- "definite algal presence"). The mean suitability for recreation ranking was also 3.1 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

The water quality database for Crystal Lake consists of nutrients and Secchi data in 1986-1988, 1992, 1994, 1997, 2001, and 2005. The lake quality grades seem to show that the lake's quality has remained rather similar through the mid-1980's through the mid-2000's.

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a

fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-1267 or randy.anhorn@metc.state.mn.us.



Crystal Lake Robbinsdale, Hennepin Co.

Lake ID: 270034

WMO: Shingle Creek




Volunteer: Wayne Sicora

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
6/11/05	22.5				31	80		0.7	3	4
6/25/05	25.6				25	122		1	2	3
7/16/05	31				110	93		0.6	3	4
8/11/05	25.4				17	126		0.5	4	4
9/15/05	22.1				34	124		1.7	2	2
9/23/05	20.9				60	136		1	2	2
10/8/05	15.4				41	139		1.4	2	1




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus							D	D	F				F
Chlorophyll a							D	D	F				D
Secchi Depth							D	D	F				D
Overall							D	D	F				D

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		F			F				D				D
Chlorophyll a		C			C				C				C
Secchi Depth		D			C				D				D
Overall		D			D				D				D

Source: Metropolitan Council and STORET data



Dean Lake (70-0074) City of Shakopee

Dean Lake is a small shallow lake located within City of Shakopee (Scott County). There is very little known morphological data available for the lake). Because of the shallowness of the lake, its entire area is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation), and the lake does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

Two thousand and four marks the third year in which Dean Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Thus, 2002-2005 are the only years of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored seven times between late-April and mid-September, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	214.3	66.0	328.0	F
CLA (µg/l)	47.8	1.7	150.0	C
Secchi (m)	0.3	0.2	0.4	F
TKN (mg/l)	3.15	1.20	6.20	
Overall Grade				F

The difference between the TP, CLA and Secchi grades in current and past years (see report grade on the lake's information page), may indicate that suspended sediments may play a large role in the inner workings of the lake. This scenario can be fairly typical for shallow lakes where wind action and storm sewer inflow either increase the influx of sediments to the system or cause the re-suspension of existing bottom sediments. That is, the suspended sediments influence the lake's phosphorus make-up (a larger portion of the in-lake phosphorus in particulate form rather than a soluble form more readily available for algal uptake), reduce water clarity, and could actually be limiting the amount of light available for algal growth, thus keeping the CLA concentrations down (resulting in a better than expected grade).


As mentioned earlier, there are no water quality data available for Dean Lake other than the 2002-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.2 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 4.0 for recreational suitability (4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Dean Lake, Shakopee, Scott Co.

Lake ID: 700074
WD: Lower Minnesota River
Volunteers: The Gerlach Family



2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/23/05	12.5				7.3	7.4		0.4	1	4
5/1/05	18.5				4.4	66		0.4	1	4
5/28/05	17.5				7.8	141		0.4	2	4
7/1/05	28				1.7	313		0.2	3	4
8/1/05	30.5				150	304		0.2	3	4
8/30/05					87	328		0.2	2	4
9/14/05	20				36	134		0.3	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus										F	F	D	F
Chlorophyll a										D	C	B	C
Secchi Depth										F	F	F	F
Overall										F	D	D	D

Source: Metropolitan Council and STORET data



DeMontreville Lake (82-0101) Valley Branch Watershed District

Lake DeMontreville, located in Lake Elmo (Washington County), has public access on its northwestern side, and is considered a “Priority Lake” due to its multi-recreational uses. The 160-acre lake with a mean and maximum depth of 2.4 m (~8 feet) and 7.3 m (24 feet), was monitored seven times from late-April to early-October, 2005. Roughly 90 percent of the lake’s area is considered (the 0-15 foot depth area of aquatic vegetation dominance). The lake’s size and mean depth results in an approximate lake volume of 1,280 ac-ft.

The lake’s surface area and watershed size (1,108 acres) translates to a 7:1 watershed-to-lake size ratio. Generally the larger the ratio, the greater the potential stress on the lake from surface runoff.

This was the third year that Lake DeMontreville has been involved in CAMP. The lake has been monitored in the past by Council staff (most recently in 2003). A search of the STORET nationwide water quality database for data on the lake revealed a moderate database since the 1980’s with nutrient and Secchi transparency data available in 1980, 1984, 1991, 1993, 1995, 2000 and 2003-2005. Additionally, Secchi transparency data are available for 1985-1986, and 1988-1989.

2005 summer (May-September) data summary




<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	24.8	14.0	39.0	B
CLA (µg/l)	12.7	2.9	36.0	B
Secchi (m)	3.2	1.4	5.6	A
TKN (mg/l)	1.00	0.57	1.70	
Overall Grade				B

Historically, 1980-2005 lake quality grades for Lake DeMontreville (see lake information sheet on the following page) show that the quality of the lake has improved over the past 25 years. The overall grades in 1980, 1984, and 1991 were all C. The overall grades in 1993, 1995, and 2005 were B, and the overall grades for 2000, and 2003-2004 were A. A recent MPCA conducted trend analysis on the lake’s Secchi transparency data, revealed a statistically significant improvement in recent water clarity.

The graphs showing the volunteer's perceptions of the lake's physical condition and recreational suitability seem somewhat correlated to the other graphs for this lake. The better the lake's clarity (also relating to lower TP and CLA concentrations), the better the lake's perceived physical condition and recreational suitability. The summertime mean recorded physical condition was 2.2 on a 1 to 5 ranking scale shown on the lake information sheet (between 2- "some algae present" and 3- “definite algae present”). The mean suitability for recreation ranking, also on a 1-to-5 scale, was 2.2 (between 2- " minor aesthetic problem" and 3- “swimming slightly impaired”).


The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

Date	Surf Temp	Bot Temp	Surf DO	Bot DO	CIA	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ig/L	ig/L	ig/L	M	1 to 5	1 to 5
4/1/05	14.9	7.1	6.65	0.1	3.3	42		4.5	3	2
5/2/05	15.3	10.2	5.57	5.57	4.2	14		4.1	2	2
6/1/05	23.9	12.5	7.56	0.57	2.9	20		3.4	2	2
7/1/05	30	13.1	7.91	0.08	3.3	20		1.5	2	2
8/1/05	26.4	15.8	5.39	0.19	17	31		1.4	3	3
9/9/05	22.1	14	8.42	0.73	36	39		1.4	3	3
10/3/05	19.4	17.2	7.47	1.98	27	31		3	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C				C							B	
Chlorophyll a	C				C							C	
Secchi Depth	C				C	C	C		C	D		C	
Overall	C				C							C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B		C					A			A	A	B
Chlorophyll a	A		B					A			B	A	B
Secchi Depth	B		B					A			A	B	A
Overall	B		B					A			A	A	B

Source: Metropolitan Consolidated STORET data



Downs Lake (82-0110) Valley Branch Watershed District

Downs Lake, located in Lake Elmo (Washington County), was monitored nine times between late-May and early-October, 2005. The mean and maximum depths of the 35-acre lake are 1.5 m (5 feet) and 2.1 m (7 feet), respectively. The lake's size and mean depth results in an approximate lake volume of 175 ac-ft. Because of the shallowness of the lake, the entire lake is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's 2,400-acre watershed translates to a large watershed-to-lake size ratio of 69:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This was the sixth year in which Downs Lake has been involved in CAMP (1999 and 2001-2004 being the others). A search through the STORET nationwide water quality database for data on the lake resulted in no data other than that collected through CAMP. Thus, 1999 and 2001-2005 are the only years where data are available. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	141.8	80.0	269.0	D
CLA (µg/l)	67.0	13.0	100.0	D
Secchi (m)	0.4	0.2	0.8	F
TKN (mg/l)	2.04	1.10	3.20	
<i>Overall Grade</i>				D

The summertime means resulted in a TP grade of D, CLA grade of D, and Secchi transparency grade of F. The overall grade, calculated from all three parameters was D. The lake's 2005 overall water quality grade is similar to that recorded in 1999 and 2003, and better than those of 2001-2002 and 2004 (overall grade of F).

As mentioned earlier, there are no water quality data available for Downs Lake other than the 1999 and 2001-2005 CAMP data. Therefore it is not possible to determine any long-term. In the short-term, the lake seems to fluctuate between overall grades of D/F. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer's opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The average user perception rankings, were 3.1 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.0 for recreational suitability (4- "no swimming - boating ok").


If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Downs Lake

Lake Elmo, Washington Co.





Lake ID: 820110
WD: Valley Branch
Volunteers: The Wesley Sly Family

● Sampling site
Contours in meters



2005 Data

	Surf Temp	Bot Temp	Surf DO	Bot DO	Chl a	Surf TP	Bot TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
5/23/05	19.9				13	80		0.8	2	4
6/12/05	29				56	110		0.4	2	4
6/30/05	27				74	119		0.4		
7/1/05	25.8				86	124		0.3	3	4
7/21/05	30.9				64	119		0.4	4	4
7/29/05	27.8				72	138		0.3	3	4
8/15/05	29				100	269		0.2	4	4
9/12/05	25.9				78	175		0.2	4	4
10/10/05	16.1				62	274		0.2	3	4



Lake Water Quality Grades Based on Summertime Averages

Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	D	F	D	F	D	D
Chlorophyll a							D	F	F	C	D	D	D
Secchi Depth							D	F	F	F	F	F	F
Overall							D	F	F	D	F	D	D

Source: Metropolitan Council and STORET data

Eagle Lake [Carver County] (10-0121) Carver County Environmental Services

Eagle Lake is located in Young America Township in Carver County. The lake has a surface area of 233-acres, and a maximum and mean depth of 7.9 m (26 feet) and 1.2 m (4 feet), respectively. Because of the shallowness of the lake, the entire area is considered littoral, (the shallow [0-15 foot depth] area dominated by aquatic vegetation) and does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has been reported on the lake.

The lake has a 1,050-acre immediate watershed, which translates to a watershed-to-lake area ratio of 4.5:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff). A 1999 water quality report on water resources in Carver County estimates land use for the watershed at: two percent residential, 63 percent agricultural, and 35 percent open/undeveloped (Carver County Planning 1999).

This was the seventh year that Eagle Lake has been involved in CAMP (previously enrolled in 1998-2003), although it has been previously monitored by Council staff (as recently as 2004). The lake was monitored 14 times between mid-April and mid-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	151.6	34.0	400.0	D
CLA (µg/l)	47.2	3.9	120.0	C
Secchi (m)	1.3	0.3	3.0	C
TKN (mg/l)	1.36	0.70	2.90	
Overall Grade				C

The lake's overall water quality grades indicate that 2005 was the lake's best recorded water quality year to date.

The perceived physical and recreational conditions of the lake, recorded by the volunteers, were ranked on a 1-to-5 scale. The rankings are shown in both tabular and graphical form on the lake's associated information sheet. The mean physical condition ranking was 3.1 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability ranking was 3.4 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Eagle Lake, Carnden Twp. Carver Co.

Lake ID: 100121
WMO: Crow River
Volunteer: Carver Co.
Env. Services

● Sampling site
Contours in meters



2005 Data





Date	Surf Temp	Bot Temp	Surf DO	Bot DO	Chl a	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/18/05	14.6		9.94		18	62		1.2	3	3
4/27/05	7.8		10.97		14	39		1.6	2	3
5/11/05	14.9		10.8		6.9	68		3	2	2
5/24/05	17				14	37		2.2	3	4
6/6/05	20.7		9.45		3.9	34		2.6	2	2
6/20/05	25		8.3		9.9	98		2.7	2	4
7/1/05	24.1		12.7		21	112		1.1	2	2
7/18/05	27.1		7.85		47	192		0.6	3	3
8/1/05	23.7				120	248		0.4	4	4
8/18/05	24		10.81		120	220		0.3	5	5
8/31/05	22.9		9.62		61	196		0.4	3	4
9/13/05	22.6		5.47		49	112		0.6	4	4
9/23/05	20				61	400		0.5	3	3
10/10/05	12.7		10.11		71	363		0.4	3	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	F	F				F							
Chlorophyll a	D	C				F							
Secchi Depth	C	C				F							
Overall	D	D				F							

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			F		F	F	F	F	F	F	F	F	D
Chlorophyll a			C		C	C	C	C	D	C	C	C	C
Secchi Depth			B		C	B	C	D	D	F	D	D	C
Overall			D		D	D	D	D	D	F	D	D	C

Source: Metropolitan Council and STORET data



Eagle Lake (27-0111-01) Shingle Creek Watershed Management Commission

This was the seventh year that Eagle Lake has been enrolled in CAMP (the lake was previously involved in CAMP in 1993, 1996-1998, 2000, and 2002). The 291-acre lake is located within the City of Maple Grove (Hennepin County). It has a 3,220-acre watershed. The lake and watershed areas translate to a watershed-to-lake area ratio of 11:1. The maximum and mean depths of the lake are 10.4 and 3.8 m (34.0 and 12.5 feet), respectively. Roughly 68 percent of the lake's area is considered littoral (the 0-15 foot depth area dominated by aquatic vegetation). The approximate volume of the lake is 3,667 acre-feet (ac-ft) and a public access to the lake is located on its eastern shores. Because of the lake's multi-recreational uses, the lake is considered a "Priority Lake" in the Metropolitan Area. One problem that may possibly hinder future recreational activity on the lake, however, is Eurasian Water Milfoil (*Myriophyllum spicatum*), which has been reported in the lake.

Eagle Lake was monitored six times from early-May to late-July, 2005. The data and resulting graphs showing seasonal variability in TP and CLA concentrations, Secchi transparency, and user perceptions are presented on the information sheet following these written comments.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	36.2	19.0	83.0	C
CLA (µg/l)	15.5	3.7	55.0	B
Secchi (m)	2.6	1.5	4.3	B
TKN (mg/l)	0.77	0.62	0.97	
Overall Grade				B

The lake's 2005 overall water quality grade is similar to those recorded in 1993, 1996-1998, and 2000, and better than those recorded in the 1980's and early 1990's.

A search for water quality data through Council, MPCA, and STORET files resulted in a good amount of data. There are nutrient data available for 1980, 1983, 1986-1987, 1991, 1993, 1996-1998, 2000, 2002, and 2005. Except for a slight decline in 2002, the lake has shown some improvement in water quality since the 1980's (D in 1980, to C's in 1983-1991 and 2002, and B's in 1996-2000 and 2005).

At each monitoring event, the volunteers' opinion of the lake condition was ranked on a 1-to-5 scale as shown on the lake information sheet. The average score for physical condition was 2.0 (2- "some algae present"), and 1.5 for recreational suitability (between 1- "crystal clear" and 2- "minor aesthetic problems").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Eagle Lake, Maple Grove, Hennepin Co.




Lake ID: 270111-01
WMO: Shingle Creek
Volunteer: Sharon Collins

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/1/05	10.8				3.8	30		2.1		
5/3/05	12				4.2	26		2.4		
5/24/05	18.3				3.7	83		4.3	2	1
6/21/05	27				6.7	28		3.1	2	2
7/3/05	23.9				8.4	19		2.1	2	
7/24/05	28.3				55	31		1.5	2	




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C			C			C	C				C	
Chlorophyll a	D			C			B	C				C	
Secchi Depth	D			C			C	C				C	
Overall	D			C			C	C				C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B			B	B	B		C		C			C
Chlorophyll a	B			B	A	B		A		A			B
Secchi Depth	B			C	C	C		B		D			B
Overall	B			B	B	B		B		C			B

Source: Metropolitan Council and STORET data



Earley Lake (19-0033) Black Dog Watershed Management Commission

Earley Lake is located within the City of Burnsville in Dakota County. The 29-acre lake receives flow from Crystal Lake (Burnsville) and the Earley Lake watershed. Most of its 1,629-acre watershed is either parkland or open space. The watershed-to-lake size ratio is a rather large 56:1. Generally, the larger the ratio the greater the potential stress on the lake from surface runoff. Earley Lake outlets at its west end to Sunset Pond.

Earley Lake has been enrolled in CAMP since 1994. The lake was monitored seven times between mid-June and mid-September, 2005. On each sampling date the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	60.6	26.0	137.0	C
CLA (µg/l)	12.9	6.0	23.0	B
Secchi (m)	1.6	1.2	1.9	C
TKN (mg/l)	0.74	0.32	1.30	
<i>Overall Grade</i>				C

Individual and overall grades for 2005 are identical to those found in 1994-2004. While there has been slight variability in individual summer means from year to year, the lake's quality has remained fairly consistent. The lake's mean TP seems to generally fall within the 50.0-60.0 µg/l range, while the CLA and Secchi means generally range between 11.0-18.0 µg/l and 1.2-1.7 m, respectively. The lake's best water quality was recorded in 2003.


Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The mean physical condition ranking was 2.7 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 4.0 (4- "no swimming – boating ok").

Statistical analysis on the lake's water quality database did not detect any long-term trends, in the short-term however, the lake seems to be very well represented by an overall water quality grade of C. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

If you know of errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Earley Lake Burnsville, Dakota Co.

Lake ID: 190033
WMO: Black Dog Lake
Volunteer: Mary Oaster



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 to 5	1 to 5
6/2/05	26				93	26		1.9	5	4
6/30/05	26				113	32		1.9		
7/2/05	24.6				89	29		1.9	3	4
7/24/05	29.7				6	137		1.2	2	4
8/1/05	26.4				23	95		1.8	3	4
8/28/05	24.6				10	62		1.3	2	4
9/18/05	26.9				30	43		1.4	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C	C	C	C	C	C	C	C	C	C	C	C	C
Chlorophyll a	B	B	B	B	B	B	B	B	B	B	B	B	B
Secchi Depth	C	C	C	C	C	C	C	C	C	C	C	C	C
Overall	C	C	C	C	C	C	C	C	C	C	C	C	C

Source: Metropolitan Council and STORET data



East Lake (19-0349) City of Lakeville

East Lake is a small lake located in Lakeville (Dakota County). There is very little morphological data available for the lake.

This marks the first year in which East Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake came up empty. Therefore, 2005 is the only complete year of water quality data available. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 13 times between late-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary




<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	202.1	92.0	318.0	F
CLA (µg/l)	120.2	53.0	270.0	F
Secchi (m)	0.3	0.2	0.5	F
TKN (mg/l)	2.42	1.50	4.00	
Overall Grade				F

Throughout the monitoring period, the volunteers ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The resulting user perception rankings are shown on the information sheet. The mean physical condition ranking was 3.7 (between 3- "definite algae present" and 4- "high algal color"), and 4.3 for recreational suitability (between 4- "no swimming - boating ok" and 5- "no aesthetics possible").

Because of the limitedness of the lake's water quality database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

East Lake Lakeville, Dakota Co.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/26/05	10.1				46	121		0.7	3	4
5/10/05	17.9				53	223		0.4	4	4
5/23/05	17.9				55	132		0.5	3	5
6/6/05					64	189		0.4	4	5
6/21/05	26.3				140	198		0.2	4	4
7/5/05	22.4				170	211		0.2	3	4
7/20/05	28				270	317		0.2	4	5
8/2/05	26.9				140	318		0.2	4	4
8/16/05	22.4				160	255		0.2	4	4
8/29/05	26.9				110	138		0.3	4	4
9/13/05	20.2				86	92		0.4	3	4
9/27/05	17.9				74	149		0.3	4	4
10/10/05	14.6				42	87		0.5	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													F
Chlorophyll a													F
Secchi Depth													F
Overall													F

Source: Metropolitan Council and STORET data



East Boot Lake (82-0034) *Carnelian - Marine Watershed district*

East Boot Lake, located in May Township (Washington County), was monitored 14 times between mid-April and mid-October, 2005. The mean and maximum depths of the 47-acre lake are 8.2 m (27 feet) and 0.9 m (3 feet), respectively. The lake's size and mean depth results in an approximate lake volume of 282 ac-ft. Because of the overall shallowness of the lake, roughly 82 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance), the majority of the lake does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's small 93-acre immediate watershed translates to a small watershed-to-lake size ratio of 2:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This was the sixth year that East Boot Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake revealed a limited amount of data (1996-2004 and now 2005).

On each sampling date, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	36.1	20.0	53.0	C
CLA (µg/l)	14.5	2.5	43.0	B
Secchi (m)	3.2	0.9	6.4	A
TKN (mg/l)	0.98	0.58	1.40	
Overall Grade				B

Similarly to that seen in 2004, the lake's 2005 minimum and maximum Secchi transparency readings represent the largest range in CAMP 2005. The lake's 2005 overall grade is identical to those recorded through CAMP in 1996-1998 and 2004, and better than the recent grades posted in 1999-2003 (C).

Statistical analysis on the lake's water quality database did not detect any trends. With this in mind however, the lake's recent water quality seems to be well represented by an overall grade of C+/B-. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.9 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 3.5 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

East Boot Lake May Twp., Washington Co.




LAKE ID: 820034

W/D: Carnelian-Marine

Volunteer: Washington Co.


SWCD

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/1/05	12.4	4.9	7.18	0.3	12	102		2.3	2	4
4/25/05	12.9	5.2	5.52	0.1	1	36		5.5	2	3
5/10/05	15.7	5.5	6.33	0.17	4.2	21		5.6	3	4
5/24/05	17.9	5.8	5.44	0.03	2.6	24		6.4	3	4
6/6/05	22.3	6.2	5.1	0.32	2.5	22		6.1	2	3
6/22/05	27	7.2	9.3	0.7	2.9	20		4.6	2	2
7/8/05	24	7.3	10.49	0.36	12	40		2.3	3	3
7/19/05	28	7.4	8.42	0.53	4.5	35		1.4	3	4
8/1/05	30	7.5	9	0.62	43	53		0.9	4	5
8/15/05	26.3	7.8	7.17	0.43	34	52		1.2	3	4
8/29/05	23.6	7.8	4.8	0.62	22	38		1.5	3	3
9/13/05	23.4	8.2	7.41	0.46	24	41		1.5	3	3
9/29/05	17.7	8.3	5.31	0.71	7.3	51		3.8	3	3
10/10/05	15.3	8.6	7.29	0.48	23	30		2.1	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				B	B	B	C	C	C	C	C	C	C
Chlorophyll a				B	C	C	C	C	C	C	C	B	B
Secchi Depth				B	A	B	C	C	C	B	B	A	A
Overall				B	B	B	C	C	C	C	C	B	B

Source: Metropolitan Council and STORET data



Edina Lake (27-0029) Conservation League of Edina

Edina Lake is a small shallow lake (a maximum depth of approximately 1.0 m (3.3 feet), located within Edina (Hennepin County). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). There is very little other known morphological data available for the lake.

This marks the second year in which Lake Edina has been involved in CAMP (2004 being the first). A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Thus, 2004-2005 are the only known years of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 10 times between mid-May and late-September, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	127.7	54.0	199.0	D
CLA (µg/l)	60.0	19.0	110.0	D
Secchi (m)	0.3	0.2	0.7	F
TKN (mg/l)	2.57	1.80	3.80	
<i>Overall Grade</i>				D

The lakes 2005 overall grade is identical to that recorded in 2004. The 2005 individual summer means, however, are worse.

As mentioned earlier, there are no water quality data available for Lake Edina other than the 2004-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.



Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.9 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 4.0 for recreational suitability (4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Lake Edina Edina, Hennepin Co.


LAKE ID: 270029
WD: Nine Mile Creek
Volunteer: Conservation League of Edina

◆ Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
5/13/05	11.3				23	67		0.7	2	4
5/26/05	19.2				19	124		0.3	2	4
6/10/05	24				24	54		0.4	3	4
6/24/05	29				41	106		0.4	3	4
7/8/05	29				29	122		0.4	3	4
7/22/05	32				86	168		0.3	2	4
8/5/05	29				96	199		0.3	4	4
8/19/05	26				110	143		0.3	3	4
9/9/05	24.7				100	146		0.2	4	4
9/23/05	20				72	148		0.2	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												D	D
Chlorophyll a												C	D
Secchi Depth												F	F
Overall												D	D

Source: Metropolitan Council and STORET data



Edith Lake (82-0004) Valley Branch Watershed District

Edith Lake is a 81-acre lake located within Afton (Washington County). The lake has a maximum depth of depth of approximately 13.0 m (43 feet). Roughly 42 percent of the lake's surface area is considered littoral zone (the 0-15 foot depth area of aquatic plant dominance).

Additionally, the lake has a 1,576-acre immediate drainage area, which results in a watershed-to-lake area ratio of 19:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This marks the first year in which Edith Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Therefore, 2004 is the only known year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored six times between mid-April and early-September, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary


<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	20.3	16.0	24.0	A
CLA (µg/l)	5.0	2.0	8.0	A
Secchi (m)	2.7	2.3	3.1	B
TKN (mg/l)	0.69	0.68	0.70	
<i>Overall Grade</i>				A

As mentioned earlier, there are no water quality data available for Edith Lake other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.0 for physical condition (2- "some algae present"), and 1.8 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	15.9				20	39		2.9	2	1
4/24/05	13.3				5.4	22		2.7	2	1
6/7/05	24.9				2	16		3.1	2	1
7/4/05	25.1				6.3	24		2.3	2	2
8/14/05	26.1				3.5	21		3.0	2	2
9/4/05	23.7				8	20		2.5	2	2

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophylla													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													A
Chlorophylla													A
Secchi Depth													B
Overall													A

Source: Metropolitan Council and STORET data



Elmo Lake (82-0106) Valley Branch Watershed District

Lake Elmo, located in Lake Elmo (Washington County), has public access associated with the Lake Elmo Regional Park located on the west side of the lake. The lake is considered a “Priority Lake” due to its multi-recreational uses. The 284-acre lake has a maximum depth of 41.7 m (roughly 140 feet [deepest in the TCMA]). Roughly 22 percent of the lake’s surface area is considered littoral zone (the 0-15 foot depth area of aquatic plant dominance).

The lake was monitored nine times from early-May to early-October, 2005.

This was the second year that Lake Elmo has been involved in CAMP (the other being 1994). The lake has been monitored in the past by Council staff (most recently in 1991).

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	22.5	8.0	68.0	A
CLA (µg/l)	2.7	1.6	6.3	A
Secchi (m)	4.1	2.5	5.5	A
TKN (mg/l)	0.46	0.30	0.73	
Overall Grade				A

The lake’s 2005 overall water quality grade is identical to those recorded in 1981, 1988, 1991, and 1994, and better than those recorded in 1980, 1982, and 1984 (B).

A search of the STORET nationwide water quality database for data on the lake revealed a moderate database since the 1980’s with nutrient and Secchi transparency data available in 1980-1982, 1984, 1988, 1991, 1994 and 2005. Additionally, Secchi transparency data are available for 1985-1987, 1989-1990 and 1992-1993. The lake’s database indicates that the lake’s recent water quality is well represented by an overall grade of A. Additionally, a recent MPCA conducted trend analysis on the lake’s Secchi transparency data, revealed a statistically significant improvement in recent water clarity.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake’s physical and recreational conditions on a 1-to-5 scale. The summertime mean recorded physical condition was 2.0 (2- "some algae present"). The mean suitability for recreation ranking, also on a 1-to-5 scale, was 2.1 (between 2- " minor aesthetic problem" and 3- “swimming slightly impaired”).

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.


If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Elmo

Lake Elmo, Washington Co.

Lake ID: 820106
WD: Valley Branch
Volunteer: Wendy Griffin

● Sampling site
Contours in meters



2005 Data





Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	Chl. a ug/L	Surf. TP ug/L	Bot. TP ug/L	Secchi M	PC	RS
5/6/05	11.4				6.3	24		2.5	3	3
6/9/05	22.6				1.6	15		5	2	2
6/21/05	24.7				1.6	18		5.5	2	2
7/3/05	23.5				3	18		4	2	3
7/12/05	27.9				1.6	68		3.5	2	2
7/17/05	27					10		4	2	2
8/7/05	27.3				2.2	19		4	1	1
9/6/05	24				2.8	8		4.5	2	2
10/10/05	15.8				2.7	11		7	1	1

Lake Water Quality Grades Based on Summertime Averages

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B	A	B		B				B						A	
Chlorophyll a	B	A	B		A				A						A	
Secchi Depth	C	B	C		B	A	B	B	A	A	A	A	A	A	A	
Overall	B	A	B		B				A						A	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		A											A
Chlorophyll a		A											A
Secchi Depth	A	A											A
Overall		A											A

Source: Metropolitan Council and STORET data



Farquhar Lake (19-0023) City of Apple Valley

Farquhar Lake, located in the City of Apple Valley (Dakota County), covers an area of 63 acres and has a maximum depth of 3.0 m (10 feet). The lake's mean depth of 1.4 m (4.6 feet) and surface area translates to an approximate lake volume of 290 ac-ft (the lake volume may have changed over the past couple years due to the lake level rising 1.5 to 2.0 feet above normal). Because the maximum depth is only 3.0 m, the entire lake area is considered littoral (the area of aquatic plant dominance), and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).


The land uses within the 353-acre contributing watershed to the lake are approximately split between agricultural uses and urban/residential. The watershed-to-lake size ratio is 6:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff).

This was the eleventh year that Farquhar Lake has been enrolled in CAMP. The lake was monitored 15 times between mid-April and mid-October, 2005.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP ($\mu\text{g/l}$)	171.5	70.0	272.0	F
CLA ($\mu\text{g/l}$)	56.0	14.0	100.0	D
Secchi (m)	0.6	0.2	1.2	F
TKN (mg/l)	2.56	0.87	3.80	
Overall Grade				F

The lake's 2005 overall grade is identical to those recorded in 1999-2004, and worse than the C's recorded in 1994 and 1996, and the D's of 1995 and 1997. The 2005 CLA grade, however, is the best recorded since 1997.



The above graph clearly depicts the lake's recent (mid-1990s to present) degradation. In fact, a recent MPCA conducted trend analysis on the lake's Secchi transparency data, revealed a statistically significant decrease in recent water clarity. The reason for the degradation in the lake's water quality is not entirely

known. A more in-depth study combining watershed as well as in-lake monitoring may help determine the areas contributing the most to the lake's degradation.

Throughout the 2005 season, the volunteer monitor ranked their perceptions of the lake's physical and recreational condition on a 1-to-5 scale. The mean perceived physical condition was 2.8 (falling between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability was 2.9 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Farquhar Lake

Apple Valley, Dakota Co.

Lake ID: 190023
WMD: Dakota County
Volunteer: Rick Bruneau

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/14/05	14.7				41	102		1.1	2	1
5/1/05	14.2				14	92		1.2	1	1
5/14/05	17.3				28	90		1	2	3
5/18/05	18.2				29	70		0.9		
5/25/05	19.1				29	136		0.7	2	3
6/5/05	29.6				39	199		0.6	3	4
6/21/05	32.1				61	269		0.4	4	4
7/10/05	28.3				97	272		0.3	4	3
7/24/05	26.9				100	201		0.3	3	3
8/7/05	27.3				89	204		0.3	4	
8/21/05	30.8				93	182		0.2	4	4
9/4/05	27.6				51	147		0.3	3	3
9/17/05	25.4				15	196		1.1	1	1
10/1/05	24.7				8.1	128		1.1	2	1
10/15/05	17.8				15	106		1.3	2	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		C	D	D	D		F	F	F	F	D	F	F
Chlorophyll a		B	C	C	D		F	F	F	F	F	F	D
Secchi Depth		C	D	C	D		F	F	F	F	F	F	F
Overall		C	D	C	D		F	F	F	F	F	F	F

Source: Metropolitan Council and STORET data



Fireman's Lake (10-0226) Carver County Environmental Services

This was the fifth year that Fireman's Lake (located within the City of Chaska [Carver County]), has been involved in CAMP (the lake was first enrolled in 2001). The 8-acre lake has a maximum depth of 7.0 m (23 feet). Roughly 88 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has been reported on the lake.

A search through the STORET nationwide water quality database determined that the 2001-2005 CAMP data are the only years of available water quality data for the lake.

The lake was monitored 14 times from mid-April to mid-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	23.4	13.0	69.0	B
CLA (µg/l)	3.9	1.0	10.0	A
Secchi (m)	3.2	2.1	4.2	A
TKN (mg/l)	0.39	0.26	0.82	
Overall Grade				A

While the lake's 2005 overall grade is identical to those of 2001-2005, the 2005 individual summer means slipped from those recorded in 2004.

As mentioned earlier, there are no water quality data available for Fireman's Lake other than the limited 2001-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term however, the lake's water quality is well represented by an overall grade of A. To better understand the lake's water quality and where it may be heading, more data are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 1.4 for physical condition (between 1- "crystal clear" and 2- "some algae present"), and 1.1 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Fireman's Lake Chaska, Carver Co.


Lake ID: 100226

WMD: Carver County

Volunteer: Carver Co. Env. Services

● Sampling site

Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/14/05	14.5		11		2.8	19		2.1	2	1
4/28/05	11.3		11.5		2.1	14		3	1	1
5/9/05	17.9		11.5		2.8	19		3.7	1	1
5/23/05	18.6		9.5		3	13		3	1	1
6/6/05	23.2		5.56		1	69		2.4	2	1
6/22/05	25.7		8.56		4.5	29		2.5	2	1
7/6/05	25.6		13.52		3.2	28		2.2	2	2
7/20/05	28.4		10.32		4.6	15		2.1	2	1
8/2/05					2.3	17		3.8	1	1
8/16/05	25.6		6.7		3.5	17		3.8	2	1
8/29/05	25.6		10.22		3.2	17		4.2	1	1
9/13/05	23.6		8.7		4.8	15		3.4	1	1
9/27/05	19.8				10	18		3.9	1	1
10/11/05	14.6				6.9	37		2.4	1	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus									A	A	B	A	B
Chlorophyll a									A	A	A	A	A
Secchi Depth									B	A	A	A	A
Overall									A	A	A	A	A

Source: Metropolitan Council and STORET data



Fish Lake [Scott County] (70-0069) *Prior Lake - Spring Lake Watershed District*

Fish Lake is located in Spring Lake Township (Scott County). This was the eighth year that the 171-acre lake has been a part of CAMP. The lake's mean and maximum depth of 4.4 m (14 feet) and 8.5 m (28 feet) translates to an approximate volume of 2,468 ac-ft. Roughly 43 percent of the lake's surface area is considered littoral, that is, the 0-15 foot depth area of the lake dominated by aquatic vegetation. The lake has a 434-acre watershed that, when divided by the surface area of the lake results in a rather small watershed-to-lake size ratio of 2.5:1 (the larger the ratio the greater the potential stress on the lake from surface runoff). The lake is considered a Metropolitan Council "Priority Lake" due to its multi-recreational uses. The lake can be accessed on the northwestern end.

The lake was monitored 13 times between mid-April and mid-October, 2005. A search for historic water quality data through Council, MPCA, and STORET files resulted in a few years of data (1980, 1984, 1990, 1995, 1997 [only two monitoring events], and 1998-2005).

The collected data and resulting graphs showing TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	40.0	22.0	71.0	C
CLA (µg/l)	23.3	15.0	36.0	C
Secchi (m)	1.3	0.8	2.4	C
TKN (mg/l)	1.27	0.80	1.60	
Overall Grade				C

The lake has received overall grades of C in 1980, 1995, 1997-2000 and 2003-2005, overall grade of B in 2001 and D's in 1984 and 2002.

During each visit, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The mean physical condition ranking was 2.7 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.0 (2- "minor aesthetic problem").

Statistical analysis on the lake's water quality database did not detect any long-term trends, in the short-term however, the lake seems to be very well represented by an overall lake water quality grade of C/C+. To better determine if this indicates a possible trend or is simply a fluctuation within the lake's normal range, more data are needed.




The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of any erroneous lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Fish Lake Spring Lake Twp., Scott Co.


Lake ID: 700069
WD: Prior Lake-Spring Lake
Volunteer: Scott Peterson

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	13.2				15	46		2.3	2	1
5/3/05	11.1				15	31		2.4	2	2
5/15/05	12.4				16	47		1.7	2	2
5/19/05	15.8				19	40		1.5		
5/29/05	16.8				36	57		1.1	3	2
6/12/05	25.2				25	48		1.9	3	2
6/26/05	26.5				16	33		1.2	2	2
7/10/05	26.7				24	71		0.9	3	2
7/27/05	27				21	28		0.9	3	2
8/7/05	27.8				25	34		0.8	3	2
8/28/05	24.7				25	22		0.9	3	2
9/17/05	21.8				30	29		0.9	3	2
10/1/05	18.7				48	72		0.8	3	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C				D								
Chlorophyll a	C				D						C		
Secchi Depth	D				D						C		
Overall	C				D								

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		C		C	C	C	C	C	C	D	C	C	C
Chlorophyll a		C		C	C	C	C	C	B	C	C	C	C
Secchi Depth		D		C	C	C	C	B	B	D	B	C	C
Overall		C		C	C	C	C	C	B	D	C	C	C

Source: Metropolitan Council and STORET data



Fish Lake [Washington County] (82-0064) *Carnelian - Marine Watershed District*

Fish Lake is located in New Scandia Township in Washington County. The lake has a surface area of 72 acres, and a maximum and mean depth of 3.0 m (10 feet) and 1.5 m (5 feet), respectively. Because of the shallowness of the lake, its entire surface area is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation, and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The mean depth and surface area of the lake translates to an approximate volume of 360 ac-ft.

The lake's watershed area of 683 acres translates to a watershed-to-lake size ratio of 9.5:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff).

This was the fifth year that Fish Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake revealed a limited amount of data collected. Water quality data were found for 1998-2004 and now 2005.

The lake was monitored seven times between mid-April and early-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	88.0	54.0	113.0	D
CLA (µg/l)	39.2	8.8	79.0	C
Secchi (m)	1.0	0.6	1.7	D
TKN (mg/l)	1.68	1.10	2.00	
Overall Grade				D

The resulting overall grade for 2005 is identical to that recorded in 2002 and 2004, and better than those recorded in 1998-2001 and 2003 (F). The 2005 CLA mean is the lakes best recorded to date.

Because of the limitedness of the lake's water quality database, the determination of any long-term trends is not possible to determine. In the short-term, the lake seems well represent by the overall grade of D/F. To better understand the lake's water quality and what direction it may be heading, more years of data collection are needed.

The perceived physical and recreational conditions of the lake, recorded by the volunteers, were ranked on a 1-to-5 scale. The rankings are shown in both tabular and graphical form on the lake's associated information sheet. The mean physical condition ranking was 3.2 (between 3- "definite algae present" and 4- "high algae color"), while the mean recreational suitability ranking was 3.6 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Fish Lake

New Scandia Twp., Washington Co.


LAKE ID: 820064

WD: Carnelian-Marine

Volunteer: Washington Co. SWCD




● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/18/05	17.4	16.6	6.55	6.48	29	110		1.1	3	3
5/16/05		13.2	6.65	6.43	19	86		1.1	3	3
6/14/05	23.9	21.3	8.01	0.33	8.8	79		1.2	2	3
7/11/05			7.63	7.35	17	54		1.7	3	4
8/9/05	26.6	26.6	6.71	0.84	72	108		0.6	4	4
9/9/05	21.6	21.4	7.96	6.7	79	113		0.6	4	4
10/6/05	16.3		5.72		54	115		1.1	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus						F	F	D	D	D	D	D	D
Chlorophyll a						D	D	F	F	D	F	F	C
Secchi Depth						F	F	F	F	D	F	D	D
Overall						F	F	F	F	D	F	D	D

Source: Metropolitan Council and STORET data



Forest Lake [West Basin] (82-0159) *Comfort Lake-Forest Lake Watershed District*

Forest Lake is divided into three distinct basins; however, only the west basin was monitored through CAMP in 2005 (the middle and eastern basins were monitored by Council staff in 2005). Because of the lake's multi-recreational uses it is considered a "Priority Lake" in the Metropolitan Area. One problem that may possibly hinder future recreational activity on the lake, however, is Eurasian Water Milfoil (*Myriophyllum spicatum*), which has been reported in the lake.

The entire 2,249-acre lake is located within the City of Forest Lake (Washington County). The acreage of each basin is as follows: west basin= 1,109 acres, middle basin= 360 acres, and the east basin= 780 acres. While the lake as a whole has a maximum and mean depths of 11.5 and 3.4 m (38 and 11 feet), the western basin itself has a mean and maximum depth of 3.0 m and 6.7 m (10 and 22 feet). The total volume of the whole lake is 24,986 ac-ft, and depending on hydrologic conditions has an 8-12 year residence time. Roughly 68 percent of the lake's surface area is considered littoral, (the shallow [0-15 feet] area dominated by aquatic plants). The 4,285-acre watershed translates to a rather small watershed-to-lake area ratio of 2:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). The lake has nine public accesses, 14 inlets and one outlet.

This was the eleventh year that the west basin of Forest Lake has been involved in CAMP (the previous being 1993, and 1996-2004). In 2005, the west basin of Forest Lake was monitored 14 times between mid-April and mid-October. Results are presented on graphs and data tables on the following page.

2005 summer (May-September) data summary




<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	37.7	16.0	84.0	C
CLA (µg/l)	21.6	4.7	60.0	C
Secchi (m)	2.2	1.4	3.8	C
TKN (mg/l)	0.91	0.50	1.30	
Overall Grade				C

Given the volatility of the lake's annual water quality (the lake received overall water quality grades of C in 1984, 1986, 1988, 1991, 1992, 1999-2000, 2002-2003, and 2005 and B in 1989, 1997-1998, 2001 and 2004), no definitive long-trends can be determined at this time. 2005 marks the first year since 1988 where the lake received a C for all three water quality parameters. The lake's overall water quality fluctuates between and B and C, depending on annual climatological conditions.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The mean perceived physical condition of the west basin of Forest Lake was 3.0 (3- "definite algae present"), while the mean recreational suitability was 2.6 (between 2- "minor aesthetics problem" and 3- "swimming slightly impaired").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you detect any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/1/05	10.9	10.4	8.96	7.88	4.5	24		2.6	2	1
4/28/05	9.9	9.5	6.82	6.8	5	60		2.9	3	2
5/13/05	11.9	11.9	6.09	5.63	4.7	19		2.9	2	2
5/24/05	15.3	14.5	5.45	4.95	5.2	38		3.2	3	2
6/9/05	21.5	20.1	4.62	3.71	5.6	25		3.4	3	3
6/21/05	26.3	22.8	7.51	5.96	5.7	16		3.0	2	2
7/7/05	22.7	22.1	8.45	6.34	9.9	26		2.3	3	2
7/22/05	26.5	24.8	8.07	1.58	14	32		1.4	3	3
8/2/05	25.5	24.4	7.26	4.68	19	38		1.2	3	2
8/16/05	23.8	22.4	8.77	0.55	35	46		1.1	4	4
9/1/05	21.9	21.6	8.57	0.67	60	84		1.1	3	3
9/14/05	21.3	21.1	7.6	0.46	39	48		0.8	3	3
9/30/05		16	8.68	0.72	40	43		0.8	4	3
10/1/05	13	12.9	7.87	7.08	23	39		1.2	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus				C		C	C	C	C	B		C	
Chlorophyll a				C		C			C	B	C	B	
Secchi Depth				C		C	C	C	C	C	C	C	
Overall				C		C			C	B		C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C		C	B	B	C	C	B	C	C	B	C	
Chlorophyll a	B		B	B	B	B	B	B	B	B	A	C	
Secchi Depth	C		C	C	C	C	C	C	C	C	B	C	
Overall	C		C	B	B	C	C	B	C	C	B	C	

Source: Metropolitan Council and STORET data



Gaystock Lake (10-0031) Carver County Environmental Services

This year marks the fourth year of CAMP monitoring on Gaystock Lake, which is located in Dahlgren Township (Carver County). Other than the 1999 (only two monitoring events), 2000, 2001 and 2005 CAMP data, a search for any historical water quality data came up empty. The 105-acre lake has a maximum depth of 5.0 m (16 feet). Because of the shallowness of the lake, the majority of its area is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation), and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake does not have a public access.

The lake was monitored 14 times between late-April and mid-October, 2005. Results are presented on graphs and data tables on the following page. During each monitoring event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as its perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	217.0	112.0	337.0	F
CLA (µg/l)	187.5	21.0	360.0	F
Secchi (m)	0.3	0.1	0.8	F
TKN (mg/l)	3.42	1.50	5.60	
Overall Grade				F

The collected data and resulting graphs showing TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake's information sheet on the following page.


Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 3.5 (between 3- "definite algal presence" and 4- "high algal color"), while the mean recreational suitability ranking was 3.7 (between 3- "swimming slightly impaired" and 4- "no swimming - boating ok").

Because 2005 was only the third year of collected data (no grades were determined in 1999 because of the lack of sufficient data), no long- or short-term trends can be determined. To better understand the lake's quality and what direction it may be heading, more years of data collection are needed.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Gaystock Lake Dahlgren Twp., Carver Co.

Lake ID: 100031
WMO: Carver County
Volunteer: Carver Co. Env. Services



2005 Data





Date	Surf Temp	Bot Temp	Surf DO	Bot DO	CIA	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ig/L	ig/L	ig/L	M	1-5	1-5
4/23/05	15.4		11.17		69	192		0.5	2	4
4/28/05	10.1		10.5		52	163			3	4
5/1/05	15.9		8.46		21	161		0.8	3	4
5/21/05	17		8.6		52	134		0.6		
6/5/05	22.9		8.21		58	112		0.5	3	3
6/23/05	25.3		15.26		71	152		0.3	4	3
7/5/05	26				210	272		0.2	3	4
7/21/05	25.9		10.24		230	228		0.1	4	4
8/2/05					280	228		0.2	4	4
8/17/05	24.3		9.54		360	266		0.2	4	4
8/30/05	23.6		13.1		340	175		0.2	3	4
9/15/05	20.9		4.52		230	337		0.2	4	4
9/27/05	18.3				210	322		0.2	3	3
10/12/05	12.6				45	287		0.5	3	2

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus									C	F			F
Chlorophyll a									F	F			F
Secchi Depth									F	D			F
Overall									D	F			F

Source: Metropolitan Council and STORET data



George Watch Lake (2-0005) Rice Creek Watershed District

This was the tenth year that George Watch Lake, located in the City of Lino Lakes (Anoka County), has been enrolled in CAMP. The lake was monitored 12 times from mid-April to mid-September, 2005. The 528-acre lake, which has a canoe access on its eastern side, has a mean and maximum depth of 1.5 m (5 feet) and 2.0 m (6.5 feet). The lake's approximate volume is 2,587 ac-ft and because of the shallowness of the lake, it is entirely littoral zone (the area of aquatic plant dominance) and never develops and maintains a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column) through the summer months. The major land use within the lake's immediate watershed is undeveloped/park.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	225.6	46.0	485.0	F
CLA (µg/l)	43.1	4.6	130.0	C
Secchi (m)	0.65	0.20	1.10	F
TKN (mg/l)	3.33	0.77	7.0	
Overall Grade				D

The lake's data reveal overall grades of D in 1982-1983, 1985, 1987-1988, 1990, 1997, 1999-2000, 2003, and 2005 and F in 1981, 1989, 1991, 1996, 1998, 2001-2002, and 2004.

A search through the STORET database for historic data on George Watch showed that the lake has been monitored several times in the past. There are nutrient data available for 1981-1983, 1985-1991, and 1996-2005. The lake's overall lake water quality grades seem to indicate that the lake water quality has remained fairly constant fluctuating between an F and D grade throughout the 20+ years of data. The TP and Secchi data has remained fairly consistent throughout the monitoring years, but the CLA seems to fluctuate greatly.

A reason for the fluctuating CLA means while the Secchi and TP numbers remain fairly constant could be the amount of sedimentation that could at times be limiting the amount of light available for algal growth thus keeping CLA low and vice versa

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The summertime mean physical condition was 3.0 (3- "definite algae present"). The mean suitability for recreation ranking was 3.9 (roughly equal to 4- "no swimming - boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you detect any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

George Watch Lake


Lino Lakes, Anoka Co.

Lake ID: 20005

W/D: Rice Creek

Volunteer:
Wargo Nature
Center

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	14				16	40		1.1	1	4
5/14/05	15				9.5	46		1.1	1	4
5/17/05	16.2				10.5	50		1.1		
5/27/05	21				16	101		1.1	1	4
6/9/05	28				17	139		1.0	2	4
6/21/05	27				11.8	154		1.0	3	4
7/25/05	26				27	320		0.4		
7/27/05	26				34	279		0.2	4	3
8/12/05	28				93	473		0.3	4	4
8/24/05	22				100	485		0.3	4	4
9/6/05	23				7.5	362		0.3	4	4
9/13/05	22				130	72		0.3	4	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	F	F	F	F	F	F	F	F	F	F	F	F	F
Chlorophyll a	F	C	B	B	C	B	D	C	F				
Secchi Depth	F	D	F	F	F	F	F	F	D	F			
Overall	F	D	D	D	D	D	D	F	D	F			

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			F	D	F	D	D	F	D	F	F	F	F
Chlorophyll a			D	C	D	C	C	F	D	C	D	C	C
Secchi Depth			F	F	F	D	F	D	F	D	F	F	F
Overall			F	D	F	D	D	F	D	D	F	D	D

Source: Metropolitan Council and STORET data



German Lake (82-0056) Carnelian – Marine Watershed District

German Lake is a 109-acre lake located in New Scandia Township (Washington County). There is very little known morphological data available for the lake.

This was the fourth year that German Lake has been involved in CAMP. A search through the STORET nationwide water quality database determined that the 2002-2005 CAMP data are the only years of available water quality data for the lake.

As part of the watershed's involvement in CAMP in 2005, the lake was monitored seven times between mid-April and early-October. During each sampling event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	25.2	18.0	40.0	B
CLA (µg/l)	4.8	2.6	6.5	A
Secchi (m)	2.4	2.1	2.6	B
TKN (mg/l)	0.70	0.51	0.84	
<i>Overall Grade</i>				B

The lake 2005 overall water quality grade is similar to those recorded in 2002-2004.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the following page. The mean physical condition ranking was 2.0 (2- "some algae present"), while the mean recreational suitability ranking was 3.8 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").


As mentioned earlier, there are no water quality data available for German Lake other than the 2002-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term however, the lake's water quality is well represented by an overall grade of B. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

German Lake Forest Lake, Washington Co.

Lake ID: 820056
WMO: Carnelian-Marine
Volunteer: Wash. Co. SWCD

◆ Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/18/05	15	14.8	5.53		7	50		2.3	2	3
5/16/05	12.7	12	6.19	5.91	5.8	40		2.3	2	4
6/15/05	23	22.7	6.52	6.38	6.5	28		2.6	2	4
7/11/05			9.13	7.46	2.6	18		2.4	2	4
8/12/05	24.2	24.2	5.3	4.98	5	18		2.3	2	4
9/9/05	21.5	21.4	7.52	6.8	4.1	22		2.1	2	3
10/6/05	15.3	15.3	7.08	6.93	4.5	17		2.4	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus									B	B	B	B	B
Chlorophyll a									A	A	A	A	A
Secchi Depth									C	B	B	B	B
Overall									B	B	B	B	B

Source: Metropolitan Council and STORET data



Goetschel Lake (82-0313) Valley Branch Watershed District

Goetschel Lake is located in Grant Township (Washington County). This was the fourth year that the 22-acre lake has been a part of CAMP. The lake's mean and maximum depth of 1.2 m (4 feet) and 4.2 m (14 feet) translates to an approximate volume of 88 ac-ft. Because of the shallowness of the lake, its entire surface area is considered littoral, that is, the 0-15 foot depth area of the lake dominated by aquatic vegetation. The lake has a 2,812-acre watershed that, when divided by the surface area of the lake results in a large watershed-to-lake size ratio of 122:1 (the larger the ratio the greater the potential stress on the lake from surface runoff).

A search through the STORET nationwide water quality database determined that the 2002-2005 CAMP data are the only years of available water quality data for the lake.

As part of the watershed's involvement in CAMP in 2005, the lake was monitored fourteen times between mid-April and early-October. During each sampling event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	35.0	24.0	52.0	B
CLA (µg/l)	7.4	1.3	13.0	B
Secchi (m)	1.9	1.2	2.4	C
TKN (mg/l)	0.74	0.52	1.00	
Overall Grade				B

The lakes 2005 overall water quality grade is identical to those recorded in 2002-2004.

Throughout the monitoring period, the volunteers' opinions of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the following page.

The mean physical condition ranking was 2.8 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 3.8 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

As mentioned earlier, there are no water quality data available for Goetschel Lake other than the 2002-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term however, the lake's water quality is well represented by an overall grade of B. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Goetschel Pond

Lake Elmo, Washington Co.




Lake ID: 820313

W/D: Valley Branch

Volunteer: John Longtin


● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	15.7				4.9	34		2	2	4
5/3/05	10.1				4.7	38		1.9	2	3
5/16/05	12.5				5.1	40		2.1	2	3
5/18/05	15				6.2	28		2		
5/26/05	17.8				2.9	44		1.8	2	3
6/12/05	26.9				1.3	24		1.9	3	4
6/26/05	27				13	26		1.8	3	4
7/10/05	26.9				7.5	24		1.2	3	4
7/24/05	27.4				12	52		1.8	3	4
7/31/05	26.3				5.8	44		1.8	3	4
8/14/05	24.1				11	39		1.7	3	4
8/28/05	23				6	31		2.4	3	4
9/11/05	23.5				10	32		2	3	4
9/25/05	19.6				10	33		2	3	4
10/9/05	12.9				7.5	36		2.3	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus										C	C	B	C
Chlorophyll a										A	A	B	A
Secchi Depth										C	B	C	C
Overall										B	B	B	B

Source: Metropolitan Council and STORET data



Goggins Lake (82-0077) Browns Creek Watershed District

Goggins Lake is an 11-acre lake located within May Township (Washington County). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This was the seventh year that Goggins Lake has been involved in CAMP (1999 being the first). Other than the CAMP data, a search through the STORET nationwide water quality database for historical water quality data for the lake came up empty. The lake was monitored 14 times between mid-April and mid-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	83.2	42.0	154.0	D
CLA (µg/l)	45.1	5.6	88.0	C
Secchi (m)	1.2	0.5	2.7	C
TKN (mg/l)	1.68	0.84	2.5	
Overall Grade				C

The 2005 overall grade is identical to that recorded 1999 and 2003-2004, and better than the D's recorded in 2000-2002. Results are presented on graphs and data tables on the following page.

Statistical analysis on the lake's water quality database did not detect any long-term trends. In the short-term however, the lake's water quality is well represented by an overall grade of D+/C. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.




The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.8 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.1 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Goggins Lake May Twp., Washington Co.


LakeID: 820077
WMO: Browns Creek
Volunteer: Wash. Co. SWCD

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/15/05	12.4	10.3	6.83	0.25	17	104		1.7	3	4
4/28/05	11	9.9	6.74	2.69	10	52		1.5	2	2
5/9/05	15.3	10.3	6.32	2.76	5.6	42		2.1	3	4
5/26/05	15.8	13.4	5.3	2.45	6.5	44		2.7	3	4
6/8/05	22.1	15.6	4.74	0.23	8.8	45		2.7	3	4
6/23/05	25.8	16.7	8.93	0.32	23	54		1.8	4	4
7/5/05	23.1	17.7	9.36	0.09	67	92		0.8	4	4
7/25/05	26.4	18.3	10.68	0.62	88	102		0.6	4	4
8/5/05	24.8	19.5	4.94	0.84	74	94		0.5	4	4
8/17/05	23.5		9.17	0.42	45	76		0.6	5	5
8/30/05	22.5	20.7	10.07	0.41	36	82		0.6	4	4
9/15/05	21.1	21.1	7.03	5.26	80	130		0.5	4	4
9/30/05	16.2	16.1	9.51	0.61	62	154		0.5	4	4
10/14/05	14	13.6	8.27	0.4	31	83		0.6	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	D	D	D	C	C	D
Chlorophyll a							C	C	C	C	C	C	C
Secchi Depth							C	D	D	D	C	D	C
Overall							C	D	D	D	C	C	C

Source: Metropolitan Council and STORM data



Golden Lake (2-0045) Rice Creek Watershed District

Golden Lake, located in the City of Circle Pines (Anoka County), was monitored 11 times between early-May and mid-October, 2005. Public access to the 57-acre lake (1.5 miles in circumference) is possible for non-motorized boats through Golden Lake County Park. The mean and maximum depths of the lake are 2.5 m (8 feet) and 7.3 m (24 feet), respectively. The lake's size and mean depth results in an approximate lake volume of 460 ac-ft. Roughly 42 percent of the lake is considered littoral zone, that is, an area of aquatic plant dominance.

The lake's 7,680-acre watershed translates to a large watershed-to-lake size ratio of 135:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

On each sampling date, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	61.0	29.0	121.0	C
CLA (µg/l)	25.2	2.5	55.0	C
Secchi (m)	1.9	0.8	3.2	C
TKN (mg/l)	2.00	1.50	2.70	
<i>Overall Grade</i>				C

The physical and recreational conditions of Golden Lake as perceived by the volunteer(s) were ranked on a 1-to-5 scale. These rankings are shown on the lake's information sheet on the next page. The summertime mean physical condition was 1.9 (between 1- "crystal clear" and 2- "some algae present"). The mean suitability for recreation ranking, was 1.4 (between 1- "beautiful" and 2- "minor aesthetic problem").

Golden Lake has a fairly extensive water quality database with Secchi and nutrient data for 1980-1981, 1984-1991, and 1993-2005. Because the lake's water quality grade has fluctuated between C, D, and F (a C in 1985-1987, 1996, 1998-2000, and 2005, D in 1980-1981, 1993, 1997 and 2001-2004, and an F in 1988-1991) throughout these 20+ years of monitoring data, no long-trends can be determined. It seems that the lake has a very wide fluctuation range in its water quality. In order to detect any possible long-term trends, more years of data collection are needed.



The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you recognize any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Golden Lake Circle Pines, Anoka Co.




Lake ID: 20045
W/D: Rice Creek
Volunteer: City of Circle Pines

• Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/2/05	9.5				9.4	53		2.3	1	1
5/17/05	13.8				8.5	30		2.2		
5/20/05	14.6				5.7	58		2.3	1	1
5/31/05	20.2				4	29		3.2	1	1
6/16/05	21.3				2.5	37		3.2	1	1
7/6/05	25.3				15	37		1.8	2	1
7/27/05	23.8				55	65		0.8	3	2
8/16/05	24.3				45	79		1	3	2
8/30/05	25.4				41	101		0.8	3	2
9/20/05	23.5				48	121		1.3	2	2
10/13/05	14.6				130	101		1.1	2	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C	D			D	F	C	F	D	D	D	D	
Chlorophyll a	D					C	C	D	F	F	F	F	
Secchi Depth	D	D				C	C	C	F	F	F	F	
Overall	D					D	C	D	F	F	F	F	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	D			C	D	C	C	C	D	D	D	D	C
Chlorophyll a	D			C	C	C	C	C	D	D	C	C	C
Secchi Depth	D			D	D	D	D	C	D	D	D	D	C
Overall	D			C	D	C	C	C	D	D	D	D	C

Source: Metropolitan Council and STORET data



Goose Lake [Scandia] (82-0059) Comfort Lake-Forest Lake Watershed District

Goose Lake, an 83-acre lake (1.9 miles in circumference) located in New Scandia Township (Washington County), was monitored seven times from mid-April to early-October, 2005. Goose Lake was enrolled in CAMP in 1994-1998 and 2004. The lake has a maximum and mean depth of 7.6 m (25 feet) and 2.4 m (8 feet), respectively. The lake's mean depth and size translate to a lake volume of approximately, 664 ac-ft. Because of the shallowness of the lake, roughly 98 percent of the lake is considered littoral (the area of aquatic vegetation dominance). A Public access is located on the western side of the lake.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP (µg/l)	65.2	34.0	146.0	C
CLA (µg/l)	40.4	6.7	110.0	C
Secchi (m)	2.16	0.61	3.81	C
TKN (mg/l)	1.40	1.10	2.00	
Overall Grade				C

The lake's 2005 overall grade was the same as those recorded in 1994-1998 and 2004.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions was ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The mean physical condition ranking was 3.0 (3- "definite algae present"), while the mean recreational suitability ranking was 3.4 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

Because 1994-1998 and 2004-2005 are the only years of available data, no long-term trends can be determined. On the short-term, however, the lake's overall water quality seems to be represented quite well by an overall grade of C. There is some normal fluctuation in each parameters annual means, however. To better understand the quality of the lake and what direction it may be heading, more years of data collection are needed.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Goose Lake,

New Scandia Twp., Washington Co.

Lake ID: 820059

WMO: Comfort Lake - Forest Lake




Volunteer: Wash. Co. SWCD

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/18/05	15.3	7.9	6.02	5.93	5.3	67		3.0	2	2
5/16/05	12.7	11.9	5.93	5.18	6.7	48		3.8	3	3
6/15/05	22.6	13.3	7.47	0.39	7.4	35		3.4	2	3
7/11/05	24.9	15	10.3	0.1	18	34		2.1	3	3
8/9/05	25.9	16.2	6.51	0.51	60	146		0.9	3	4
9/7/05	22.4	18.1	11.7	9.37	110	63		0.6	4	4
10/3/05	18.2	16.4	9.37	0.55	160	103		0.5	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		C	D	C	C	C						C	C
Chlorophyll a		C	B	C	C	C						C	C
Secchi Depth		D	C	C	C	C						B	C
Overall		C	C	C	C	C						C	C

Source: Metropolitan Council and STORET data



Goose Lake [Waconia] (10-0089) Carver County Environmental Services

Goose Lake, located in Waconia Township in Carver County, was monitored 14 times between mid-April and mid-October, 2005. The lake has been involved in CAMP since 1995. Because the maximum depth of the 407-acre lake is only 3.0 m (10 feet), the entire lake area is considered littoral zone (the 0-15 foot depth area of the lake dominated by aquatic vegetation). Additionally, because of the lake's shallowness it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake's mean depth of 1.5 m (roughly 5 feet) and its surface area translate to an approximate lake volume of 2,035 ac-ft.

The lake has a 1,100-acre immediate watershed, which translates to a watershed-to-lake area ratio of 27:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff). A 1999 water quality report on water resources in Carver County estimates land use for the watershed at: four percent residential, 61.0 percent agricultural, and 35.0 percent open/undeveloped (Carver County Planning 1999).

On each sampling date, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	109.8	47.0	242.0	D
CLA (µg/l)	108.5	29.0	200.0	F
Secchi (m)	0.4	0.2	0.5	F
TKN (mg/l)	2.32	0.91	3.70	
Overall Grade				F

The physical and recreational conditions of Goose Lake as perceived by the volunteer were ranked on a 1-to-5 scale. These rankings are shown on the lake's information sheet on the next page. The mean physical condition ranking was 3.3 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability ranking was 3.2, (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

Because of the variability among the nine years of data (grades ranging from C to F), no long-term trends can be determined. In the short-term however, the lake fluctuates greatly, with an overall grade of C in 1996 and 1998, D in 1995, 1999, 2001-2002, and 2004, and an overall grade of F in 1997, 2000, 2003 and 2005.


To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Goose Lake, Waconia Twp., Carver Co.

• Sampling site
Contours in meters



2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 to 5	1 to 5
4/18/05	15.2		10.3		45	117		0.6	3	3
4/27/05	10.2		10.66		46	174		0.7	3	3
5/11/05	15.7		10.85		66	98		0.4	3	3
5/24/05	19.7		11.01		79	86		0.5	3	3
6/6/05	21.7		8.24		29	242		0.5	4	3
6/20/05	24.6		9.22		48	177		0.5	3	3
7/7/05	23.8		11.2		68	77		0.4	4	4
7/8/05	26.8		6.16		93	61		0.4	4	4
8/1/05	23.7				130	105		0.2	3	3
8/18/05	23.5		7.21		200	47		0.3	3	3
8/31/05	23		8.39		180	148		0.4	3	3
9/13/05	22.9		7.61		160	69		0.3	3	3
9/21/05	22.1				140	98		0.4	3	3
10/10/05	12.2				63	103		0.5	3	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			D	C	F	D	D	F	D	D	F		D
Chlorophyll a			C	C	D	C	D	F	C	C	F		F
Secchi Depth			F	C	F	C	F	F	D	F	F		F
Overall			D	C	F	C	D	F	D	D	F		F

Source: Metropolitan Council and STORET data



Hafften Lake (27-0199) Pioneer-Sarah Watershed Management Commission

Hafften Lake, located in Greenfield (Hennepin County), has public access on the eastern side of the lake. The 43-acre lake has a maximum depth of 13.4 m (roughly 44 feet). Roughly 60 percent of the lake's surface area is considered littoral zone (the 0-15 foot depth area of aquatic plant dominance).

The lake was monitored 15 times from mid-April to mid-October, 2005.

While this was the first year that Hafften Lake was monitored through CAMP, the lake has been monitored by Council staff in the past (most recently in 2004). A search of the STORET nationwide water quality database for data on the lake revealed a limited database with nutrient and Secchi transparency data available in 2000-2001, and 2004-2005.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	48.6	31.0	86.0	C
CLA (µg/l)	20.9	4.0	98.0	C
Secchi (m)	1.9	0.8	3.4	C
TKN (mg/l)	1.33	1.00	1.70	
Overall Grade				C

While the lake's 2005 overall grade is the same as those recorded in 2000-2001 and 2004, the individual parameter means seem to indicate that 2005 represents the lake's best monitored water quality year to date.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions was ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The mean physical condition ranking was 2.6 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 3.4 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

Because 2000-2001 and 2004-2005 are the only years of available data, no long-term trends can be determined. On the short-term, however, the lake's overall water quality seems to be well represented by an overall grade of C. To better understand the quality of the lake and what direction it may be heading, more years of data collection are needed.

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Hafften Lake,

Greenfield, Hennepin Co.



LAKE ID: 270199

W/D: Pioneer-Sarah Creek

Volunteers: Todd Fellman and
Jim Van Someren

● Sampling site

Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/17/05	15.8				2.6	52		3	1	2
5/3/05	11.8				7.8	50		3.1	2	2
5/16/05	13.7				5.4	48		3.4	2	2
5/25/05	18.5				6.5	31		3.2		
5/30/05	20.5				5.3	33		3	2	2
6/12/05	26.2				4	38		2.8	2	2
6/26/05	27				25	60		1.3	3	2
7/11/05	29.2				18	51		1.1	3	3
7/27/05	25.5				17	86		1	3	3
8/7/05	27.6				17	58		1	3	3
8/21/05	24.3				17	31		1	3	3
9/5/05	23.9				15	35		1.2	3	3
9/18/05	23.5				98	62		0.8		3
10/2/05	19.6				47	54		0.9	3	3
10/11/05	14.1				38	117		1.4	3	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus									C	C		C	C
Chlorophyll a									C	C		C	C
Secchi Depth									C	C		D	C
Overall									C	C		C	C

Source: Metropolitan Council and STORET data



Half Breed Lake [Sylvan] (82-0080) *Comfort Lake-Forest Lake Watershed District*

Half Breed Lake (also known as Sylvan Lake) is a 75-acre lake located in Forest Lake Township (Washington County). The lake's mean and maximum depth of 1.7 m (5.6 feet) and 10.3 m (34 feet) translates to an approximate volume of 420 ac-ft. Roughly 67 percent of the lake's surface area is considered littoral, that is, the area dominated by aquatic vegetation. The lake has a 303-acre watershed which, when divided by the surface area of the lake results in a rather small watershed-to-lake size ratio of 4:1 (the larger the ratio the greater the potential stress on the lake from surface runoff). The lake has no inlets and no public access to the lake.

Half Breed Lake was monitored 14 times from mid-April to mid-October, 2005. The collected data and resulting graphs showing the seasonal variability in TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake's information sheet on the following page.

Similar to prior years, the lake's data and graphs reveal that the water quality of Half Breed Lake rates in the top 10 percent of lakes again in the area in 2005.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	20.6	16.0	32.0	A
CLA (µg/l)	3.6	2.6	4.5	A
Secchi (m)	4.8	3.4	6.1	A
TKN (mg/l)	0.68	0.46	0.98	
Overall Grade				A

The lake's 2005 overall grade is identical to those recorded in 1987-1989, 1991, 1993, 1996, and 1998-2002-2004, and better than the overall grade of B recorded in 1986. Historic water quality data and resulting lake quality grades indicate that the lake has maintained its high quality over the past 20+ years. Additionally, the MPCA recently conducted a trend analysis on the lake's Secchi transparency data, which revealed a statistically significant improvement in recent water clarity.

Throughout the monitoring period, the volunteer's opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The mean physical condition ranking was 2.0 (2- "some algae present"), while the mean recreational suitability ranking was 1.8 (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.stat.mn.us.

Half Breed Lake

(Sylvan Lake)



Forest Lake Twp./
New Scandia Twp.,
Washington Co.

LAKE ID: 820080

WMO: Comfort Lake - Forest Lake

Volunteer: Washington Co.
SWCD

● Sampling station
Contours in meters



2005 Data





Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	Chl μg/L	Surf. TP μg/L	Bot. TP μg/L	Secchi M	PC 1-5	RS 1-5
4/14/05	11.8	5	5.97	0.19	5.3	27		4.3	2	2
4/25/05	12.2	5.1	6.01	0.21	5.1	19		4.7	2	2
5/13/05	12.5	5.5	5.96	0.16	3.9	17		4.9	2	2
5/24/05	17.5	5.6	5.61	0.06	4.2	37		5.0	2	2
6/6/05	24.1	5.9	4.49	0.37	4.5	26		4.4	2	3
6/23/05	25.9	6.1	8.76	0.74	4.1	18		3.4	2	1
7/1/05	24.1	6.3	9.08	0.66	3.7	20		5.0	2	2
7/22/05	27.7	6.4	7.96	0.68	3.2	17		4.0	2	2
8/2/05	27.3	6.4	6.94	0.57	2.6	20		3.8	2	1
8/16/05	25.2	6.4	7.17	0.46	2.9	16		4.5	2	2
9/1/05	22.6	6.7	7.12	0.59	3.3	16		6.1	2	2
9/14/05	22.2	7	6.64	0.44	3.7	19		5.2	2	2
9/30/05	16.1	7.2	8.08	0.72	3.3	20		6.1	2	1
10/14/05	13.5	7.2	8.42	0.57	3.1	4		5.0	2	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	B	A					C	B	A	A		A	
Chlorophyll a							B	A	A	A		A	
Secchi Depth	A	A	A	A	A	A	A	A	A	A	A	A	A
Overall							B	A	A	A		A	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	A			A		A	A	A	A	A	A	A	A
Chlorophyll a	A			A		A	A	A	A	A	A	A	A
Secchi Depth	A			A		A	A	A	A	A	A	A	A
Overall	A			A		A	A	A	A	A	A	A	A

Source: Metropolitan Council and STORET data



Hart Lake (02-0081) Rice Creek Watershed District

Hart Lake is an eight-acre lake located within the city of Columbia Heights (Anoka County). There is very little known morphological data available for the lake.

This marks the second year in which Hart Lake has been involved in CAMP (2004 being the first). A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Thus, 2004 is the only complete, year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored four times between late-May and late-July, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	195.3	156.0	238.0	F
CLA (µg/l)	76.3	15.0	220.0	D
Secchi (m)	0.4	0.3	0.6	F
TKN (mg/l)	4.15	2.20	5.50	
<i>Overall Grade</i>				D

As mentioned earlier, there are no water quality data available for Hart Lake other than the 2004-20045 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.8 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.0 for recreational suitability (4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Hart Lake


Columbia Heights,
Anoka County

LAKE ID: 20081

W/D: Rice Creek

Volunteer: Ray Muno





● Sampling station
Contours in meters



*Bathymetry
Unknown*

2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/23/05	22.4				26	156		0.6	3	4
6/21/05	23.5				44	229		0.4	4	4
7/14/05	24.6				220	158		0.4	4	4
7/22/05	31				15	238		0.3	4	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												NA	F
Chlorophyll a												NA	D
Secchi Depth												NA	F
Overall												NA	F

Source: Metropolitan Council and STORET data

Hay Lake (82-0065) *Marine on St. Croix Watershed Management Organization*

This was the seventh year of CAMP monitoring on Hay Lake, located in New Scandia Township (Washington County). The lake was monitored seven times between mid-April and early-October, 2005. The only known morphological data available for the 33-acre lake is its maximum depth (6.1 m [20 feet]). Other than the 1998-2001, 2003-2004, and 2005 CAMP data for the lake, a search for historical water quality data and any physical information came up empty.

During each monitoring event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented on graphs and data tables on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	96.2	78.0	124.0	D
CLA (µg/l)	82.8	28.0	140.0	F
Secchi (m)	0.9	0.6	1.4	D
TKN (mg/l)	1.70	1.20	2.10	
Overall Grade				D

The lake's overall 2005 lake quality grade of D was identical to those recorded in 1998-2001, and 2004, and worse than that recorded in 2003 (C).

Statistical analysis on the lake's water quality database did not detect any long-term trends. In the short-term however, the lake seems well represented with an overall water quality grade of D/C. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.




Throughout the monitoring period, the volunteer(s) ranked the perceived physical condition of the lake on a 1-to-5 scale. The mean perceived physical condition of Hay Lake was 2.8 (between 2- "some algae present" and 3- "definite algal presence"), while the mean recreational suitability was 3.4 (between 3- "swimming slightly impaired" and 4- "high algal color").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Hay Lake **New Scandia Twp.,** **Washington Co.**


LAKE ID: 820065
WMD: Marine-on-St Croix
Volunteer: Washington Co. SWCD

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1-5	1-5
4/15/05	16.9	16.8	6.14	5.85	47	88		1.1	3	2
5/16/05	13	12.8	6.64	6.32	45	90		0.9	2	3
6/14/05	21.7	19.7	7.87	0.39	120	98		0.6	3	3
7/11/05			8.3	0.14	81	91		0.8	3	4
8/9/05	25.8		5.94	0.41	140	124		0.6	3	3
9/7/05	21.7		7.89	0.55	25	78		1.4	3	4
10/3/05	18.3	18.2	5.86	0.54	15	55		1.2	2	3




Lake Water Quality Grades Based on Summertime Averages

Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus						D	D	D	D		D	D	D
Chlorophyll a						F	F	F	F		C	D	F
Secchi Depth						D	D	D	D		C	D	D
Overall						D	D	D	D		C	D	D

Source: Metropolitan Council and STORRET data



Hazeltine Lake (10-0014) Carver County Environmental Services

Hazeltine Lake is a 236-acre lake located within the City of Chaska (Carver County). The maximum depth of the lake is 2.0 m (roughly six-and-a-half feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This was the third year that Hazeltine Lake has been involved in CAMP (1999;[where it was only monitored twice] and 2000 were the others). Other than the past CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty.

The lake was monitored 14 times from mid-April to mid-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	176.9	80.0	247.0	F
CLA (µg/l)	208.6	26.0	430.0	F
Secchi (m)	0.3	0.1	0.7	F
TKN (mg/l)	4.34	2.30	6.50	
Overall Grade				F

Similar to that recorded in 2000 and 2001, the individual grades result in an overall water quality grade of F for Hazeltine Lake in 2005.

As mentioned earlier, there is very little water quality data available for Hazeltine Lake. Therefore it is impossible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, more data are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.6 for physical condition (between 3- "definite algal presence" and 4- "high algal color"), and 4.1 for recreational suitability (roughly equal to 4- "no swimming - boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.




Lake Hazeltine

Chaska, Carver Co.

LAKE ID: 100014


WMO: Carver County

Volunteer: Carver Co. Env. Series



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CIA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ig/L	ig/L	ig/L	M	1-5	1-5
4/14/05	14.2		13.8		96	106		0.8	2	4
4/28/05	9.4		11.9		96	115		0.5	2	4
5/10/05	15.8		10.5		80	247		0.4	3	4
5/23/05	19.4		9.15		63	141		0.4	3	4
6/7/05	25.7		11.18		26	80		0.6	2	4
6/22/05	25.1		8.99		64	130		0.7	4	5
7/5/05	23.5		10.62		72	178		0.4	4	4
7/20/05	26.4		9.64		230	188		0.2	4	5
8/2/05					390	145		0.1	4	4
8/16/05	24.4		15.9		400	228		0.2	4	4
8/29/05	24.9		14.08		430	195		0.1	4	4
9/15/05	20.5		8.98		300	210			4	4
9/25/05	18.4				240	204		0.2	4	4
10/11/05	12.3				72	144		0.3	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								F	F				F
Chlorophyll a								F	F				F
Secchi Depth								F	F				F
Overall								F	F				F

Source: Metropolitan Council and STO RET data



Henry Lake (27-0175) *Elm Creek Watershed Management Commission*

Henry Lake is a 77-acre lake located within Hassan Township (Hennepin County). Because the maximum depth of the lake is only 1.5 m (5 feet), the entire lake area is considered littoral zone (the 0-15 foot depth area of the lake dominated by aquatic vegetation). Additionally, because of the lake's shallowness it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This marks the second year that Henry Lake has been involved in CAMP (1995 being the other). Other than for the 1995 and 2005 CAMP data, a search through the STORET nationwide water quality database for historic data on the lake came up empty. Therefore, 1995 and 2005 is the only known year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 11 times between early-May and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	76.1	46.0	119.0	D
CLA (µg/l)	33.5	6.4	88.0	C
Secchi (m)	0.9	0.2	1.6	D
TKN (mg/l)	1.36	0.95	1.70	
Overall Grade				D

The lake's 2005 individual and overall grades are identical to those recorded in 1995.

As mentioned earlier, there are no water quality data available for Henry Lake other than the 1995 and 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.9 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.6 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Lake Henry





Hassan Twp., Hennepin Co.

Lake ID: 270175
WMO: Elm Creek
Volunteer: Tom Hoverson



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 to 5	1 to 5
5/3/05	10.6				11	53		1.3	2	5
5/17/05	19.9				6.4	51		1.6	3	5
5/31/05	21.4				7.2	46		1.5	3	5
6/14/05	24.1				20	52		1.0	5	4
7/5/05	26.9				25	101		1.0	5	5
7/19/05	30.5				26	97		0.8	5	5
8/2/05	31.5				28	90		0.8	5	5
8/30/05	23				73	67		0.2	3	4
9/13/05	23				88	85		0.2	4	4
9/27/05	20.2				50	119		0.3	4	4
10/14/05	15.1				16	98		0.3	3	5



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			D										D
Chlorophyll a			C										C
Secchi Depth			D										D
Overall			D										D

Source: Metropolitan Council and STORET data

Herber's Pond (82-0015-01) *Carnelian – Marine Watershed District*

Herber's Pond is a small (13-acre) shallow lake (a maximum depth of approximately 2.0 m (6.6 feet), located in Hugo (Washington County). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). There is very little other known morphological data available for the waterbody.

This was the second year that Herber's Pond has been involved in CAMP. The lake was monitored 14 times between mid-April and mid-October, 2005. On each of the sampling days the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	93.8	52.0	129.0	D
CLA (µg/l)	62.8	13.0	190.0	D
Secchi (m)	1.1	0.6	1.4	D
TKN (mg/l)	1.33	0.92	1.80	
Overall Grade				D

The lake's 2005 individual summer means and overall water quality grade is worse than those recorded in 2004.

As mentioned earlier, there are no known nutrient data available for Herber's Pond other than the 2004-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

The perceived physical and recreational conditions (ranked on a 1-to-5 scale) are shown on the lake's information sheet on the next page. The average user perception rankings, were 2.6 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 3.9 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Herber's Pond

Stillwater Twp.,
Washington Co.

LAKE ID: 820015-01
W/D: Carnelian-Marine
Volunteer: Washington Co. SWCD

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/11/05	15.5	7.5	5.62	3.25	30	149		0.6	3	4
4/27/05	10.8	10.8	4.64	4.03	18	110		1.1	3	4
5/11/05	16.5	12.2	5.13	1.49	24	70		1.4	2	4
5/24/05	22.1	14.9	4.6	5.45	13	64		1.4	3	4
6/6/05	25.6		4.33	3.91	18	52		1.4	2	4
6/23/05	25.6	21.9	11.67	0.28	69	97		0.6	3	4
7/8/05		17.9	7.99	0.11	40	111		1.4	2	4
7/25/05	25.2		4.56	0.5	190	104		1.2	2	3
8/5/05	25.3	22.3	3.09	0.47	54	117		0.9	3	4
8/19/05	22.2	21.6	3.54	0.39	68	93		1.2	3	4
8/31/05	22.4	20.6	6.98	1.37	120	129		1.1	3	4
9/15/05	20	19.6	1.9	0.34	52	124		0.8	2	4
9/28/05	17	16.9	7.77	4.25	43	71		0.9	3	4
10/11/05	13.8	11.8	6.21	4	47	111		1.1	2	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												D	D
Chlorophyll a												C	D
Secchi Depth												C	D
Overall												C	D

Source: Metropolitan Council and STORET data



Highland Lake (2-0079) Anoka County Parks

Highland Lake is a 22-acre lake located within the City of Columbia Heights (Anoka County). The maximum depth of the lake is approximately only 1.0 m (roughly 3 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This was the seventh year that Highland Lake has been involved in CAMP (the lake was initially enrolled in 1999). Other than the past CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty.


The lake was monitored 14 times between mid-April and mid-October, 2005. During each monitoring event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented on graphs and data tables on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP ($\mu\text{g/l}$)	303.3	154.0	418.0	F
CLA ($\mu\text{g/l}$)	223.4	65.0	460.0	F
Secchi (m)	0.3	0.2	0.6	F
TKN (mg/l)	3.01	0.94	4.30	
Overall Grade				F

The lake's recent water quality (2002-2005), is quite a bit worse than that recorded in 1999-2001. In fact, the lake's 2005 summer means are not only its worst recorded to date, they are the worst recorded of all the lakes in CAMP 2005.

As mentioned earlier, there are no water quality data available for Highland Lake other than the 1999-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term, however, the lake's water quality seems to be degrading. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.



The above graph clearly depicts the lakes recent degradation. The reason for the degradation in the lake's water quality is not entirely known. A more in-depth study combining watershed as well as in-lake monitoring may help determine the areas contributing the most to the lake's degradation.


The last two graphs on the information sheet show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception ranking, on a 1-to-5 scale, was 3.4 for physical condition (between 3- "definite algae present" and 4- "high algal color") and 3.8 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Highland Lake Columbia Heights, Anoka Co.




Lake ID: 20079
WMO: Six Cities
Volunteer: Anoka Co. Parks

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/11/05	14.2				69	168		0.5	3	
4/26/05	10.6				71	204		0.4	3	
5/12/05					83	223		0.6	3	
5/27/05	18.7				76	154		0.4	3	
6/9/05	21.5				120	251		0.4	3	3
6/20/05	26				170	248		0.4	3	4
7/6/05	24.3				280	324		0.2	4	4
7/19/05	27.4				460	340		0.2	3	4
8/3/05	24.7				400	418		0.3	4	4
8/15/05					340	374		0.2	4	5
8/30/05	17.4				240	336		0.2	4	4
9/15/05	18.6				65	365		0.5	3	2
10/5/05	18.8				56	222		0.4	2	2
10/12/05	13				24	65		0.4	2	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	D	D	F	F	F	F
Chlorophyll a							C	B	D	F	F	F	F
Secchi Depth							D	D	F	F	F	F	F
Overall							D	C	D	F	F	F	F

Source: Metropolitan Council and STORET data



Hydes Lake (10-0088) Carver County Environmental Services

Hydes Lake, a 215-acre lake located within Waconia Township (Carver County) is considered a Metropolitan Area “Priority Lake” because of its multi-recreational uses. A public access is located on the lake’s northeastern shore. The mean and maximum depth of the lake is 3.0 (roughly 10 feet) and 5.5 m (18 feet). Because of the shallowness of the lake, 88 percent of the total lake area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake’s water column). The lake’s surface area and mean depth result in an approximate lake volume of 2,150 ac-ft.

The lake has a 430-acre immediate watershed, which translates to a watershed-to-lake area ratio of 2:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff). A 1999 water quality report on water resources in Carver County estimates land use for the watershed at: seven percent residential, 76 percent agricultural, and 17 percent open/undeveloped (Carver County Planning 1999).

This was the seventh year that Hydes Lake has been involved in CAMP (the lake was initially enrolled in 1999). The lake has been monitored by Council staff in the past (the last year being 1996). A search of the STORET nationwide water quality database for data on the lake revealed a moderate database throughout the 1990’s with nutrient data available in 1985, 1991, 1993, 1996 and now 1999-2005.

The lake was monitored 14 times between mid-April and mid-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake’s perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	154.3	62.0	340.0	F
CLA (µg/l)	71.8	3.0	470.0	D
Secchi (m)	2.0	0.4	4.0	C
TKN (mg/l)	2.31	1.50	4.10	
Overall Grade				D




The lake’s 2005 overall grade is identical to those of worse recorded in 1985, 1991, 1993, 1996, 1999-2000, 2002, and 2004, and worse than that of 2001 and 2003 (C).

Statistical analysis on the lake’s water quality database did not detect any long-term trends. In the short-term however, the lake’s water quality seems to be well represented by an overall grade of D. In order to detect any possible long-term trends, additional years of data collection are needed.

The last two graphs show seasonal variation in the lake’s perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.4 for physical condition (between 3- “definite algae present” and 4- “high algal color”) and 3.3 for recreational suitability (between 3- “swimming slightly impaired” and 4- “no swimming – boating ok”).


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/18/05	14		9.44		3.7	66		3.9	1	2
4/22/05	10.6		9.4		3.8	127		2.3	2	2
5/11/05	14.7		10.15		7	81		1.9	2	2
5/24/05	17.6		9.4		4.1	62		4	3	4
6/6/05	21.1		10.15		6.2	96		2.6	4	4
6/20/05	24.9		8.66		15	168		2.6	4	4
7/7/05	24.9		15.91		130	139		1	3	3
7/18/05	27		6.34		470	140		0.9	3	3
8/1/05	23.7				23	144		1.9	5	4
8/18/05	24.2		5.52		17	135		2.3	3	3
8/31/05	23.3				3	210		3.1	3	2
9/13/05	22.6				40	182		1.5	4	4
9/23/05	20.3		4.33		75	340		0.4	3	3
10/10/05	13.5		7.42		110	279		1	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus						F						F	
Chlorophyll a						D						D	
Secchi Depth						D						D	
Overall						D						D	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	F		F				F	F	D	D	D	D	F
Chlorophyll a	C		C				C	C	C	C	C	D	D
Secchi Depth	C		C				C	C	C	F	C	D	C
Overall	D		D				D	D	C	D	C	D	D

Source: Metropolitan Council and STORET data



Island Lake (2-0022) *Anoka County Parks*

This was the third year of CAMP monitoring on Island Lake, which is located in Linwood Township (Anoka County). The lake has a surface area of 67 acres and a maximum depth of 6.7 m (22 feet). Roughly 87 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

A search through the STORET nationwide water quality database for data on the lake provided only one prior year of water quality data for the lake (1983) prior to the 2003-2005 CAMP data. The lake was monitored 14 times between mid-April and mid-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	33.1	17.0	97.0	C
CLA (µg/l)	14.6	6.3	63.0	B
Secchi (m)	1.8	1.3	2.4	C
TKN (mg/l)	0.79	0.20	1.40	
Overall Grade				C

The lake's overall 2005 lake quality grade of C is identical to that recorded in 1983 and worse than the B calculated from the 2003-2004 data.

Because 2005 is only the fourth year of available data, no long- or short-term trends can be determined. To better understand the quality of the lake and what direction it may be heading, more years of data collection are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.0 for physical condition (between 2- "some algae present") and 1.6 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").



The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of any errors in the lake's data/physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Island Lake Linwood Twp., Anoka Co.

Lake ID: 20022
WMD: Sunrise River
Volunteer: Anoka Co. Parks

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/11/05	13.2				11	52		1.4	2	
4/25/05	11.1				5.6	69		1.4	2	
5/12/05					8.9	97		1.5	2	
5/31/05					8.2	22		2.1	2	
6/9/05					63	20		2.4	2	1
6/21/05	24				6.3	40		2.4	2	3
7/6/05	24.2				10	41		2.0	2	1
7/19/05	29				11	21		1.4	2	1
8/3/05	24.6				7	18		2.0	2	3
8/15/05	23				11	17		1.3	2	2
8/30/05	17.9				8.4	25		1.5	2	1
9/15/05	21				12	30		1.7	2	1
10/5/05	17.8				15	54		1.2	2	1
10/12/05	13.4				30	43		1.4	2	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus				C									
Chlorophyll a				C									
Secchi Depth				D									
Overall				C									

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus											B	C	C
Chlorophyll a											B	A	B
Secchi Depth											C	C	C
Overall											B	B	C

Source: Metropolitan Council and STORM data



Jane Lake (82-0104) Valley Branch Watershed District

Lake Jane, which has a surface area of roughly 155 acres, is located in the northwest corner of the City of Lake Elmo (Washington County). The maximum and mean depths of the lake are 12.0 and 3.7 m (39 and 12 feet), respectively (roughly 72 % of the lake is considered littoral; the area of aquatic plant dominance). The approximate volume of the lake is 1,860 acre-feet (ac-ft) and its residence time (the estimated time it would take the lake to replenish itself if it were drained), is roughly 1.4 years. The size of the lake's immediate watershed is approximately 1,402 acres.

The lake has a public access located on its south end, which gets heavy use by area fishermen (the MNDNR manages the lake for largemouth bass, bluegill and crappie, and reports good reproduction) and boaters during the summer months. Furthermore, Lake Jane is considered a "Priority Lake" in the Metropolitan Area because of its multi-recreational uses.

This is the third year the lake has been a part of CAMP (1994 being the first). In addition to the CAMP monitoring, the lake has been monitored in past years by Council staff. As part of the 2005 volunteer monitoring program, Lake Jane was monitored 13 times from mid-April to early-October. Graphs as well as the actual data collected by volunteers show the seasonal variability in TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability). The graphs and data tables are presented on the information sheet on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	13.2	9.0	21.0	A
CLA (µg/l)	3.0	1.6	4.7	A
Secchi (m)	5.3	3.5	7.0	A
TKN (mg/l)	0.64	0.47	0.83	
Overall Grade				A

Data retrieved from the MPCA's STORET water quality database revealed an extensive historical database for Lake Jane. Varying amounts of water quality data were available representing each year since 1980. Out of the 18 years of data, Secchi transparencies were recorded annually, phosphorus was measured in 10 of those 18 years, and CLA eight years. The lake's best water quality has been recorded in 2000 and 2004-2005.

The average user perception rankings of Lake Jane correspond to the overall good quality of the lake. On a 1 to 5 ranking scale, the mean physical condition ranking was 1.0 (1- "crystal clear"), while the mean recreational suitability was 1.0 (1- "beautiful").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of any errors in the lake's data/physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Jane

Lake Elmo, Washington Co.



LAKE ID: 820104

W/D: Valley Branch

Volunteer: Chuck Taylor


● Sampling site

Contours in meters



2005 Data

Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	CLA ug/L	Surf. TP ug/L	Bot. TP ug/L	Secchi M	PC 1 thru 5	RS 1 thru 5
4/16/05	13.1				2.1	15		6.5	1	1
5/2/05	11.2				2.3	12		6.5	1	1
5/15/05	13.5				1.8	11		6.8		
5/19/05	13.7				1.6	15		7	1	1
6/1/05	20.1				1.6	16		7	1	1
6/15/05	23.9				3.1	10		4.7	1	1
7/3/05	24.3				3.3	21		4	1	
7/16/05	32					9		4.3	1	
8/1/05	28.3				3.4	13		4.5		
8/15/05	25.4				4.1	12		3.5	1	
9/4/05	22.9				2.5	12		5.2	1	
9/18/05	24.8				4.7	14		4.8		
10/3/05	20.3				6.9	21		4.7	1	




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	B	B			C		B	B				B	
Chlorophyll a					C		B	B				B	
Secchi Depth	A	A	A	A	B	B	B	B	B	B	B	B	C
Overall					C		B	B				B	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		A						A				A	A
Chlorophyll a		A						A				A	A
Secchi Depth	B	B						A				A	A
Overall		A						A				A	A

Source: Metropolitan Council and STORET data



Jellum's Bay [Site-1] (82-0052-02) *Carnelian - Marine Watershed District*

Jellum's Bay, located in New Scandia Township in Washington County. This was the sixth year the lake has been involved in CAMP. Because the maximum depth of the 72-acre lake is only 4.9 m (16 feet), the majority of the lake's area is considered littoral zone (the 0-15 foot depth area of the lake dominated by aquatic vegetation). Additionally, because of the lake's shallowness it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake's mean depth of 2.4 m (roughly 8 feet) and its surface area translate to an approximate lake volume of 569 ac-ft. The lake has a 333-acre immediate watershed, which translates to a watershed-to-lake area ratio of 4.6:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff).

A search through the STORET nationwide water quality database for data on the lake provided historical water quality data on the lake for years 1996-2004. The resulting data and graphs appear on the next page.

The lake was monitored 14 times between mid-April and mid-October, 2005. Results are presented on graphs and data tables on the following page. During each monitoring event the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	116.7	72.0	180.0	D
CLA (µg/l)	57.1	12.0	150.0	D
Secchi (m)	1.0	0.3	2.1	D
TKN (mg/l)	2.08	1.20	2.80	
Overall Grade				D

The lake's 2005 overall grade of D (calculated from the three individual grades) is identical to those recorded in 1996-1999, and 2001-2004, and better than that of 2000 (F). The lakes individual summer means in 2005 however, were much worse than those recorded in 2004.

Statistical analysis on the lake's water quality database did not detect any long-term trends. In the short-term however, the lake's water quality seems to be well represented by an overall grade of D. In order to detect any possible long-term trends, additional years of data collection are needed.

Throughout the monitoring period, the volunteers' opinions of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The mean perceived physical condition of Jellum's Bay was 3.3 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability was 3.9 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Jellum's Lake, Site 1 New Scandia Twp., Washington Co.


LAKE ID: 820052-02

W/D: Camelian-Marine

Volunteer: Wash. Co. SWCD




● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/13/05	13.1	10.7	5.65	4.86	38	74		1.1	3	4
4/25/05	13	12.8	5.14	4.67	20	88		1.4	2	3
5/10/05	16.7	14.3	5.64	2.23	19	72		1.5	2	4
5/23/05	16.8	15.2	4.91	3.87	26	80		1.2	2	4
6/6/05	22.6	21.1	4.26	0.24	12	142		2.1	2	4
6/22/05	26.9	21.7	9.14	0.35	15	110		1.8	2	2
7/7/05	24.4	22	12.4	0.13	110	109		0.6	4	5
7/18/05	28.2	19	7.42	0.06	64	98		0.8	3	4
8/1/05	29.3	18.3	9.66	0.43	150	105		0.3	4	4
8/15/05	25.6	19.5	6.62	0.39	65	105		0.6	4	4
8/30/05	24.3	20.5	9.88	0.36	62	154		0.8	3	4
9/13/05	23.1	22.6	6.54	0.51	43	128		0.9	5	4
9/29/05	16.9	16.1	6.62	4.33	62	180		0.8	5	4
10/10/05	16	12.8	8.85	4.46	80	118		0.6	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				F	D	D	D	D	D	C	D	D	D
Chlorophyll a				D	D	D	D	F	D	D	F	C	D
Secchi Depth				D	D	F	F	F	D	D	D	D	D
Overall				D	D	D	D	F	D	D	D	D	D

Source: Metropolitan Council and STORET data



Keller Lake [Burnsville] (19-0025) Black Dog Watershed Management Commission

Keller Lake, located in both the cities of Apple Valley and Burnsville (Dakota County), covers an area of 63 acres and has a maximum depth of 3.0 m (10 feet). The lake's mean depth of 1.4 m (4.6 feet) and surface area translates to an approximate lake volume of 290 ac-ft (the lake volume may have changed over the past couple years due to the lake level rising 1.5 to 2.0 feet above normal). Because the maximum depth is only 3.0 m, the entire lake area is considered littoral (the area of aquatic plant dominance), and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The land uses within the 353-acre contributing watershed to the lake are approximately split between agricultural uses and urban/residential. The watershed-to-lake size ratio is 6:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff).

This was the seventh year that Keller Lake has been enrolled in CAMP. The lake had been monitored by Council-staff in the past as part of a study on Crystal Lake (which Keller flows into). In 2005, the lake was monitored 14 times between mid-May and mid-October. The collected data and resulting graphs showing TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	56.3	21.0	81.0	C
CLA (µg/l)	11.6	1.5	34.0	B
Secchi (m)	1.5	0.9	2.0	C
TKN (mg/l)	1.10	0.53	1.70	
Overall Grade				C

The lake's overall grade in 2005 (C) is similar to that recorded in 2002 and 2004, better than those recorded in 1996-1997, 1999-2001 and 2003 (D), and worse than the B recorded in 1998. Because of the variability of the lake's grades, no statistically significant long-term trend is evident from the lake's water quality database. The lake's water quality seems to be best represented by an overall grade of D+/C.

Similar to past years, the lake's Secchi transparency in 2005 would have been greater except on many monitoring events the lake's excessive submergent macrophyte growth got in the way. Therefore, the lake's 2005 water clarity was actually better than that represented by the summer mean and resulting grade.

Throughout the 2005 season, the volunteer monitor ranked their perceptions of the lake's physical and recreational condition on a 1-to-5 scale. The mean perceived physical condition was 3.0 (3- "definite algae present"), while the mean recreational suitability was 4.4 (between 4- "no swimming - boating ok" and 5- "no aesthetics possible").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Keller Lake

Burnsville, Dakota Co.




Lake ID: 190025

WMO: Black Dog

Volunteer: Glen Gramse


● Sampling site

Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/15/05	15					49		1.5	3	4
5/23/05	18.6				8.8	51		1.0	3	4
6/6/05	23.4				6.2	50		2.0	3	4
6/14/05	24.7				1.5	26		2.0	3	4
6/22/05	30.1				2.5	21		2.0	2	5
6/30/05	26				5.9	63		1.8		
7/5/05	23.4				8.7	81		1.5	3	5
7/19/05	29				9.7	56		1.7	3	5
8/3/05	27.3				4.2	50		2.0	3	5
8/15/05	25.4				16	78		1.2	3	4
8/30/05	26				24	62		1.2	3	4
9/14/05	23.5				34	78		0.9	3	4
9/25/05	20.6					67		1.0	5	4
10/10/05	14.5				6.6	58		2.4	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			D	D	C	D	D	D	C	D	C	C	C
Chlorophyll a			F	C	A	C	C	C	B	C	B	B	B
Secchi Depth			D	D	C	D	D	D	D	D	C	C	C
Overall			D	D	B	D	D	D	C	D	C	C	C

Source: Minnesota Consolidated STORET data



Kingsley Lake (19-0030) *Black Dog Watershed Management Commission*

This was the eighth year that Kingsley Lake has been monitored as part of CAMP (1995-1997 and 2000-2005). Additionally, the lake was monitored by Council-staff in 1993. The lake is located in the northwestern corner of the City of Lakeville in Dakota County. The lake has a surface area of 44 acres (shoreline length of 1.7 miles), a maximum depth of 4.0 m (13 feet), and a contributing watershed of 193 acres. The resulting watershed-to-lake size ratio is a rather small 4:1 that no doubt contributes to the good water quality of the lake. Because of the shallowness of the lake, the entire lake is considered littoral (area of aquatic vegetation dominance), and never develops and maintains a thermocline.

Kingsley Lake was monitored 16 times between mid-April and early-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	14.6	8.0	18.0	A
CLA (µg/l)	3.4	1.5	6.9	A
Secchi (m)	2.5	1.6	3.0	B
TKN (mg/l)	0.64	0.33	0.81	
Overall Grade				A

Similar to past years, the Secchi transparency in 2005 would have been greater except that on many monitoring events, the lake's excessive submergent macrophyte growth got in the way. For this reason, if it weren't for the macrophyte interference, the water clarity conditions may have actually been that of an A grade.

The physical and recreational conditions of Kingsley Lake as perceived by the volunteer(s) were ranked on a 1-to-5 scale. The mean physical condition ranking was 2.0 (2- "some algae present"), while the mean recreational suitability ranking was 2.6 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's water quality seems to be represented by a water quality grade of A/high B.


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake ID: 190030
WMO: Black Dog
Volunteers: The Green Family

Kingsley Lake

Lakeville, Dakota Co.

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	14				8.8	91		3	1	1
4/29/05	9				3.5	18		3.1	2	1
5/15/05	10				3.5	16		3	2	2
5/19/05	14				5.5	14		3		
5/27/05	16				4.5	17		3	2	2
6/10/05	24				6.9	18		2.4	2	2
6/22/05	24				1.5	13		2.6	2	3
7/9/05	25				2.5	14		3	2	3
7/24/05	29				2.9	16		1.9	2	3
8/2/05	24				4.2	13		2.8	2	3
8/3/05					1.9					
8/21/05	23				2.4	18		2.4	2	3
9/3/05	20				2.4	13		1.6	2	3
9/15/05	19				5.2	17		2.1	2	2
9/28/05	16				2.7	8		2.1	2	2
10/13/05	12				4.5	10		2.5	1	2

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B		B	A	A			A	A	A	B	A	A
Chlorophyll a	A		A	A	A			A	A	A	A	A	A
Secchi Depth	A		B	B	B			B	C	B	B	B	B
Overall	A		B	A	A			A	B	A	B	A	A

Source: Metropolitan Council and STORET data



Kismet Lake (82-0333) Browns Creek Watershed District

Kismet Lake is located in Washington County. The relatively small lake has a maximum depth of approximately 3.7 m (12 feet). Because of the shallowness of the lake the whole lake is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation.

This was the eighth year that Kismet Lake has been involved in CAMP (in was initially enrolled in 1998). The only available lake data found through a search for historical water quality was the 1998-2005 CAMP data. The lake was monitored 13 times between mid-April and mid-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	34.1	24.0	46.0	C
CLA (µg/l)	19.7	7.7	41.0	B
Secchi (m)	2.0	1.5	2.4	C
TKN (mg/l)	0.87	0.72	1.10	
Overall Grade				C




The lake's 2005 overall grade is identical to those recorded in 1998-2002, and worse than those recorded in 2003-2004 (B).

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's quality seems well represented by an overall grade of B+/C. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

The perceived physical and recreational conditions of the lake, recorded by the volunteers, were ranked on a 1-to-5 scale. The rankings are shown in both tabular and graphical form on the lake's associated information sheet. The mean physical condition ranking was 2.3 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 3.3 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/11/05	13	7.5	6.28	7.85	12	26		2.6	2	3
4/28/05	11.2	10.6	6.3	6.4	13	33		2.1	2	2
5/11/05	16.3	14.2	5.59	6.64	9.5	31		2.4	3	4
5/25/05	17.3	16.9	4.67	4.12	10	32		2.3	3	4
6/8/05	23.5	18.4	4.13	0.21	7.7	32		2.3	2	4
6/23/05	27.7	22	8.13	0.25	8	27		2.1	2	2
7/5/05	24.1	23.2	6.89	3.7	17	46		2.0	2	4
7/18/05	28.4	28	7.51	5.76	11	33		2.3	2	3
8/5/05	26.4	23.7	4.3	0.35	18	38		2.0	2	3
8/19/05	23.5	23	5.72	0.22	39	34		1.5	3	4
8/31/05	24.7	21.6	7.63	0.59	27	40		1.5	2	3
9/15/05	21.5	20.7	4.72	0.26	41	38		1.5	2	3
10/14/05	14.1	13.1	8.83	2.23	28	36		1.8	2	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus						C	C	D	C	C	B	B	C
Chlorophyll a						C	C	C	B	B	B	A	B
Secchi Depth						C	C	C	C	C	B	B	C
Overall						C	C	C	C	C	B	B	C

Source: Metropolitan Council and STORET data



Klawitter Lake (82-0368) Valley Branch Watershed District

Klawitter Lake is a small 4.5-acre lake located within the boundaries of Lake Elmo (Washington County). Because of the shallowness of the lake, it is considered entirely littoral (the 0-15 foot depth zone of a lake dominated by aquatic vegetation), and does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's surface area and watershed size (168 acres) translates to an 37:1 watershed-to-lake size ratio. Generally the larger the ratio, the greater the potential stress on the lake from surface runoff.

This was the fourth year that Klawitter Lake has been involved in CAMP. Other than for the 2002-2005 CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty.

As part of the watershed's involvement in CAMP in 2005, the lake was monitored 14 times between late-April and mid-October. During each sampling event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	103.1	70.0	191.0	D
CLA (µg/l)	21.8	12.0	45.0	C
Secchi (m)	0.8	0.5	1.0	D
TKN (mg/l)	1.99	1.20	3.5	
<i>Overall Grade</i>				D

The 2005 overall grade determined through the calculation of all three parameters was D, is similar to that recorded in 2003-2004, and worse than the overall grade of C of 2002.

The lake's TP mean does include a point deemed an outlier (691.0 µg/l) on one sampling date (September 1). The point is roughly four-to-six times greater than the expected value and a reason for the erroneous number is not known (in-lab processes and QA/QC runs within the batch where the questionable sample was analyzed, were checked and okayed).

Throughout the monitoring period, the volunteers' opinions of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the following page. The mean physical condition ranking was 2.9 (between 2- "some algae present" 3- "definite algae present"), while the mean recreational suitability ranking was 2.9 (between 2- "minor aesthetic problems" and 3- "swimming slightly impaired").

As mentioned earlier, there are no water quality data available for Klawitter Lake other than the 2002-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Klawitter Pond Lake Elmo, Washington Co.




Lake ID: 820368
WD: Valley Branch
Volunteer: Bonnie Jurand

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 thru 5	1 thru 5
4/24/05	14.4				9.1	85		1.0	2	1
5/15/05	13.9				16	103		1.0	2	2
5/23/05	21.6				17	77		1.0	2	2
6/12/05	25.7				19	135		0.9	3	3
6/29/05	25.3				15	96		0.8	3	3
7/16/05	30.2				13	77		0.9	3	4
7/27/05	26.7				29	70		0.8	3	3
8/1/05	25.9				24	77		0.7	3	3
8/13/05	23.5				28	102		0.7	3	3
9/1/05	22.4				12	691		0.5	4	3
9/14/05	22.8				45	191		0.5	3	3
10/1/05	20.5				53	189		0.4	3	3
10/16/05	14.5				43	124		0.6	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus										D	D	D	D
Chlorophyll a										B	C	C	C
Secchi Depth										D	F	D	D
Overall										C	D	D	D

Source: Metropolitan Council and STORET data



La Lake (82-0097) City of Woodbury

La Lake, located in the City of Woodbury (Washington County), has been monitored through CAMP since 1994. The lake has a surface area of approximately 35 acres (1.3 miles around) and a maximum depth of 3.5 m (11 feet). Because of the shallowness of the lake, it is considered entirely littoral (the 0-15 foot depth zone of a lake dominated by aquatic vegetation), and does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

As part of the lake's involvement in CAMP in 2005, the lake was monitored 13 times between late-April and mid-October. During each sampling event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	63.0	53.0	74.0	C
CLA (µg/l)	16.1	3.1	66.0	B
Secchi (m)	1.8	0.9	2.4	C
TKN (mg/l)	1.04	0.70	1.30	
<i>Overall Grade</i>				C


The lake's 2005 overall grade was identical to those recorded in 1994, 1996-1997, and 1999-2003, and worse than those recorded in 1995 and 1998 (B).

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's overall water quality seems to be well represented by a water quality grade of high-C/low-B. With this in mind, however, some concern should be given to the recent (late-1990's and early-2000's) poor water quality years.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.




La Lake

Woodbury, Washington Co.



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/25/05	12.3				10	51		1.5	1	2
5/15/05	11.8				13	60		1.4	1	2
5/23/05	15.7				12			1.2	2	2
6/9/05	23.2				66	68		0.9	3	3
6/24/05	25.9				31	53		1.3	4	4
7/6/05	24.08				18	57		1.4	4	4
7/22/05	25.3				10	71		1.7	4	4
8/5/05	25.5				6.8	67		2.0	3	3
8/19/05	24.1				4.2	55		2.2	2	3
9/1/05	22.4				3.8	59		2.4	2	3
9/13/05	23.5				5.7	66		2.0	2	3
9/27/05	19.6				3.1	74		2.0	2	3
10/14/05	13.4				4.6	61		2.4	2	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		C	C	D	D	C	D	D	D	D	C	NA	C
Chlorophyll a		B	A	B	C	B	C	C	C	B	C	NA	B
Secchi Depth		C	B	C	C	B	C	C	C	C	B	NA	C
Overall		C	B	C	C	B	C	C	C	C	C	NA	C

Source: Metropolitan Council and STORET data



Lac Lavon Lake (19-0446) Black Dog Watershed Management Commission

This was the ninth year that Lac Lavon has been involved in CAMP. The only known water quality data available for the lake were Secchi transparency data in 1989-1991 and CAMP data for 1997-2005.

The lake, located within the City of Apple Valley in Dakota County, is actually an abandoned gravel pit maintained by groundwater (MDNR 1996). The lake is a unique resource in the Twin Cities Metropolitan Area because it is one of only six lakes in the seven-county area stocked with trout (rainbows). The 55-acre lake (2.3 miles in circumference) has a maximum depth of 9.8 m (32 feet) and 65 percent of the lake is considered littoral zone (the 0-15 foot depth zone of the lake dominated by aquatic vegetation). The lake's fishing pier is located on the eastern end of the lake. An area of concern and need for future management is the recent detection of Eurasian Water Milfoil (*Myriophyllum spicatum*) in the lake.

Lac Lavon was monitored 12 times between mid-May and mid-October, 2005. The data and resulting graphs showing seasonal variability in TP and CLA concentrations, Secchi transparency, and user perceptions are presented on the information sheet following these written comments.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	12.6	7.0	17.0	A
CLA (µg/l)	3.0	1.0	9.8	A
Secchi (m)	4.4	3.6	5.5	A
TKN (mg/l)	0.67	0.34	1.20	
Overall Grade				A

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The mean physical condition ranking was 1.0 (1-"crystal clear"), while the mean recreational suitability ranking was 1.0 (1- "beautiful").

No statistically significant long-term trend is evident from the lake's overall water quality database, in the short-term however, the lake's water quality seems well represented by an overall grade of A. A recent MPCA conducted trend analysis on the lake's Secchi transparency data, however, revealed a statistically significant decrease in recent water clarity. The reason for the degradation in the lake's water clarity is not entirely known. A more in-depth study combining watershed as well as in-lake monitoring may help determine the areas contributing the most to the lake's degradation.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you detect any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lac Lavon Apple Valley/Burnsville, Dakota Co.

Lake ID: 190446
WMO: Black Dog
Volunteer: Wally Shaver

● Sampling site
Contours in meters



2005 Data





Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
5/15/05	12				9.8	16		3.6	1	1
5/18/05	14.2				6.2	14		3.8	1	1
5/29/05	19.9				12	13		3.8	1	1
6/10/05	24.1				1.9	17		4.6	1	1
6/26/05	24.1				1.3	10		5.5	1	1
7/10/05	26.8				1	15		5	1	1
7/24/05	28.1				4.9	14		4.4	1	1
8/7/05	26.2				2.4	12		4.5	1	1
8/21/05	24				1.7	8		4.3	1	1
9/5/05	23.2				2.4	7		4.7	1	1
10/2/05	19.1				6.3	11		3.7	1	1
10/16/05	15				7.7	11		2.9	1	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth										A	A	A	
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus					A	A	A	A	B	A	A	A	A
Chlorophyll a					A	A	A	A	A	A	A	A	A
Secchi Depth					A	A	A	A	A	A	A	A	A
Overall					A	A	A	A	A	A	A	A	A

Source: Metropolitan Council and STORET data



Langton Lake [North Basin] (62-0204) Rice Creek Watershed District

Langton Lake is divided into three distinct basins. For this reason there were three monitoring sites in 2005. Additionally, the results will be discussed individually for each of the three sites.

The entire 30-acre lake is located within the City of Roseville (Ramsey County). The maximum and mean depths of the lake are 1.5 and 1.2 m (5 and 4 feet), which along with the surface area, translate to an approximate volume of approximately 120 ac-ft. Because of the shallowness of the lake, its entire surface area is considered littoral, (the shallow [0-15 feet] area dominated by aquatic plants). The lake's contributing watershed is 257 acres, which translates to a watershed-to-lake area ratio of 9:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff).

This marks the first year in which any of the three Langton Lake sites have been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Therefore, 2005 is the only known year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

Langton Lake (North Basin) was monitored 14 times between mid-April and early-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	173.6	100.0	335.0	F
CLA (µg/l)	123.3	29.0	400.0	F
Secchi (m)	0.69	0.3	1.0	F
TKN (mg/l)	1.91	0.90	3.10	
Overall Grade				F

Langton (North Basin) had the worst recorded water quality of the lake three monitored sites.

As mentioned earlier, there are no water quality data available for Langton Lake (North Basin) other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.1 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 4.4 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").



The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Langton Lake, North Basin Roseville, Ramsey Co.


Lake ID: 620204
W/D: Rice Creek
Volunteer: Yul Yost

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	18.9				29	131		0.6	1	2
5/1/05	10.5				38	113		0.8		
5/15/05	13.7				49	100		1	3	2
5/29/05	19.4				53	102		1	3	4
6/12/05	26.5				29	112		1	1	3
6/26/05	28.2				56	166		0.3	5	5
7/10/05	29.3				160	144		0.6	3	5
7/24/05	27.8				400	237		0.5	3	5
8/7/05	24.2				360	249		0.5	4	5
8/21/05	23.6				38	335		0.7	3	5
9/4/05	19.1				61	168		0.6	3	5
9/18/05	25.5				92	183		0.6	3	5
10/2/05	18.8				170	105		0.9	2	5
10/9/05	13.6				86	72		1	1	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth					F	F		D	D		C		
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													F
Chlorophyll a													F
Secchi Depth													F
Overall													F

Source: Metropolitan Council and STORET data



Langton Lake [Site-1] (62-0049-01) Rice Creek Watershed District

Langton Lake (Site-1) was monitored 14 times between mid-April and early-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	53.4	30.0	107.0	C
CLA (µg/l)	22.7	7.3	70.0	C
Secchi (m)	1.1	0.9	1.4	D
TKN (mg/l)	1.28	0.49	1.90	
Overall Grade				C

As mentioned earlier, there are no water quality data available for Langton Lake (Site-1) other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.0 for physical condition (2- "some algae present"), and 3.0 for recreational suitability (3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Langton Lake, Site 1 Roseville, Ramsey Co.

Lake ID: 620049-01

W/D: Rice Creek

Volunteer: Yul Yost





● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/17/05	17.4				8.6	42		1.5	1	1
5/1/05	10.7				9.4	53		1.4		
5/15/05	13				7.3	38		1.2	2	2
5/29/05	19				9.9	34		1.2	2	1
6/12/05	25.6				13	46		1.3	1	1
6/26/05	26.9				28	64		1.2	3	4
7/10/05	28				22	61		1.2	2	4
7/24/05	28.1				70	107		0.9	1	2
8/3/05	26.3				51	75		1.0	2	4
8/21/05	23.9				21	48		1.1	3	5
9/4/05	22				11	31		0.9	2	5
9/18/05	22.8				7.3	30		1.0	2	2
10/2/05	18.1				6.3	22		1.0	1	1
10/9/05	13.7				6.1	25		1.5	1	1



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													C
Chlorophyll a													C
Secchi Depth													D
Overall													C

Source: Metropolitan Council and STORET data

Langton Lake [Site-2] (62-004-02) Rice Creek Watershed District

Langton Lake (Site-2) was monitored 14 times between mid-April and early-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	43.8	30.0	87.0	C
CLA (µg/l)	15.7	8.1	48.0	B
Secchi (m)	1.1	0.6	1.5	D
TKN (mg/l)	0.81	0.37	1.50	
Overall Grade				C

Langton (Site-2) had the best recorded water quality of the lake three monitored sites.

As mentioned earlier, there are no water quality data available for Langton Lake (Site-2) other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. A recent MPCA conducted trend analysis on the lake's Secchi transparency data, revealed a statistically significant improvement in recent water clarity.

To better understand the lake's overall water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.0 for physical condition (2- "some algae present"), and 3.2 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Langton Lake, Site 2 Roseville, Ramsey Co.

Lake ID: 620049-02

W/D: Rice Creek

Volunteer: Yul Yost

Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/17/05	17.1				8.5	37		1.2	1	1
5/1/05	9.8				8.1	37		1.5		
5/15/05	12.8				8.7	36		1.3	2	2
5/29/05	19				12	36		1.2	1	1
6/12/05	27.3				6.8	43		1	1	1
6/26/05	28				16	49		1.1	1	4
7/10/05	29.9				15	38		1.1	2	4
7/24/05	29.4				48	87		1	3	4
8/3/05	27				17	39		1.3	2	4
8/21/05	24.7				19	47		0.8	5	5
9/4/05	22.3				17	36		0.6	2	5
9/18/05	24				5.5	30		1.3	1	2
10/2/05	18.3				5.3	24		1.4	1	1
10/9/05	13.1				7.2	28		1.2	1	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													C
Chlorophyll a													B
Secchi Depth													D
Overall													C

Source: Metropolitan Council and STORET data



Lee Lake (19-0029) Black Dog Watershed Management Commission

Lee Lake, a 25-acre land-locked lake with a maximum depth of 5.2 m (17 feet), is located in Lakeville (Dakota County). The shoreline length of the lake is 1.0 miles. The majority of its 324-acre watershed (which translates to a watershed-to-lake size ratio of 13:1) is now developed with urban uses; however, past cattle farming is the primary phosphorus source to the lake and may have left behind an internal loading problem. To determine if this is the case, a more in-depth monitoring program is needed. An abundance of submerged aquatic vegetation (Curlyleaf pondweed) has been a continuing problem in the lake. Not only is it an aesthetic and recreational problem, but the decaying of plants in late-summer adds to concentrations of phosphorus in the water column.

The lake has been involved in CAMP in 1994-1997 and 2000-2005. In an attempt to inhibit algal populations within the lake, barley straw has been added since 2003. Barley straw has been used for algal control in the United Kingdom for many years. A recent study on Valley Lake-Lakeville, Minnesota (discussed later in Valley Lake section of this report), has suggested that carbon from the decaying barley straw inhibits algal populations via microbial competition for phosphorus (McComas and Anhorn 2004). Therefore, in an attempt to determine if the straw method successfully reduced algal biomass in 2005, Lee Lake TP, TKN, CLA and Secchi transparencies were tested nine times between late-April and late-August. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	70.1	30.0	165.0	D
CLA (µg/l)	17.0	4.2	44.0	B
Secchi (m)	1.5	0.9	2.8	C
TKN (mg/l)	1.33	0.65	2.50	
Overall Grade				C

While the lake's overall water quality grade is identical to those recorded in 1994-1997, 1999, and 2001-2004, and better than that recorded in 2000 (D), the CLA and Secchi parameter means indicate that the lake's 2005 water quality was its best since 2001. No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake seems well represented by an overall grade of C. In order to determine any long-term trends or to better define the lake's normal water quality range, more data are needed.

After a recent fish survey on the lake suggested that the unusually high fish densities may be impacting the barley treatment on the lake, roughly 80 pounds of fish per acre were recently removed. The survey had revealed that bluegill sunfish, black crappies, and black bullheads dominated the lake's fishery. In fact, the number of bluegills caught per net was high, with the average haul of bluegills per net averaging 465 per net. The local average range for bluegills per net is 3-25 bluegills (McComas 2004). The recent fish removal coincides with the improved water quality suggesting not only that the barley straw treatment was successful at improving the lake's 2005 water quality, but the unbalanced fishery may have been negating the benefits of previous barley straw treatments.

Throughout the course of the study, the volunteer monitors ranked their perceptions of the lake's physical and recreational condition on a 1-to-5 scale. These rankings, as well as the data and graphs discussed above, are shown on the lake's information sheet on the following page. The mean physical condition ranking was 2.4 (between 2- "3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability ranking was 3.4 (3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.





If you know of any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lee Lake

Lakeville, Dakota Co.


Lake ID: 190029
WMO: Black Dog
Volunteer: Jamie Cooper

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/26/05	11.2				11	43		1.6	2	3
5/10/05	15.7				42	47		2.8	2	2
5/23/05	18.5				59	30		1.7	2	2
6/10/05	27				44	62		1.2	2	4
6/21/05	29.7				83	45		0.9	3	4
7/19/05	28.6				73	44		1.9	2	3
7/27/05	27				26	105		0.9	3	4
8/8/05	26.9				36	63		0.9	3	4
8/21/05	23				44	165		1.1	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		C	C	C	C			D	C	C	C	C	D
Chlorophyll a		C	B	B	B			C	B	B	C	C	B
Secchi Depth		C	C	C	C			D	C	C	C	D	C
Overall		C	C	C	C			D	C	C	C	C	C

Source: Metropolitan Council and STORET data



Legion Pond (82-0462) Valley Branch Watershed District

Legion Pond is a small 16-acre lake located within Lake Elmo (Washington County). The lake has a 224-acre immediate drainage area, which results in a watershed-to-lake area ratio of 14:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This marks the first year in which Legion Pond has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Therefore, 2004 is the only known year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	105.3	10.0	193.0	D
CLA (µg/l)	39.5	8.0	130.0	C
Secchi (m)	0.9	0.5	1.2	D
TKN (mg/l)	1.25	0.60	1.80	
Overall Grade				D

When comparing the lakes TP (nutrient), CLA (algal biomass estimator), and Secchi (water clarity) grades, it is apparent that the TP and Secchi grades (and summer means) are worse than the CLA grade. In a most cases, the three should be fairly comparable. One possible explanation for the lake's 2005 findings may be that the majority of the lake's TP comes from either in-lake suspended sediments (re-suspension), or the intrusion of sediment-laden runoff to the lake, which in turn lessens the clarity of the water and inhibits algal growth.

As mentioned earlier, there are no water quality data available for Legion Pond other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.1 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 1.0 for recreational suitability (1- "beautiful").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Legion Pond Lake Elmo, Washington Co.

Lake ID: 820462
W/D: Valley Branch
Volunteer: Molly Winkels

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/17/05	17.9				13	51		1.2	1	1
5/1/05	8.7				9	51		1.2	1	
5/15/05	13.6				12	58		1.2	1	
5/18/05	15.2				16	52		1.0		
5/30/05	21.3				29	125		1.1	2	1
6/17/05	29.8				21	88		0.8	2	
6/25/05	30				25	193		0.6	2	
7/10/05	29.7				67	182		0.8	2	
7/24/05	30.5				130	140		0.5	2	
8/7/05	27.4				57	144		0.6	3	
8/21/05	27.8				8	10		1.0	3	
9/22/05	23				37	114		0.8	3	1
10/5/05	18.5				65	174		0.8	2	
10/16/05	14.6				41	114		0.7	2	1

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													D
Chlorophyll a													C
Secchi Depth													D
Overall													D

Source: Metropolitan Council and STORET data



Lily Lake (82-0023) City of Stillwater

Lily Lake, located in the City of Stillwater in Washington County, was monitored seven times between mid-April and early-October, 2005. The lake has been monitored through CAMP since 1995.

The 52-acre lake has a maximum depth of 17.4 m (57 feet), and has public access located on the lake's northern shore and a fishing pier on its southern shore. On each sampling date Lily Lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	42.6	31.0	53.0	C
CLA (µg/l)	19.0	5.7	36.0	B
Secchi (m)	2.0	1.2	3.8	C
TKN (mg/l)	0.99	0.54	1.30	
<i>Overall Grade</i>				C

The lake's 2005 overall water quality grade is similar to those recorded in 1996-2000 and 2002-2004, and worse than those of 1995 and 2001 (B).

The physical and recreational conditions of Lily Lake as perceived by the volunteer(s) were ranked on a 1-to-5 scale. These rankings are also graphed on the lake's information sheet. The mean physical condition ranking was 2.8 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.8 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

A search for water quality data through Council, MPCA, and STORET files resulted in a moderate amount of data. While 1995-2005 are the only years for which nutrient data are available, Secchi transparencies were collected through the MPCA's Citizen Lake Monitoring Program in 1985, and 1987-1992. The data seem to show a wide fluctuation in the lake's mean CLA concentration and water clarity. The best conditions were recorded in 1995 and 2001 (A's and B's), while 1996-2000 and 2002-2005 conditions were mainly represented by C's.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lily Lake Stillwater, Washington Co.




Lake ID: 820023
WMO: Middle St. Croix River
Volunteer: Wash. Co. SWCD

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	10/15	10/15
4/19/05	15.9	3.7	6.41	0.15	11	66		2.0	3	3
5/19/05	14	3.8	5.72	0.23	15	51		1.2	2	3
6/16/05	24.6	3.8	7.54	0.6	7.5	36		3.8	2	2
7/11/05	23.8	3.9	9.23	0.06	5.7	31		2.1	3	3
8/6/05	27.6	4	9.17	3.87	36	53		1.2	4	3
9/12/05	23.4	4.1	7.43	1.83	31	43		1.4	3	3
10/3/05	19.6	4.2	8.54	0.98	23	28		1.5	3	3




Lake Water Quality Grades Based on Summer Time Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth						D		C	C	C	C	C	B
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			C	C	C	C	C	C	C	C	C	C	C
Chlorophyll a			B	C	B	C	C	C	A	B	B	B	B
Secchi Depth			A	B	C	C	C	C	B	C	C	C	C
Overall			B	C	C	C	C	C	B	C	C	C	C

Source: Metropolitan Council and STORET data



Little Carnelian Lake (82-0014) *Carnelian - Marine Watershed District*

This was the sixth year of CAMP monitoring in Little Carnelian Lake which is located in Stillwater Township (Washington County). The lake was first enrolled in the program in 2000. The 162-acre lake (which has a shoreline length of 1.7 miles), has a mean and maximum depth of 10.7 m (35 feet) and 21.3 m (70 feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 5,686 ac-ft. The lake does not have a public access and its 565-acre watershed translates to a meager 3.5:1 watershed-to-lake size ratio (the greater the ratio, the greater the potential stress on the lake from surface runoff).

The lake was monitored 14 times between mid-April and mid-October, 2005. Results are presented on graphs and data tables on the following page. During each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	13.6	9.0	25.0	A
CLA (µg/l)	2.8	2.0	4.0	A
Secchi (m)	6.9	5.8	7.6	A
TKN (mg/l)	0.54	0.42	0.80	
<i>Overall Grade</i>				A

Similar to all past years of CAMP monitoring, the individual grades result in overall lake grade of A for Little Carnelian Lake. This places the lake's water quality within the top 10 percent of Metro Area lakes for the years 2000-2005. In fact, similar to that reported in 2004, the lake's 2005 Secchi mean was again the best mean water clarity in CAMP.

The collected data and resulting graphs showing TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake's information sheet on the following page.

Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 1.8 (between 1- "crystal clear" and 2- "some algae present"), while the mean recreational suitability ranking was 1.1 (between 1- "beautiful" and 2- "minor aesthetic problem").

A search of the STORET nationwide water quality database for data on the lake revealed a moderate database throughout the 1990's with nutrient data available in 1991-1996 and 1998-2005. The lake's database indicates that the lake's water quality is well represented by an overall grade of A. Furthermore, a recent MPCA conducted trend analysis on the lake's Secchi transparency data, revealed a statistically significant improvement in recent water clarity.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Little Carnelian Lake

Stillwater Twp., Washington Co.

LAKE ID: 820014




WCD: Carnelian-Marine

Volunteer: Wash. Co.

SWCD


◆ Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/13/05	7.6	4.2	6.75	1.76	2.6	13		6.1	2	1
4/25/05	11.5	4.6	6.78	0.08	4.9	12		6.9	1	1
5/10/05	13.8	4.8	6.71	0.21	3.6	18		7.6	2	1
5/20/05	12.6	4.9	6.14	0.44	3	13		7.6	1	1
6/6/05	20.1	5.2	5.1	0.39	2.6	25		7.6	2	1
6/21/05	24.1	5.4	8.24	0.71	2.7	10		6.4	2	1
7/6/05	23.8	5.6	7.9	0.69	2.1	15		6.7	2	1
7/19/05	26.9	5.7	7.22	0.05	2.5	12		5.8	2	1
8/1/05	26	5.7	6.34	0.73	2	9		7.6	2	1
8/15/05	24.7	5.8	8.55	3.31	2	12		7.0	2	2
8/29/05	23.3	5.8	7.85	3.44	4	11		6.7	2	1
9/13/05	23.1	6.1	8.11	1.07	3.4	12		7.2	1	1
9/29/05	19	6.1	7.76	1.25	3.3	12		6.1	2	1
10/10/05	16	6.2	8	0.89	4.1	6		6.1	2	1




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus												A	A
Chlorophyll a												A	A
Secchi Depth												A	A
Overall												A	A

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				A	A			A	B	A	A	A	A
Chlorophyll a				A	A			A	A	A	A	A	A
Secchi Depth	A	A	A	A	A	A		A	A	A	A	A	A
Overall				A	A			A	A	A	A	A	A

Source: Metropolitan Council and STORET data



Little Johanna Lake (62-0058) Rice Creek Watershed District

This was the fifth year that Little Johanna Lake, which is located on the boundary between the Cities of Arden Hills and Roseville (Ramsey County), was monitored as part of CAMP. The 35-acre lake has a maximum depth of 12.0 m (39 feet). A search through the STORET nationwide water quality database for data on the lake came up empty other than for the 2001-2005 CAMP data.

The lake was monitored six times from mid-April to early-September, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	65.6	50.0	82.0	C
CLA (µg/l)	24.0	11.0	33.0	C
Secchi (m)	1.8	1.1	2.6	C
TKN (mg/l)	1.36	0.91	1.70	
Overall Grade				C

The lake's 2005 overall grade is identical to those of 2001-2004.

Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 3.3 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability ranking was 4.0 (4- "no swimming – boating ok").

As mentioned earlier, there are no water quality data available for Little Johanna Lake other than the recent 2001-2005 data. Therefore it is not possible to determine any long-term trends. In the short-term, however, the lake seems well represented by an overall grade of C. To better understand the lake's water quality and where it may be heading, more data are needed.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.



If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Little Johanna Lake

Arden Hills/Roseville,
Ramsey Co.





Lake ID: 620058
W/D: Rice Creek
Volunteer: Jason Johnson

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	15				54	70		0.8	1	4
6/10/05	25				11	50		2.5	2	4
6/23/05						55				
6/24/05	28				23	62		2.4	4	4
7/22/05	27				33	79		1.2	5	4
9/2/05	22				29	82		1.1	2	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus									C	D	D	C	C
Chlorophyll a									C	C	C	B	C
Secchi Depth									C	C	C	C	C
Overall									C	C	C	C	C

Source: Metropolitan Council and STORMET data

Long Lake [Apple Valley] (19-0022) *Vermillion River Watershed Management Commission*

Long Lake, which has a surface area of roughly 36 acres, is located within the City of Apple Valley (Dakota County). Other than the fact that the maximum depth of the lake is approximately 3.5 m (10 feet), there is no known morphological data available for the lake. Because the lake is relatively shallow, it does not develop and maintain a thermocline (a density gradient owed to changing water temperatures throughout the water column), and the entire lake is considered littoral, (the shallow [0-15 feet] area dominated by aquatic plants).

This is the fifth year in which Long Lake was involved in CAMP (1997 and 2002-2004 being the others). A search for historical water quality data for the lake came up empty.

As part of the 2005 volunteer monitoring program, Long Lake was monitored 15 times from early-May to mid-October. Graphs as well as the actual data collected by the volunteer(s) show the seasonal variability in TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability). The graphs and data tables are presented on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	234.9	74.0	366.0	F
CLA (µg/l)	81.3	27.0	240.0	F
Secchi (m)	0.4	0.2	0.6	F
TKN (mg/l)	3.57	1.50	8.90	
Overall Grade				F

The lake's 2005 overall grade is identical to those recorded in 2002-2004, and worse than that recorded in 1997 (D)

Because 2005 is only the fifth year of available data, no long-term trends can be determined. In the short-term however, the lake's overall water quality is well represented by an overall grade of F. To better understand the quality of the lake and what direction it may be heading, more years of data collection are needed.





Throughout the course of the study, the volunteer monitors ranked their perceptions of the lake's physical and recreational condition on a 1-to-5 scale. These user perception rankings are shown on the lake's information sheet on the following page. The mean physical condition ranking was 2.5 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability was 4.0 (4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Long Lake

Apple Valley, Dakota Co.

Lake ID: 190022
WMO: Dakota County
Volunteers: Cherie Serie and
Al Kettelkamp



2005 Data

Date	Surf Temp C	Bot Temp C	Surf DO mg/L	Bot DO mg/L	Chl a ug/L	Surf TP ug/L	Bot TP ug/L	Secchi M	PC 1-5	RS 1-5
4/14/05	15				66	113		0.6	2	2
5/1/05	9				45			0.5	2	4
5/6/05	19				90	128		0.5	2	4
5/11/05	20				48	136		0.6		
5/24/05	22				44	361.5		0.6	2	4
6/10/05	26.8				27	74		0.5	3	4
6/22/05	30.5				43	106		0.4	2	4
7/24/05	30.7				120	274		0.3	3	4
8/2/05	28.2				240	232		0.2	3	4
8/16/05	25.8					355		0.2	2	4
8/29/05	26.9					319		0.2	4	4
10/1/05	18.6				81	314		0.2	4	4
10/12/05	13.9				110	176		0.2	2	4
10/16/05					410					
10/29/05					390					

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus					D					F	F	F	F
Chlorophyll a					D					F	F	F	F
Secchi Depth					F					F	F	F	F
Overall					D					F	F	F	F

Source: Metropolitan Council and STO RET data

Long Lake [Mahtomedi] (82-0130) Rice Creek Watershed District

Long Lake, a 48-acre lake with a maximum depth of 7.7 m (25 feet), is located within City of Mahtomedi (Washington County). Roughly 92 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This marks the third year in which Long Lake has been involved in CAMP. Other than for the 2003-2005 CAMP data, a search through the STORET nationwide water quality database for historic data on the lake was unsuccessful.

On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The lake was monitored 12 times between early-May and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	23.2	18.0	57.0	B
CLA (µg/l)	4.3	2.7	5.8	A
Secchi (m)	2.6	2.2	3.2	B
TKN (mg/l)	0.64	0.42	0.95	
<i>Overall Grade</i>				B

The lake's 2005 overall water quality grade is slightly worse than that recorded in 2004.

As mentioned earlier, there are no water quality data available for Long Lake other than the 2004-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.6 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 2.6 for recreational suitability (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us



Long Lake [May Township] (82-0030) *Marine on St. Croix WMO*

Long Lake is an 88-acre lake located in May Township (Washington County). There is little morphological data available for the lake. Because the maximum depth is only 3.7 m (12 feet), the entire lake area is considered littoral (the area of aquatic plant dominance), and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake, which was monitored through CAMP in 1993-1997 and 1999-2004, was sampled seven times between late-April and early-October, 2005.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	43.8	29.0	69.0	C
CLA (µg/l)	14.2	4.7	42.0	B
Secchi (m)	2.6	1.5	3.1	B
TKN (mg/l)	0.82	0.56	1.00	
Overall Grade				B




Statistical analysis on the lake's overall database fails to reveal any "statistically significant" long-term trends. The lake's 2005 overall grade was identical to those recorded in 2000-2001 and 2003-2004, and better than those of 1993-1997, 1999, and 2002 (C). Overall, the lake's water quality is representative of a high-C/B grade.

A recent MPCA conducted trend analysis on the lake's Secchi transparency data, however, revealed a statistically significant improvement in recent water clarity.

Throughout the course of the study, the volunteer monitor ranked their perceptions of the lake's physical and recreational condition on a 1-to-5 scale. These rankings as well as the data and graphs discussed above are shown on the lake's information sheet on the following page. The mean physical condition ranking was 2.2 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 3.0 (between 3- "swimming slightly impaired").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/20/05	15.7	11.1	4.92	50.23	5.8	94		2.7	2	2
5/19/05	13.8	12.7	5.14	2.33	8.8	44		3.0	2	3
6/15/05	23.5	18.7	6.64	3.05	4.7	29		2.9	2	2
7/13/05	21.8	21.8	11.22	0.37	6.1	29		2.9	2	3
8/11/05	24.8	22	5.45	0.35	42	69		1.5	3	4
9/8/05	22.9	20.4	8.82	0.51	9.6	48		2.4	2	3
10/6/05	16.1	16	6.45	0.67	6.6	32		3.0	2	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C	C	C	C	C		C	C	C	C	C	B	C
Chlorophyll a	C	C	C	B	C		B	B	B	B	A	A	B
Secchi Depth	B	C	C	C	C		C	B	B	C	B	B	B
Overall	C	C	C	C	C		C	B	B	C	B	B	B

Source : Metropolitan Council and STORM data



Long Lake [Pine Springs] (82-0118) Valley Branch Watershed District

Long Lake is a 62-acre lake located in Pine Springs Township (Washington County). The mean and maximum depth of the lake is 3.6 m (roughly 12 feet) and 10.4 m (34 feet), respectively. Roughly 55 percent of the lake's area is considered littoral (the 0-15 foot depth area of aquatic vegetation dominance). The lake's size and mean depth results in an approximate lake volume of 744 ac-ft.

The lake's surface area and watershed size (2,060 acres) translates to a 33:1 watershed-to-lake size ratio. Generally the larger the ratio, the greater the potential stress on the lake from surface runoff.

Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has recently been found in the lake.

This was the third year that Long Lake has been involved in CAMP (the other being 1993 and 2004). The lake has been monitored in the past by Council staff (most recently in 2003). The lake was monitored 14 times between mid-April and mid-October, 2005. The volunteer data and resulting graphs showing the seasonal variability in TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	39.1	18.0	98.0	C
CLA (µg/l)	19.0	6.3	45.0	B
Secchi (m)	1.8	0.9	4.0	C
TKN (mg/l)	1.25	0.62	2.00	
Overall Grade				C

A search for water quality data on Long Lake uncovered a very small database. The only year other than 2005 where water quality data was available was 1984, 1993, and 2003-2004. While the limited database restricts the ability to determine long-term trends, the lake seems to fluctuate between an overall grade of B and C. The lake's best recorded water quality was observed in 2003. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

The average user perception rankings, on a 1-to-5 scale, was 2.2 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 2.5 for recreational suitability (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Long Lake

Pine Springs, Washington Co.



Lake ID: 820118

W/D: Valley Branch

Volunteer: Bill Feely





● Sampling station

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/13/05	9.1				29	66		0.9	1	2
4/27/05	10.4				32	52		0.9	1	2
5/10/05	14.2				19	38		1.2	1	2
5/23/05	14.4				12	96		1.7	1	2
6/6/05	20.8				7	28		2.4	2	2
6/21/05	23.8				6.3	18		4.0	3	3
7/5/05	22.4				10	25		2.0	3	3
7/19/05	24				11	24		2.6	2	3
8/1/05	24.9				17	34		1.6	3	3
8/16/05	25.5				33	36		0.9	3	3
8/30/05	22.7				16	46		1.7	2	2
9/14/05	21.6				33	36		1.1	2	2
9/27/05	18.5				45	47		0.9	2	2
10/15/05	15.4				35	46		1.0	2	2



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					C								
Chlorophyll a					B								
Secchi Depth					C								
Overall					C								

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B										B	C	C
Chlorophyll a	B										A	B	B
Secchi Depth	C										B	C	C
Overall	B										B	C	C

Source: Metropolitan Council and STORM data

Long Lake [Stillwater] (82-0021) Browns Creek Watershed District

Long Lake, which has a surface area of roughly 96 acres, is located on the western boundary of the City of Stillwater (Washington County). Its maximum depth is 6.7 m (22 feet).

As part of the 2005 volunteer monitoring program, Long Lake was monitored 14 times from mid-April to mid-October. This was the ninth year that Long Lake has been involved in CAMP. The lake was also involved in the program in 1995-1996, and 1998-2004. Graphs as well as the actual data collected by volunteers show the seasonal variability in TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability). The graphs and data tables are presented on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	72.7	44.0	121.0	D
CLA (µg/l)	55.1	17.0	130.0	D
Secchi (m)	0.9	0.3	1.7	D
TKN (mg/l)	1.55	0.77	2.30	
Overall Grade				D

A search for water quality data through Council, MPCA, and STORET files resulted in a moderate amount of data. Nutrient data are available for the lake in 1995-1996, and 1998-2005. Additionally, Secchi transparencies collected through the MPCA's Citizen Lake Monitoring Program are available for 1987, 1989, and 1991-1994. When these data are analyzed, it reveals that the lake's water clarity, prior to the C recorded in 2004, had been fairly constant with grades of F in 1987, 1991-1995, 1998-2003, and D in 1989, 1996 (although the 1996 database is limited), and 2005.

A recent MPCA conducted trend analysis on the lake's Secchi transparency data, revealed a statistically significant decrease in recent water clarity

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The mean physical condition ranking was 3.5 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability was 3.7 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Long Lake, Stillwater, Washington Co.





Lake ID: 820021
WMD: Browns Creek
Volunteer: Wash. Co. SWCD

• Sampling site
Contours in meters



2005 Data

Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	Chl. a µg/L	Surf. TP µg/L	Bot. TP µg/L	Secchi M	PC 1-5	RS 1-5
4/11/05	13	4.6	7.02	0.64	18	61		1.5	2	3
4/27/05	11.8	5.1	5.8	0.06	22	58		1.2	2	4
5/11/05	16.3	5.8	5.57	0.14	22	73		1.4	3	3
5/26/05	17.6	6.2	4.68	0.04	17	44		1.7	3	3
6/7/05	25.4		4.66		19	47		1.2	3	4
6/23/05	26.1	7.5	10.03	0.53	46	59		0.8	3	3
7/6/05	22.5	8.1	10.96	0.5	40	80		0.8	4	4
7/18/05	26.5	8.4	7.58	0.09	43	66		0.8	3	4
8/3/05	27.5	9.1	9.45	0.51	97	121		0.5	4	4
8/17/05	25.4	9.6	11.86	0.4	130	76		0.3	4	4
8/31/05	23.8	10.5	9.29	0.58	66	82		0.6	4	4
9/15/05	21.8	10.8	6.78	0.32	72	79		0.6	4	4
10/3/05	19.5	11.8	8.64	0.48	49	66		0.6	4	4
10/13/05	14.7	11.4	10.71	0.43	45	48		0.9	3	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth								F		D		F	F
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			D	D		D	D	F	D	D	D	C	D
Chlorophyll a			D	D		F	F	F	F	D	D	C	D
Secchi Depth	F	F	F	D		F	F	F	F	F	F	C	D
Overall			D	D		F	F	F	F	D	D	C	D

Source: Metropolitan Council and STO RET data

Long Lake [Washington Co.] (82-0068) *Carnelian - Marine Watershed District*

Long Lake is a 35-acre lake located within New Scandia Township (Washington County). The maximum and mean depths of the lake are 2.1 m (roughly seven feet) and 1.1 m (three-and-a-half feet), respectively. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake's surface area and mean depth translates to an approximate volume of 126 ac-ft.

The majority of the land within the 381-acre watershed is undeveloped. The watershed-to-lake size ratio is 11:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). There is no formal boat access point on the lake.

This was the sixth year that Long Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake was very limited. The only years in which data are available other than the 2000-2005 CAMP data, were 1998-1999. The lake was monitored seven times between mid-April and early-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	89.8	58.0	118.0	D
CLA (µg/l)	42.2	16.0	110.0	C
Secchi (m)	1.0	0.5	1.4	D
TKN (mg/l)	1.44	1.20	2.00	
Overall Grade				D

The lake's 2005 overall grade, which is identical to that recorded in 2001 and 2004, is better than those recorded in 1998-2000, and 2003(F), and worse than the C observed in 2002.

As mentioned earlier, there is a limited amount of water quality data available for Long Lake. Therefore it is not possible to determine any long-term or short-term trends. The lake's quality has fluctuated between an overall grade of C and F. To better understand the lake's water quality and where it may be heading, more data are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.8 for physical condition (between 2- "some algae presnt" and 3- "definite algae present"), and 4.0 for recreational suitability (2- "no swimming – boating ok").


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Long Lake New Scandia Twp., Washington Co.

LAKE ID: 820068




WD: Carnelian-Marine

Volunteer: Wash. Co. SWCD



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/29/05	16.4	15.1	4.81	2.09	17	115		1.7	2	3
5/16/05	14.8	12.2	5.86	5.25	21	71		1.1	3	4
6/14/05	24.2	22	7.39	0.34	16	108		1.4	2	4
7/11/05			8.59	9.27	18	58		1.4	3	4
8/9/05	27	26.8	7.53	0.43	110	118		0.5	3	4
9/7/05	22.7	22.7	11.12	1.43	46	94		0.8	3	4
10/3/05			7.89	5.96	25	75		0.9	5	5




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus						D	D	D	C	C	D	D	D
Chlorophyll a						F	F	F	D	C	F	D	C
Secchi Depth						F	F	F	D	D	F	D	D
Overall						F	F	F	D	C	F	D	D

Source: Metropolitan Council and STORET data



Loon Lake (82-0015-02) *Carnelian - Marine Watershed District*

This was the sixth year of CAMP monitoring in Loon Lake, which is located in the Stillwater Township (Washington County). The 64-acre lake has a mean and maximum depth of 2.4 m (eight feet) and 4.9 m (16 feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 206 ac-ft. Because of the shallowness of the lake, the majority of its area is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation), and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake does not have a public access and its 407-acre watershed translates to a 6.4:1 watershed-to-lake size ratio (the greater the ratio, the greater the potential stress on the lake from surface runoff).

The lake was monitored seven times between mid-April and early-October, 2005. Results are presented on graphs and data tables on the following page. During each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	142.6	103.0	192 .0	D
CLA (µg/l)	96.4	41.0	240.0	F
Secchi (m)	0.6	0.3	0.9	F
TKN (mg/l)	2.68	2.00	3.30	
Overall Grade				F

The lake's 2005 overall grade was identical to those recorded in 1996-1998 and 2003-2004, and worse than those in 2000-2002 (D).

The collected data and resulting graphs showing TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake's information sheet on the following page.

Throughout the summer, the volunteer ranked the lake's physical and recreational conditions on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 3.2 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability ranking was 4.0 (4- "no swimming – boating ok").

Because of the limitedness of the lake's water quality database, no long-trend can be determined. In the short-term however, the lake's water quality seems to be well represented by D/F+. To better understand the quality of the lake and what direction it may be heading, more years of data collection are needed.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Loon Lake

Stillwater Twp., Washington Co.


LAKE ID: 8200 15-02

W/D: Camelian-Marine

Volunteer: Wash. Co. SWCD





◆ Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/20/05	16.5	11	5.93	0.15	53	93		1.2	2	3
5/17/05	13.1	13.1	6.43	6.76	41	137		0.9	2	3
6/15/05	23.8	14.7	7.06	0.33	46	103		0.8	3	4
7/14/05	28	23.6	9.93	0.11	240	171		0.3	4	5
8/9/05	25.1	23.3	6.2	0.32	86	192		0.5	3	4
9/6/05	24.5	21.3	10.07	0.45	69	110		0.6	4	4
10/4/05	19.3	17.1	6.56	0.81	49	98		0.8	3	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				F	F	F	F	D	D	D	D	D	D
Chlorophyll a				D	D	D	D	D	D	D	F	F	F
Secchi Depth				F	F	F	F	D	D	F	F	F	F
Overall				F	F	F	F	D	D	D	F	F	F

Source: Metropolitan Council and STORET data

Lotus Lake (10-0006) City of Chanhassen

While Lotus Lake has previously been monitored by Council staff (1985, 1990 and 1999-2000) and the MPCA's volunteer Secchi program (1980, 1988-1991), 2005 marks the third year the lake has been monitored through CAMP. Lotus Lake, with a surface area of 246 acres, is located within the City of Chanhassen (Carver County) [public access to the lake is possible on the southern end of the lake]. The lake's surface area and its 1,033-acre watershed translates to a 4:1 watershed-to-lake size ratio (the greater the ratio, the greater the potential stress on the lake from surface runoff).

The lake's maximum and mean depths of 8.9 and 4.3 (29.2 and 14.2 feet), along with its surface area, translates to a lake volume of approximately 3,500 ac-ft. Roughly 74 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake is considered a "Priority Lake" due to its multi-recreational uses. Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has been reported on the lake.

In 2005, Lotus Lake was monitored eight times between late-May and early-October. Results are presented on graphs and data tables on the following page. During each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	47.0	22.0	85.0	C
CLA (µg/l)	30.5	7.2	66.0	C
Secchi (m)	1.5	0.7	2.5	C
TKN (mg/l)	1.47	0.78	2.50	
Overall Grade				C

The lake's 2005 overall grade of C is identical to those recorded in 1985, 1999-2000, and 2004, and worse than the D recorded in the minimal monitoring of 2003 (D).

Throughout the summer, the volunteer ranked their opinion of the lake's physical and recreational conditions on a 1-to-5 scale (see lake information sheet). The mean physical condition was 2.6 (between 2- "some algae present" and 3- "definite algae present"), while the recreational suitability ranking was 2.2 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lotus Lake **Chanhassen, Carver Co.**


Lake ID: 100006

WD: Riley Purgatory-Bluff Creek

Volunteer: Shelly Strohmaier




● Sampling site

Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/25/05	16.5				8.2	22		2.4		
5/30/05	18.6				7.2	24		2.5	2	2
7/9/05	25				8.3	31		1.8	2	2
7/25/05	25				21	77		1.5		
7/26/05	26.3				29	49		1.35	2	2
8/21/05	25.8				66	41		0.8	4	3
9/8/05	23				42	85		0.7	3	2
10/2/05	20.2				49	52		1	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus						C							
Chlorophyll a						C					C		
Secchi Depth	D					C			D	C	C	C	
Overall						C							

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							C	C			D	C	C
Chlorophyll a							C	C			C	C	C
Secchi Depth							C	C			D	C	C
Overall							C	C			D	C	C

Source: Metropolitan Council and STORET data



Louise Lake (82-0025) *Carnelian - Marine Watershed District*

Louise Lake is a 48-acre lake located within Stillwater Township (Washington County). The maximum and mean depths of the lake are 3.7 m (roughly 12 feet) and 1.8 m (six feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 283 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's 616-acre watershed and surface area translates to a watershed-to-lake size ratio of 13:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). There is no formal boat access point on the lake.

This was the sixth year that Louise Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake provided limited information (1996-2004).

The lake's Secchi transparency was monitored seven times from mid-April to early-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.


Water samples to be analyzed for TP, TKN and chlorophyll were not collected for the lake in 2005. Because Secchi transparency was the only data collected there are no nutrient or chlorophyll concentration means to compare to previous years. The lake's 2005 summertime (May through September) mean Secchi transparency was 0.7 m (minimum of 0.3 m and a maximum of 1.5 m). This translates to a grade of D for water clarity. The lake's 2005 water clarity was dramatically worse than those recorded in 2003-2004 (2.0m and 2.5 m), and similar to that of 2001 (0.9 m).

Because of the limitedness of the lake's water quality database, no long-term can be determined. In the short-term however, the data seems to show that the lake, consistently fluctuates between an overall C and D grade. To better understand the lake's water quality and where it may be heading, more data are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.8 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.2 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").




If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Louise **Stillwater Twp., Washington Co.**



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 to 5	1 to 5
4/21/05	15.7	13.7	6.72	4.71				2.3	3	4
5/17/05	12.7	11.7	6.98	3.57				1.5	4	4
6/16/05	23.3	16.7	11	0.35				0.5	3	4
7/14/05	29.1	23.5	11.04	0.4				0.3	5	5
8/9/05	27.4	27.3	7.33	0.32				0.5	3	4
9/6/05	24.3	24.3	10.65	1.95				0.8	4	4
10/4/05	19.6	17.9	7.09	3.96				0.8	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				D	D	B	C	D	D	D			
Chlorophyll a				D	D	D	F	B	D	C			
Secchi Depth				B	C	C	C	C	D	D	B	C	D
Overall				C	D	C	D	C	D	D			

Source: Metropolitan Council and STORET data



MacDonald's Pond (82-0062) *Carnelian – Marine Watershed District*

MacDonald's Pond is an approximate 12-acre land-locked lake located within New Scandia Township (Washington County). The maximum depth of the lake is 2.7 m (roughly 9 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). There is very little other known morphological data available for the waterbody.

This was the second year that MacDonald's Pond has been involved in CAMP (2004 being the other). On each of the sampling days the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The lake was monitored 14 times between mid-April and mid-October, 2005.

The following are the averages of the three events for each of the parameters tested.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	26.5	20.0	34.0	B
CLA (µg/l)	4.4	2.8	7.1	A
Secchi (m)	2.8	2.4	3.2	B
TKN (mg/l)	0.82	0.60	1.00	
Overall Grade				B

The lake's 2005 overall grade is worse than the A recorded in 2004.

Other than for the 2004-2005 CAMP data, there are no known water quality data available for MacDonald's Pond. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.




The perceived physical and recreational conditions (ranked on a 1-to-5 scale) are shown on the lake's information sheet on the next page. The average user perception rankings, were 2.4 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 3.6 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

MacDonald's Pond


New Scandia Twp., Washington Co.

LAKE ID: 820062
WCD: Camelian-Marine
Volunteer: Wash. Co.
SWCD



2005 Data

Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	CLA ug/L	Surf. TP ug/L	Bot. TP ug/L	Secchi M	PC 1 to 5	RS 1 to 5
4/13/05	13	12.1	6.08	6.67	2.3	30		3.0	3	4
4/27/05	11.4	11.3	6.48	5.55	2.1	26		3.2	3	4
5/10/05	17.4	12.4	5.7	5.29	2.8	24		3.2	2	4
5/23/05	17.7	15.6	5.26	6.54	3.8	20		2.7	3	4
6/6/05	23.7	23.7	4.46	3.34	2.8	28		3.2	2	4
6/22/05	27.8	27.8	8.22	0.41	4.5	29		3.2	2	2
7/7/05	24.5	24.5	7.63	7.92	3.7	29		2.9	2	4
7/18/05	29.7	28.3	7.85	4.34	5.4	34		2.7	3	4
8/1/05	29.2	25.2	7.18	2	5	27		2.7	2	4
8/15/05	24.3	23.6	6.72	3.53	7.1	25		2.4	3	4
8/30/05	23.8	22.7	7.83	1.8	5	31		2.9	2	3
9/13/05	23.2	23.1	7.54	5.71	4.6	24		2.4	3	4
9/28/05	18	18	13.01	13.28	3.6	20		2.4	2	3
10/10/05	15	13	7.38	6.89	3.9	17		3.0	2	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												A	B
Chlorophyll a												A	A
Secchi Depth												A	B
Overall												A	B

Source: Metropolitan Council and STORET data



Maple Marsh (82-0038) *Carnelian - Marine Watershed District*

Maple Marsh Lake is a 38-acre lake located within May Township (Washington County). The maximum and mean depths of the lake are 3.4 m (roughly 11 feet) and 1.7 m (five-and-a-half feet), respectively. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake's surface area and mean depth translates to an approximate volume of 126 ac-ft.

The majority of the land within the 148-acre watershed is undeveloped. The watershed-to-lake size ratio is 4:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff).

This was the fifth year that Maple Marsh Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake provided limited information (1997-2005).

The lake's Secchi transparency was monitored seven times from mid-April to early-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

Water samples to be analyzed for TP, TKN and chlorophyll were not collected for the lake in 2005. Because Secchi transparency was the only data collected there are no nutrient or chlorophyll concentration means to compare to previous years. The lake's 2005 summertime (May through September) mean Secchi transparency was 0.8 m (minimum of 0.6 m and a maximum of 1.2 m). This translates to a grade of D for water clarity. The lake's 2005 Secchi grade is identical to those recorded in 1997 and 1999-2001, better than the F in 1998, but worse than the C's of 2002 and 2004.

Because of the limited nature of the lake's water quality database the determination of any statistically significant long-term trend is not possible. With this in mind, the lake's water quality data seems to show a consistent fluctuation between an overall grade of C and D. To better understand the lake's overall water quality and where it may be heading, more monitoring is suggested.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.2 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.5 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Maple Marsh May Twp., Washington Co.

LAKE ID: 820038
WD: Carnelian-Marine
Volunteer: Wash. Co. SWCD




◆ Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/21/05	15.1	14.9	5.09	4.68				1.5	1	4
5/19/05	11.7	13.1	6.23	0.79				1.2	3	4
6/14/05	24	23.9	8.7	2.5					2	4
7/13/05	21.4	21.4	8.07	0.34				0.6	3	5
8/12/05	23.9	23.9	8.18	7.54				0.6	4	5
9/9/05	21.9	21.5	8.86	8.43				0.8	4	5
10/6/05	15	14.9	7.67	0.5				0.8	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus					F	F	C	D	F				
Chlorophyll a					D	F	F	C	D				
Secchi Depth					D	F	D	D	D	C	D	C	D
Overall					D	F	D	D	D				

Source : Metropolitan Council and STORET data



Maria Lake (10-0058) Carver County Environmental Services

Maria Lake is a 169-acre lake located within Benton and Dahlgren Townships (Carver County). The maximum depth of the lake is 1.0 m (roughly three feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This was the fourth year that Maria Lake has been involved in CAMP. Other than the 1999 CAMP data (no grades determined because it was only monitored twice) and 2000-2001 CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty.

The lake was monitored 13 times from late-April to mid-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	158.9	37.0	368.0	F
CLA (µg/l)	83.5	3.4	240.0	F
Secchi (m)	0.7	0.1	1.3	D
TKN (mg/l)	2.59	0.94	5.7	
Overall Grade				F

The lake's Secchi transparency in 2005 would have been greater except for the shallowness of the lake. On numerous monitoring events, The Secchi disk was clearly noticeable while resting on the lake's bottom. Therefore, the lake's 2005 water clarity was actually better than that represented by the summer mean.

As mentioned earlier, there is no water quality data available for Maria Lake other than the 1999-2001 and 2005 CAMP data. Therefore it is impossible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, more data are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.3 for physical condition (between 2- "some algae present" and 3- "definite algal presence"), and 3.8 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming - boating ok").



The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Maria


Benton Twp. and Dahlgren Twp.,
Carver Co.

Lake ID: 100058
WMD: Carver County
Volunteer: Carver Co. Env. Services



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl a	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1-5	1-5
4/2/05	15.6		10.95		10			1.2	3	4
5/1/05	17		11.63		6.1	39		1.2	2	4
5/2/05	16.2				3.4	39		1.2	1	3
6/6/05	23.1		11.38		19	44		1.3	1	4
6/16/05	24.7		10.79		7.5	118		1.1	2	4
7/5/05	22.5		15.5		12	89		1	2	4
7/20/05	26.9		6.77		35	142		0.6	2	3
8/3/05	26.7		12.61		210	241		0.3	3	4
8/15/05	23.4		13.7		240	368		0.2	4	4
8/30/05	24.3		12.58		190	241		0.2	3	4
9/15/05	20		7.48		110	239		0.4	3	4
9/26/05	17.2		15.97		86	194		0.5	3	4
10/1/05	13		12.72		47	127		0.7	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								F	F				F
Chlorophyll a								F	C				F
Secchi Depth								F	D				D
Overall								F	D				F

Source: Metropolitan Council and STORET data



Marion Lake (19-0026) City of Lakeville

This was the eighth year that Marion Lake has been a part of CAMP (the others were 1994 and 1999-2004). The area around Lake Marion, located in the City of Lakeville (Dakota County), is rapidly developing. The lake covers an area of roughly 560 acres and has a maximum depth of 6.4 m (21 feet). There are two public accesses to the lake. One access is located in the park while the other is located on the western side of the lake off of 195th Street West. Lake Marion is considered a "Priority Lake" by the Metropolitan Council because of its multi-recreational uses. One problem that may possibly hinder future recreational activity on the lake, however, is Eurasian Water Milfoil (*Myriophyllum spicatum*), which has been reported in the lake.

The lake gets heavy use by area fishermen and other lake users during the winter and summer months. The MDNR manages the lake for northern pike-panfish, and has stocked the lake with walleye over the past decade. Because of past winterkills, the lake's oxygen levels are monitored throughout the winter, and the lake is aerated when needed.

As part of the 2005 volunteer monitoring program, Lake Marion was monitored 16 times from mid-April to mid-October. During each monitoring event the lake was monitored for TP, CLA, TKN, and Secchi transparency, and the lake's perceived physical condition and recreational suitability. Graphs as well as the actual data collected by volunteers, show the seasonal variability of the collected data (see lake information sheet on the next page).

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	33.0	12.0	104.0	C
CLA (µg/l)	23.5	3.0	75.0	C
Secchi (m)	2.0	0.8	4.0	C
TKN (mg/l)	1.15	0.45	1.90	
Overall Grade				C




The resulting overall grade in 2005 (C) (similar to those recorded in 2002-2004), represents a decrease in water quality as compared to the overall grade of B the lake received in 1994, and 1999-2001.

The physical and recreational conditions of the lake, as observed by the volunteer monitors, were ranked on a 1 to 5 ranking scale. The volunteer's user perception rankings are shown on the lake's information sheet. The mean physical condition ranking was 2.1 (between 2- "some algae present" and 3- "definite algae present"), while the lake's mean recreational suitability ranking was 1.3 (between 1- "beautiful" and 2- "minor aesthetic problem").

While Lake Marion does have 15 years of data (13 of which contain some nutrient measurements) over the past 25 years, it is difficult to determine what is happening with the lake's water quality. The available data shows a wide range in the lake's quality with the water quality showing an improvement in the 1990's as compared to the 1980's. The lake received an overall water quality grade of D in 1981; C in 1980, 1983, 1987, and 2002-2005; and finally received a B in 1994, and 1999-2001.


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

Date	Surf Temp C	Bot Temp C	Surf DO mg/L	Bot DO mg/L	CLA ug/L	Surf TP ug/L	Bot TP ug/L	Secchi M	PC 1 2 3 4 5	RS 1 2 3 4 5
4/14/05	8				3.7	48		3.8	1	1
4/24/05	10				6.2	36		2.3	1	1
5/10/05	12				4.4	22		2.5	1	1
5/16/05	17				5.2	12		2.7		
5/24/05					4.5	17			2	1
6/7/05	19				3	18		4.0	2	1
6/21/05	23				3.1	14		3.1	2	1
7/6/05	22				4.9	21		2.0	2	1
7/19/05	24				9.2	23		2.5	2	1
8/2/05	24				21	30		1.1	2	2
8/16/05	23.2				30	27		1.3		
8/30/05	21				36	36		1.2	3	1
9/12/05	22				75	104		0.8	3	2
10/1/05	20				64	71		1.4	2	2
10/9/05	15				68	63		1.3	2	1
10/9/05	12				18	34		2.3	2	1




Lake Water Quality Grades Based on Summertime Averages

Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1990	1991	1992
Total Phosphorus	C	C		C				C		C			
Chlorophyll a	C	D		C				C		C			
Secchi Depth	C	D		B				C		C	C	C	
Overall	C	D		C				C		C			

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		B					B	B	B	C	B	C	C
Chlorophyll a		A					B	A	B	B	C	C	C
Secchi Depth		B					C	B	B	C	C	C	C
Overall		B					B	B	B	C	C	C	C

Source: Metropolitan Council and STORET data



Markgrafs Lake (82-0089) City of Woodbury

Markgrafs Lake, located within the City of Woodbury (Washington County), has a surface area of approximately 46 acres (2.6 miles around), and a maximum depth of 2.4 m (8 feet). The lake, which is used by the MDNR Fisheries as a rearing pond for walleyes, has a piped outlet on the southern end. Downstream from the outlet is a valve that can direct the overflow to either Powers or Wilmes lakes.

The 413-acre drainage area to the lake is presently made up of open/undeveloped areas. Future land uses are projected to be 11.5 percent single-family residential, 14.8 percent multi-family residential, 51.8 percent commercial/retail, 15.1 percent parks/open space, and 6.8 percent ponds/wetlands. The lake's watershed-to-lake size ratio is 10:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). Because of the lake's shallowness, much of the lake is considered littoral zone (the 0-15 foot depth area of the lake dominated by aquatic vegetation). It does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake has been involved in CAMP since 1994. Between mid-April and mid-October, 2005, the lake was monitored 14 times. During each monitoring event; TP, CLA, TKN, and Secchi transparency were measured, as was the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	136.5	88.0	174.0	D
CLA (µg/l)	46.4	8.5	76.0	C
Secchi (m)	0.5	0.3	0.6	F
TKN (mg/l)	2.20	1.40	2.90	
Overall Grade				D

The lake's 2005 overall lake quality report card grade is identical to those recorded in 1994, 1997, 1999, and 2001-2004, better than the F of 1998, and worse than the C's observed in 1995-1996.

A moderate amount historical water quality data is available for Markgrafs Lake. Data found were collected through CAMP in 1994-2005. While no statistically significant long-term trend is evident from the lake's entire water quality database (including TP, CLA and Secchi data), a recent MPCA conducted trend analysis using just the lake's Secchi transparency data, revealed a statistically significant decrease in recent water clarity. The lake's overall quality generally fluctuates between a low C (1995-1996, and 2000) and a D (1994, 1997, 1999, and 2001-2005). The lake experienced its worst recorded overall water quality (F) in 1998 and its best water quality in 1995.


Throughout the course of the monitoring season the volunteer monitor ranked the lake's perceived physical and recreational conditions on a 1-to-5 scale. The mean physical condition was 4.0 (between 4- "high algal color") while the mean recreational suitability was 4.0 (4- "no swimming – boating ok").

If you know of any errors in the lake's data/physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Markgrafs Lake Woodbury, Washington Co.

Lake ID : 820089
WD: South Washington
Volunteer: Terry Riley

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/10/05	15.1				26	115		0.6	4	4
4/24/05	15				22	122		0.6	4	4
5/16/05	15.1				52	168		0.6	4	4
5/28/05	16.2				76	118		0.6	4	4
6/11/05	21.8				8.5	92		0.5	4	4
6/22/05	23				18	88		0.5	4	4
7/10/05	27.4				52	119		0.5	4	4
7/24/05	26.9				19	174		0.4	4	4
8/6/05	26.3				72	160		0.3	4	4
8/20/05	26.9				56	152		0.4	4	4
9/3/05	26.9				66	153		0.4	4	4
9/12/05	25.8				62	161		0.4	4	4
9/27/05	17.4				29	116		0.4	4	4
10/16/05	15.1				21	114		0.4	4	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	D	C	D	D	F	D	D	F	F	D	D	D	D
Chlorophyll a	C	B	B	C	F	C	C	C	C	C	D	C	C
Secchi Depth	D	C	C	D	F	D	C	D	F	D	F	F	F
Overall	D	C	C	D	F	D	C	D	D	D	D	D	D

Source : Metropolitan Council and STORM data



McDonald Lake (82-0010) Valley Branch Watershed District

McDonald Lake is a 54-acre land-locked (no outlet) lake located within Baytown Township (Washington County). The mean and maximum depth of the lake is 1.8 m (nearly 6 feet) and 3.7 m (roughly 12 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake's size and mean depth results in an approximate lake volume of 324 ac-ft.

The lake's surface area and watershed size (1,051 acres) translates to a 12:1 watershed-to-lake size ratio. Generally the larger the ratio, the greater the potential stress on the lake from surface runoff.

This was the sixth year in which McDonald Lake has been involved in CAMP (the lake was enrolled in the program in 1999 and 2001-2004 as well). The only historical water quality data found for McDonald Lake were Secchi transparency data for 1998 and 2000, and CAMP data from 1999 and 2001-2004. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	39.3	25.0	65.0	C
CLA (µg/l)	14.2	2.7	37.0	B
Secchi (m)	1.8	0.7	3.0	C
TKN (mg/l)	1.00	0.64	1.70	
Overall Grade				C

The lake's 2005 overall grade of C is identical to those recorded in 1999, and 2001-2003, and worse than the B recorded in 2004. The lake's 2004 overall grade is the best recorded to date.

Similar to past years, the Secchi transparency in 2005 would have been greater except for the shallowness of the lake. On numerous monitoring events, The Secchi disk was clearly noticeable while resting on the lake's bottom. Therefore, the lake's 2005 water clarity was actually better than that represented by the summer mean and resulting grade.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's quality seems well represented by an overall grade of C. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.



The perceived physical and recreational conditions (ranked on a 1-to-5 scale) are shown on the lake's information sheet on the next page. The average user perception rankings, were 2.4 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 2.5 for recreational suitability (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

McDonald Lake Baytown Twp., Washington Co.

Lake ID: 820010
W/D: Valley Branch
Volunteer: Steve Groves

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	15.5				9.5	50		2.3	1	1
5/4/05	12.4				6.3	65		2.1	2	1
5/16/05	12.9				5.4	41		2.3	1	1
5/18/05	18				5.5	28		2.3		
5/30/05	21.6				2.7	25		2.3	1	1
6/12/05	24.8				8	30		3	2	1
6/26/05	27.2				18	29		1.8	3	2
7/11/05	29.7				37	52		1	3	4
7/27/05	26.7				16	38		1.8	3	4
8/7/05	28.4				20	33		1.5	2	4
8/24/05	24.1				18	46		0.8	4	4
9/11/05	25.7				22	45		0.7	4	4
10/3/05	20.7				6.3	49		1.6	4	4
10/16/05	14.6				18	148		2	3	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							C		C	C	C	C	C
Chlorophyll a							B		C	C	C	B	B
Secchi Depth						C	C	C	C	C	C	B	C
Overall							C		C	C	C	B	C

Source: Metropolitan Council and STORM data



McKusick Lake (82-0020) *Middle St. Croix Watershed Management Organization*

Lake McKusick, a 46-acre lake located within the City of Stillwater (Washington County) has a maximum depth of 4.7 m (roughly 15.5 feet). The lake has been involved in CAMP since 1994. In 2005, the lake was monitored 14 times between mid-April and mid-October.

On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	54.4	26.0	166.0	C
CLA (µg/l)	18.5	3.8	81.0	B
Secchi (m)	2.0	1.5	2.9	C
TKN (mg/l)	0.90	0.55	1.50	
Overall Grade				C




The lake's 2005 overall grade of C is identical to those recorded in 1997, and 2002-2003, better than the D's of 1994-1996 and 1998-1999, but worse than the B's of 2000-2001 and 2004. The overall grade of B recorded in 2000 and 2001 is the lake's best-recorded overall grade to date. A closer look at the three years that the lake received an overall grade of B, reveals that the best parameter means were recorded in 2004.

Throughout the monitoring period, the volunteer(s) ranked their opinions of physical and recreational conditions of the lake on a 1-to-5 scale. The resulting user perception rankings are shown on the lake information sheet. The mean physical condition ranking was 3.4 (3- "definite algae present"), while the mean recreational suitability ranking was 4.0 (4- "no swimming – boating ok").

Because of the wide variation in the lake's 1994-2005 overall water quality database, no long-term trends can be determined. In the short-term however, it seems that the lake was well represented by an overall grade of D/C until recently (2000-2004) when the lake's overall grade has improved to C+/B. In order to detect any possible long-term water quality trends, additional years of data collection are needed.


A recent MPCA conducted trend analysis on the lake's Secchi transparency data, however, revealed a statistically significant improvement in recent water clarity.

If you know of any errors in the lake's data/physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/1/05	12.9	5.2	6.56	0.4	15	62		1.5	2	2
4/27/05	10.4	6.7	5.56	0.16	11	53		2.0	2	3
5/1/05	9.2	8.1	6.1	1.38	9.4	46		2.1	3	4
5/26/05	16.8	16.8	5.53	0.07	4.3	30		2.9	3	5
6/7/05	17.7	9	4.77	0.3	4.2	34		2.1	5	5
6/23/05	26.7	9.3	7.04	0.61	33	66		1.5	5	4
7/8/05		10.8	8.3	0.08	81	166		1.7	2	5
7/25/05	26.3	12.7	8.25	0.68	12	35		2.0	2	3
8/5/05	26.3	18.2	4.28	0.31	9.6	37		1.5	3	4
8/19/05	22.7	10.4	6.7	0.31	5.7	26		2.0	2	4
8/31/05	22.7	13.7	10.23	0.5	3.8	40		2.4	2	3
9/15/05	20.8	18.3	4.56	0.28	22	64		1.5	3	3
10/3/05	20	16.1	7.98	4.74	22	63		1.5	2	3
10/14/05	14.2	12.8	8.86	0.84	21	46		0.9	2	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		D	D	D	C	D	D	C	C	C	C	C	C
Chlorophyll a		D	C	C	C	D	D	B	B	C	B	A	B
Secchi Depth		D	D	D	C	D	D	B	B	D	C	B	C
Overall		D	D	D	C	D	D	B	B	C	C	B	C

Source: Metropolitan Council and STORET data



Meadow Lake (27-0057) Shingle Creek Watershed Management Commission

This was the fourth year of CAMP monitoring on Meadow Lake, which is located in the City of New Hope (Hennepin County). The lake, also monitored as part of CAMP in 1996, 1999 and 2002, was monitored 10 times between early-May and mid-October, 2005.

The 11-acre lake has a 440-acre immediate watershed, which translates to a 40:1 watershed-to-lake size ratio (generally the larger the ratio, the greater the potential stress on the lake from surface runoff). Because of the shallowness of the lake (maximum depth of 1.2 m [4 feet]), the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	248.9	101.0	502.0	F
CLA (µg/l)	68.8	1.4	190.0	D
Secchi (m)	0.5	0.2	0.9	F
TKN (mg/l)	3.70	1.50	6.70	
<i>Overall Grade</i>				F

The lake's overall 2005 lake quality grade of F is identical to those recorded in 1996, 1999, and 2002.




Because 2005 is only the fourth year of available data, no long-trends can be determined. In the short-term however, the lake is well represented by an overall grade of F. To better understand the quality of the lake and what direction it may be heading, more years of data collection are needed.

Throughout the monitoring period, the volunteers ranked the perceived physical condition of the lake on a 1-to-5 scale. The mean perceived physical condition of Meadow Lake was 2.8 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability was 4.1 (between 4- "no swimming - boating ok" and 5- "no aesthetics possible").

If you know of any errors in the lake's data/physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us


Meadow Lake New Hope, Hennepin Co.

Lake ID: 270057
WMO: Shingle Creek
Volunteer: Diane Stauner



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
5/1/05	10.1				5.1	168		0.3	1	4
5/30/05	20.3				1.4	282		0.9	3	4
6/20/05	25				28	101		0.5	2	4
7/9/05	28.6				110	162		0.8	1	
7/24/05	29.7				30	212		0.6	4	4
8/5/05	29.7				100	255		0.3	5	5
8/13/05	23.5				86	502		0.2	4	4
9/11/05	29.7				190	309		0.3	2	4
10/2/05	20.2				96	260		0.4	1	4
10/16/05	14.6				69	205		0.4	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				F			F			F			F
Chlorophyll a				F			F			F			D
Secchi Depth				F			F			F			F
Overall				F			F			F			F

Source: Metropolitan Council and STORET data



Mergen's Lake (82-0482) Valley Branch Watershed District

Mergen's Lake is a 12-acre land-locked lake located within West Lakeland Township (Washington County). The maximum depth of the lake is 1.3 m (roughly 4 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake's surface area and watershed size (1,383 acres) translates to a 115:1 watershed-to-lake size ratio. Generally the larger the ratio, the greater the potential stress on the lake from surface runoff.

This was the fourth year that Mergen's Lake has been involved in CAMP. On each of the sampling days the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. In 2005, the lake was monitored six times between mid-April and mid-July. Because of the lack of August and September monitoring events, the lake's 2005 means, and grades are skewed to the early-summer months.

The lake's 2005 raw data and resulting graphs are presented on the associated lake information page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	142.8	108.0	195.0	D
CLA (µg/l)	10.1	3.8	15.0	B
Secchi (m)	1.0	0.4	1.3	D
TKN (mg/l)	0.98	0.75	1.30	
<i>Overall Grade</i>				C

When comparing the lakes TP (nutrient), CLA (algal biomass estimator), and Secchi (water clarity) grades, it is apparent that the TP and Secchi grades (and summer means) are worse than the CLA grade. In a most cases, the three should be fairly comparable. One possible explanation for the lake's 2005 findings may be that the majority of the lake's TP comes from either in-lake suspended sediments (re-suspension), or the intrusion of sediment-laden runoff to the lake, which in turn lessens the clarity of the water and inhibits algal growth.

Because of the limited nature of the Mergen's Lake water quality database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteers ranked the perceived physical condition of the lake on a 1-to-5 scale. The mean perceived physical condition of Mergen's Lake was 2.8 (between 2- "some algae present" and 3- "definite algae present").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Mergens' Pond

West Lakeland Twp.,
Washington Co.


Lake ID: 820482

WID: Valley Branch

Volunteer: Chris Moosbrugger




● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/18/05	20.4				8.7	150		1.2	1	
5/3/05	10.9				15	108		1.2	2	
5/31/05	22.3				6.6	109		1.3	1	
6/21/05	30				3.8	121		0.4	3	
7/14/05	28				17	195		0.8		
7/19/05	29.8				15	181		1	3	




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								D			NA	NA	D
Chlorophyll a								C			NA	NA	B
Secchi Depth								D			NA	NA	D
Overall								D			NA	NA	C

Source: Metropolitan Council and STORM data



Miller Lake (10-0029) Carver County Environmental Services

Miller Lake, a 145-acre lake located within Dahlgren Township (Carver County) is considered a Metropolitan Area “Priority Lake” because of its multi-recreational uses. The mean and maximum depths of the lake are 3.1 m (10 feet) and 4.3 m (roughly 14 feet), respectively. The lake’s mean depth and surface area translate to an approximate lake volume of 1,479 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake’s water column).

The lake has a 16,701-acre immediate watershed, which translates to a watershed-to-lake area ratio of 115:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff). A 1999 water quality report on water resources in Carver County estimates land use for the watershed at: four percent residential, 71 percent agricultural, two percent commercial/industrial, and 23 percent open/undeveloped (Carver County Planning 1999).

This was the tenth year that Miller Lake has been involved in CAMP. A search through the STORET nationwide water quality database revealed a limited water quality database with water quality data available for 1995-1997, and 1999-2004.

The lake was monitored 14 times between mid-April and mid-October, 2005. Results are presented on graphs and data tables on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	133.2	74.0	268.0	D
CLA (µg/l)	53.8	7.70	140.0	D
Secchi (m)	0.7	0.4	1.60	D
TKN (mg/l)	2.01	0.91	3.30	
Overall Grade				D

The lake’s 2005 overall grade is identical to those recorded in 1997, and 1999-2002, and better than the F’s recorded in 1995-1996 and 2003-2004.

No statistically significant long-term trend is evident from the lake’s water quality database, in the short-term however, it seems that the lake is well represented by an overall D/F grade. Also, the lake’s CLA grade had steadily improved from F’s in 1995-1996, D’s in 1997 and 1999, to C’s in 2000-2002 before falling back to a D in 2003-2005.

Throughout the monitoring period, the volunteer(s) ranked the perceived physical condition of the lake on a 1-to-5 scale. The mean perceived physical condition of Miller Lake was 2.9 (between 2- “some algae present” and 3- “definite algae present”), while the mean recreational suitability was 2.7 (between 2- “minor arsthetic problem” and 3- “swimming slightly impaired”).

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.



If you notice any errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Miller Lake

Dahlgren Twp., Carver Co.




Lake ID: 100029
WMO: Carver County
Volunteer: Carver Co.
Env. Services

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	12		13.1		74	147		0.7		
4/28/05	9.6		13.7		77	184		0.7	2	2
5/10/05	15.4		12.6		56	119		0.6	2	2
5/23/05	17.9		8.8		49	113		0.9	2	2
6/7/05	24		11.07		17	74		0.9	2	2
6/22/05	25.2		6.55		7.7	85		1.6	3	3
7/5/05	25.1				67	121		0.7	3	2
7/21/05	26.9		12.57		74	154		0.5	3	3
8/3/05	27.7				42	144		0.5	3	3
8/17/05	24.8		10.03		28	115		0.8	3	3
8/29/05					54	84		0.6	3	3
9/15/05	21.3		8.95		140	188		0.4	3	3
9/26/05	19		5.73		57	268		0.5	5	4
10/11/05	12.2		7.75		21	328		0.7	2	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			F	F	F		F	F	F	F	F	F	D
Chlorophyll a			F	F	D		D	C	C	C	D	D	D
Secchi Depth			F	F	D		D	D	C	C	F	F	D
Overall			F	F	D		D	D	D	D	F	F	D

Source: Metropolitan Council and STORET data



Mitchell Lake (27-0070) City of Eden Prairie

While Mitchell Lake has previously been monitored by Council staff, 2005 marks the second year the lake has been monitored through CAMP (2004 being the first). Mitchell Lake, with a surface area of 112 acres, is located with the City of Eden Prairie (Hennepin County). The maximum and depths of the lake are 5.8 (19 feet), respectively. Because of the shallowness of the lake, roughly 97 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

Because of its multi-recreational uses, the lake is considered a "Priority Lake". The lake has a public access and fishing pier on its southern end. One problem that may possibly hinder future recreational activity on the lake, however, is Eurasian Water Milfoil (*Myriophyllum spicatum*), which has been reported in the lake.

In 2005, Mitchell Lake was monitored 14 times between late-April and early-October. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP (µg/l)	72.3	28.0	152.0	D
CLA (µg/l)	32.0	6.5	63.0	C
Secchi (m)	1.3	0.4	3.9	C
TKN (mg/l)	1.65	0.79	2.50	
Overall Grade				C

The lake's 2005 overall grade of C is identical to those recorded in 1991, 1995 and 2004, and better than the D's recorded in 1999-2000 and 2003.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's quality seems well represented by an overall grade of C/D+. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

The average user perception rankings, on a 1-to-5 scale, were 1.8 for physical condition (between 1- "crystal clear" and 2- "some algae present"), and 1.1 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.




Lake Mitchell

Eden Prairie, Hennepin Co.

Lake ID: 270070
WD: Riley-Purgatory Bluff
Creek
Volunteer: Gordon Warner


● Sampling site

Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/24/05	13.8				7	40		2.1	1	1
5/8/05	15.9				11	44		2.0	2	1
5/22/05	15.5				6.5	28		2.2	1	1
5/25/05	16				8	33		2.4		
6/7/05	22.1				7.9	31		3.9	2	2
6/19/05	25.5				6.6	65		1.2	4	1
7/3/05	24.4				30	152		0.7	1	1
7/19/05	27.4				59	102		0.6	2	1
8/11/05	24.9				49	84		0.5	2	1
8/21/05	23				38	80		0.6	2	1
8/31/05	22.5				63	84		0.4	2	1
9/10/05	22.7				49	90		0.7	1	1
9/23/05	17					74		1.2	1	1
10/6/05	14.7				8	105		1.2	3	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus												D	
Chlorophyll a												C	
Secchi Depth												C	
Overall												C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	D			D	C	D
Chlorophyll a			C				D	D			D	C	C
Secchi Depth			C				D	C			C	C	C
Overall			C				D	D			D	C	C

Source: Metropolitan Council and STORET data



Moody Lake (13-0023) *Comfort Lake-Forest Lake Watershed District*

Moody Lake is a 35-acre lake located near Chisago City (Chisago County). The lake has a maximum and mean depth of approximately 14.6 m (48 feet). Roughly 63 percent of the lake's surface area is considered littoral zone (the 0-15 foot depth area of aquatic plant dominance).

This marks the first year in which Moody Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Therefore, 2005 is the only known year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	151.1	40.0	284.0	D
CLA (µg/l)	51.3	16.0	150.0	D
Secchi (m)	0.9	0.3	2.0	D
TKN (mg/l)	1.92	1.00	3.10	
Overall Grade				D

As mentioned earlier, there are no water quality data available for Moody other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.8 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.1 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Moody Lake

Chisago Lake Twp., Chisago Co.



LAKE ID: 130023

WMO: Comfort Lake-Forest Lake

Volunteer: Wash. Co. SWCD


● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/18/05	14.2	4.8	8.6	0.25	43	96		1.2	3	2
4/26/05	11.7	4.9	6.23	0.06	24	80		1.5	3	4
5/16/05	11.4	5.6	8.07	1.95	16	78		2.0	2	3
5/24/05	17.2	5.8	5.76	0.03	17	88		1.8	3	4
6/9/05	23.4	6.5	4.9	0.36	19	40		1.2	4	4
6/22/05	25.8	6.9	15.9	0.74	150	270		0.3	5	5
7/7/05	24.5	7.2	17.25	0.65	140	284		0.3	5	5
7/22/05	27.4	7.3	8.86	0.64	39	233		0.5	4	4
8/2/05	29	7.6	6.68	0.51	30	189		0.6	3	4
8/16/05	26.4	7.6	9.43	0.43	46	145		0.8	4	4
8/30/05	23	7.7	8.14	0.65	29	100		0.9	4	4
9/14/05	21.5	7.8	7.09	0.44	27	84		1.1	4	4
10/3/05	18.3	8	7.83	0.94	41	114		0.6	3	4
10/11/05	14	7.7	8.07	0.36	27	167		1.1	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													0
Chlorophyll a													0
Secchi Depth													0
Overall													0

Source: Metropolitan Council and STORET data



Mud Lake (82-0026) *Carnelian - Marine Watershed District*

Mud Lake is a 62-acre lake located within May Township (Washington County). The maximum and mean depths of the lake are 2.1 m (roughly seven feet) and 1.1 m (three-and-a-half feet), respectively. The lake's size and mean depth results in an approximate lake volume of 224 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's small 93-acre immediate watershed translates to a small watershed-to-lake size ratio of 2:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This was the sixth year that Mud Lake has been involved in CAMP (2000-2004 being the others). A search through the STORET nationwide water quality database for historical data on Mud Lake provided data for eight years (1995-2003).

The lake's Secchi transparency was monitored seven times from mid-April to early-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

Water samples to be analyzed for TP, TKN and chlorophyll were not collected for the lake in 2005. Because Secchi transparency was the only data collected there are no nutrient or chlorophyll concentration means to compare to previous years. The lake's 2005 summertime (May through September) mean Secchi transparency was 0.9 m (minimum of 0.6 m and a maximum of 1.2 m). This translates to a grade of D for water clarity.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's water quality seems to be well represented by a overall grade of F. To better understand the lake's water quality and where it may be heading, more data are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.4 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.8 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Mud Lake

May Twp., Washington Co.


LAKE ID: 820026

W/D: Camelian-Marine

Volunteer: Wash. Co. SWCD

◆ Sampling site

Contours in meters






Bathymetry
Unknown

2.1

Island

2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/21/05	15.6	15.6	3.47	3.51				1.5	2	4
5/19/05	14	13.9	4.17	3.93				1.2	2	4
6/14/05	24.5		9.17	0.3				0.8	3	4
7/13/05	31.7		14.84					0.8	4	5
8/12/05	24	23.6	5.03	3.15				0.6	4	5
9/9/05	21.7	21.6	7.75	7.56				1.1	4	5
10/6/05	14.2	14.2	8.6	8.2				0.3	5	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				D	F	F	F	F	D				
Chlorophyll a				D	D	F	D	F	F				
Secchi Depth			F	F	F	F	F	F	F	D	D	C	D
Overall				D	F	F	F	F	F				

Source: Metropolitan Council and STORET data



North Twin Lake (82-0018) *Carnelian - Marine Watershed District*

North Twin Lake is a 69-acre lake located in Stillwater Township (Washington County). The maximum and mean depths of the lake are 1.8 m (roughly six feet) and 0.9 m (three feet), respectively. The lake's size and mean depth results in an approximate lake volume of 207 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's 187-acre immediate watershed translates to a small watershed-to-lake size ratio of 3:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This was the sixth year that North Twin Lake has been involved in CAMP (2000-2004 being the others). A search through the STORET nationwide water quality database for data on the lake provided limited information (1996-2004).

The lake's water quality was monitored seven times from mid-April to early-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	24.0	20.0	28.0	B
CLA (µg/l)	6.5	4.9	8.2	A
Secchi (m)	1.19	1.07	1.37	D
TKN (mg/l)	0.89	0.75	1.20	
Overall Grade				B

The 2005 overall grade was identical to those recorded in 1997, 1999-2001, and 2003-2004, and better than the C's recorded in 1996 and 1998.

This overall grade is skewed however, due to the shallowness of the lake. When examining the lake's mean TP and CLA concentrations, it seems that the lake's Secchi readings were limited by the shallowness rather than algal abundance. So, while the lake only received an overall grade of B, the actual water quality may have been better.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's quality seems well represented by an overall grade of B. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 1.8 for physical condition (between 1- "crystal clear" and 2- "some algae present"), and 2.4 for recreational suitability (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

North Twin Lake

Stillwater Twp., Washington Co.



LAKE ID: 820018

WD: Carnelian-Marine

Volunteer: Wash. Co. SWCD





◆ Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/20/05	18	17.8	5.04	5.09	9.1	32		1.2	2	3
5/17/05	13.4		5.87	6.12	4.9	20		1.1	1	2
6/16/05			4.82	5.33	5.8	23		1.4	2	2
7/14/05	29.5		5.78	6.5	7.1	28		1.1	2	3
8/12/05	24.6	24.4	6	6.61	8.2	24		1.2	2	2
9/6/05	24.8		8.37	11.12	6.4	25		1.2	2	3
10/6/05	14.3	14.4	7.95	7.7	7.1	29		1.2	2	3



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				C	B	B	A	B	B		B	A	B
Chlorophyll a				D	C	D	B	A	B		A	A	A
Secchi Depth				B	B	B	B	C	C	C	C	C	D
Overall				C	B	C	B	B	B		B	B	B

Source: Metropolitan Council and STORET data

Northwood Lake (27-0627) Bassett Creek Watershed Management Organization

Northwood Lake is a 15-acre lake located within the City of New Hope (Hennepin County). The mean and maximum depths of the lake are 0.8 m (roughly 2.5 feet) and 1.5 m (roughly five feet), respectively. The lake's size and mean depth results in an approximate lake volume of 41 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake's 1,341-acre immediate watershed translates to a small watershed-to-lake size ratio of 89:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This was the sixth year that Northwood Lake has been involved in CAMP. The lake was also enrolled in the program in 2000-2004. Other than the 2000-2004 CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty. Thus, 2000-2005 are the only years of available data.

The lake was monitored 13 times from mid-April to early-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	131.7	41.0	195.0	D
CLA (µg/l)	18.5	7.3	54.0	B
Secchi (m)	1.1	0.7	1.3	D
TKN (mg/l)	1.39	0.47	2.60	
Overall Grade				C




The lake's 2005 overall grade is similar to those recorded in 2002 and 2004, and better than those of 2000-2001 and 2003 (D).

Similar to past years, the Secchi transparency in 2005 would have been greater except for the shallowness of the lake. On numerous monitoring events, The Secchi disk was clearly noticeable while resting on the lake's bottom. Therefore, the lake's 2005 water clarity was actually better than that represented by the summer mean and resulting grade.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's quality seems well represented by an overall grade of D/C. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.


The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 4.2 for physical condition (between 4- "high algal color" and 5- "severe bloom"), and 4.3 for recreational suitability (between 4- "no swimming - boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/17/05	13.1				14	119		0.6	4	4
5/1/05	9.3				12	41		1.2	3	4
5/15/05	12.3				99	79		1.2	4	4
5/27/05	14				125	100		1		
6/5/05	18				17	107		0.9	3	4
6/26/05	22.2				54	179		0.8	4	4
7/23/05	26.4				73	168		1.2	5	4
8/7/05	28.5				12	158		1.3	5	5
8/21/05	22.9				85	155		1.3	5	5
9/4/05	21.1				73	195		1.2	5	5
9/18/05	22.6				75	145		0.7	4	4
10/1/05	18.6				49	122		1.1	4	4
					15	75		1.1	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								F	F	D	F	D	D
Chlorophyll a								B	C	B	C	B	B
Secchi Depth								D	D	D	D	D	D
Overall								D	D	C	D	C	C

Source: Metropolitan Council and STORET data



Oak Lake (10-0093) Carver County Environmental Services

Oak Lake is 339-acre lake located within Watertown Township (Carver County). The maximum depth of the lake is 3.4 m (roughly 11 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This was the fifth year in which Oak Lake has been involved in CAMP (2001-2004 being the others). The 2001-2004 CAMP data are the only known available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	122.2	26.0	209.0	D
CLA (µg/l)	56.0	4.3	150.0	D
Secchi (m)	1.1	0.3	3.0	C
TKN (mg/l)	2.07	0.73	2.80	
Overall Grade				C

The lake's 2005 overall grade was identical to those of 2001-2002 and 2004, and better than the D recorded in 2003.

As mentioned earlier, there are no water quality data available for Oak Lake other than 2001-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term however, the lake's overall grade seems well represented by a grade of C-/D+. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.2 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 3.1 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Oak Lake Watertown Twp., Carver Co.




Lake ID: 100093
WMO: Carver County
Volunteer: Ed Foley



● Sampling site
Contours in meters

2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/20/05	15.3		12.4		8.3	44		2.8	2	1
4/22/05	10.4		10.9		3.6	31		1.9	2	2
5/9/05	16.6		11.5		5.6	41		2	2	2
5/24/05	19.9		11.6		4.3			2.9	2	3
6/1/05	22.6		11.17		4.5	26		3	3	3
6/21/05	26		8.13		6.6	40		3	2	2
7/6/05	23.2		11.45		150	87		0.4	4	3
7/15/05	25.4		7.69		72	144		0.4	4	4
8/4/05	25.5				93	155		0.4	4	4
8/15/05	24.8		10.33		55	184		0.5	4	4
8/29/05					81	160		0.4	3	3
9/15/05	20.6		5.92		80	209		0.4	3	3
9/27/05	18.4		8.25		64	176		0.3	3	3
10/11/05	12.6				41	135		0.6	3	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus									D	D	F	D	D
Chlorophyll a									C	C	C	C	D
Secchi Depth									C	C	C	C	C
Overall									C	C	D	C	D

Source: Metropolitan Council and STORET data



O'Connor Lake (82-0002) *Lower St. Croix Valley Watershed Management Organization*

O'Connor Lake is a 38-acre lake located within Denmark Township (Washington County). There is very little known morphological data available for the lake.

This marks the first year in which O'Connor Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Therefore, 2005 is the only known year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored eight times between early-June and late-September, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary




<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	51.2	33.0	112.0	C
CLA (µg/l)	14.3	1.0	28.0	B
Secchi (m)	1.9	1.1	2.7	C
TKN (mg/l)	0.84	0.48	1.50	
Overall Grade				C

As mentioned earlier, there are no water quality data available for Lake O'Connor other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.3 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 3.9 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").


The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
6/5/05	23.1				10	33		2.4	2	4
6/19/05	27.4				15	47		1.9	2	4
7/10/05	28.9				28	38		2	2	4
7/24/05	27.8				24	59		1.1	2	4
8/6/05	26.2				22	57		1.5	3	3
8/14/05	24.3				10	35		1.2	3	4
9/5/05	20.9				8.9	34		2.7	2	4
9/25/05	17.8				5.3	112		2.4	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													C
Chlorophyll a													B
Secchi Depth													C
Overall													C

Source: Metropolitan Council and STORRET data



Olson Lake (82-0103) Valley Branch Watershed District

While Olson Lake has previously been monitored by Council staff, 2005 marks only the third year the lake has been monitored through CAMP (1993 being the first). The 89-acre lake with a mean and maximum depth of 2.1 (6.9 feet) and 4.5 m (14.8 feet), was monitored seven times from mid-April to early-October, 2005. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake's size and mean depth results in an approximate lake volume of 623 ac-ft.

The lake's surface area and watershed size (200 acres) translates to an 2:1 watershed-to-lake size ratio. Generally the larger the ratio, the greater the potential stress on the lake from surface runoff.

During each monitoring event the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	30.2	18.0	41.0	B
CLA (µg/l)	11.9	3.2	33.0	B
Secchi (m)	2.8	1.2	4.6	B
TKN (mg/l)	0.83	0.59	1.30	
Overall Grade				B

The physical and recreational conditions of the lake, as perceived by the volunteer monitor, were ranked on a 1-to-5 scale. The rankings are shown on the lake's information sheet on the next page. The mean physical condition ranking was 2.2 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.0 (2- "minor aesthetic problem").

Available data for Olson Lake reveals that lake water quality grades have improved since the 1980's. The lake water quality report card shown on the information sheet indicates that the lake received an overall C grade in 1984, as well as receiving Secchi grades of C in 1984-1986, and 1988-1990, before receiving overall grades of B in 1991, 1993, and 1995. More recently, the lake has recorded overall grades of A in 2000 and 2003-2004, before falling back to an overall grade of B in 2005. A recent MPCA conducted trend analysis on the lake's Secchi transparency data, revealed a statistically significant improvement in recent water clarity.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Olson

Lake Elmo, Washington Co.

Lake ID: 820103

WD: Valley Branch

Volunteer: Wash. Co. SWCD




● Sampling site

Contours in meters



2005 Data

Date	Surf Temp	Bot Temp	Surf DO	Bot DO	Chl a	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/21/05	15	13	6.78	6	3.1	23		4.7	2	2
5/23/05	15.7	14	5.65	5.07	3.2	18		4.3	2	2
6/16/05	24	18.1	6.57	3	4.2	29		4.6	2	1
7/14/05	29.9	23.1	9.07	0.11	6.2	38		2.3	2	2
8/12/05	26.1	25.1	5.08	0.89	13	25		1.8	2	2
9/9/05	22.1	21.8	7.92	0.6	33	41		1.2	3	3
10/3/05	18.2	17.8	7.26	2.16	25	40		1.4	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					C							B	
Chlorophyll a					C							B	
Secchi Depth					C	C	C		C	C	C	B	
Overall					C							B	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B		C					A			A	A	B
Chlorophyll a	A		B					A			B	A	B
Secchi Depth	B		B					A			A	A	B
Overall	B		B					A			A	A	B

Source: Metropolitan Council and STORET data



Orchard Lake (19-0031) City of Lakeville

Orchard Lake, managed by the MDNR as a centrachid lake (bass and panfish), is lake located within the City of Lakeville (Dakota County). The 250-acre lake has a 2,012-acre watershed, which translates to a 8:1 watershed-to-lake size ratio (generally the larger the ratio, the greater the potential stress on the lake from surface runoff). The maximum and mean depths of the lake are 10.0 m (roughly 33 feet) and 3.0 m (10 feet), respectively. The lake's surface area and mean depth translate to an approximate volume of 2,500 acre-feet. Approximately 75 percent of the lake's surface area are considered littoral zone (area of aquatic plant dominance). A public access is located within the City Park on the lake's southeastern end, and because of its multi-recreational uses, it is considered a "Priority Lake" in the Metropolitan Area.

This was the sixth year that Orchard Lake has been involved in CAMP (also involved in 1999-2001 and 2003-2004). Council staff has also monitored the lake in the past. A search through the STORET nationwide water quality database for data on the lake resulted in nutrient and Secchi transparency information for 1980-1981, 1983, 1989, 1993, 1998-2001, and 2003-2004, as well as just Secchi data for 1987-1988.

As part of the city's involvement in CAMP in 2005, the lake was monitored 14 times between late-April and mid-October. During each sampling event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	30.4	14.0	64.0	B
CLA (µg/l)	11.6	2.4	25.0	B
Secchi (m)	2.4	1.3	3.9	B
TKN (mg/l)	1.28	0.60	1.70	
Overall Grade				B

The lake's 2005 overall grade was similar to those recorded in 1981, 1983, 1989, 2001, and 2004, and better than the C's recorded in 1980, 1993, 1998-2000 and 2003. The lake's water quality seems to be well represented by an overall grade of C+/B.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the following page.

The mean physical condition ranking was 1.8 (between 1- "crystal clear" and 2- "some algae present"), while the mean recreational suitability ranking was 1.5 (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Orchard Lake

Lakeville, Dakota Co.

LAKE ID: 190031




WMO: Black Dog

Volunteers:

Tom and Dorothy
Goodwin


● Sampling site

Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CIA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	deg/L	ug/L	ug/L	M	1-5	1-5
4/29/05	9				2.6	29		3.3	1	1
5/1/05	10.1				4.8	18		3.0	1	1
5/19/05	13				5.7	18		3.5		
5/26/05	15.1				2.4	14		3.8	2	1
6/1/05	22				3.2	21		3.9	1	1
6/23/05	25				8.7	64		3.0	1	1
7/9/05	26				11	46		1.6	2	2
7/24/05	27				9.4	15		1.6	2	2
8/2/05	25.8				12	25		1.9	3	3
8/12/05	24				20	43		1.6	2	1
8/27/05	22				19	33		1.3	2	1
9/1/05	22				25	37		1.8	2	2
10/2/05	16				36	60		1.3	2	1
10/17/05	12				10	49		2.1	1	1




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C	B		B						B			
Chlorophyll a	B	B		B						B			
Secchi Depth	C	B		B				C	C	C	D	C	
Overall	C	B		B						B			

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C					C	C	C	B		C	C	B
Chlorophyll a	B					C	C	C	B		C	B	B
Secchi Depth	C					C	C	C	B		C	B	B
Overall	C					C	C	C	B		C	B	B

Source: Metropolitan Council and STORET data



Pamela Lake (27-0675) Conservation League of Edina

Pamela Lake is a 18-acre shallow lake (maximum depth of a 1.5 m [roughly 5 feet], located within Edina (Hennepin County). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This marks the second year in which Pamela Lake has been involved in CAMP (2004 being the first). Other than for the 2004 CAMP data, a search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 10 times between mid-May and late-September, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	80.8	42.0	200.0	D
CLA (µg/l)	18.8	5.0	48.0	B
Secchi (m)	1.1	0.7	1.4	D
TKN (mg/l)	1.36	0.67	2.20	
Overall Grade				C

When comparing the lakes TP (nutrient), CLA (algal biomass estimator), and Secchi (water clarity) grades, it is apparent that the TP and Secchi grades (and summer means) are quite a bit worse than the CLA grade. In a most cases, the three should be fairly comparable. One possible explanation for the lake's 2005 findings may be that the majority of the lake's TP comes from either in-lake suspended sediments (re-suspension), or the intrusion of sediment-laden runoff to the lake, which in turn lessens the clarity of the water and inhibits algal growth.

As mentioned earlier, there are no water quality data available for Pamela Lake other than the 2004-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.5 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 3.9 for recreational suitability (between 3- "swimming slightly impaired and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Pamela Lake

Edina, Hennepin Co.


LAKE ID: 270675

WD: Minnehaha Creek

Volunteer: Conservation League
of Edina

◆ Sampling site

Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
5/13/05	12.8				48	84		0.7	3	
5/26/05	21.3				11	143		1.1	2	4
6/10/05	24				20	53		1.4	3	2
6/24/05	30				21	200		1.3	2	2
7/8/05	29				5.1	50		1.1	2	5
7/22/05	31				15	60		0.8	5	5
8/5/05	29				5	42		1.2	2	
8/19/05	25				7.4	42		1.0	2	5
9/9/05	24.5				42	81		1.0	2	4
9/23/05	20				13	53		1.3	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												D	D
Chlorophyll a												B	B
Secchi Depth												C	D
Overall												C	C

Source: Metropolitan Council and STORM data



Parkers Lake (27-0107) Bassett Creek Watershed Management Organization

This was the fifth year that Parkers Lake has been involved in CAMP (it was first enrolled in 2000). The 97-acre lake, located within the City of Plymouth (Hennepin County), has a public access located within a city park on the lake's north end. One problem that may possibly hinder future recreational activity on the lake, however, is Eurasian Water Milfoil (*Myriophyllum spicatum*), which has been reported in the lake.

The mean and maximum depths of the lake are 3.7 m (roughly 12 feet) and 11.3 m (roughly 37 feet), respectively. The lake's size and mean depth result in an approximate lake volume of 1,164 ac-ft. Approximately 70 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance). The lake's 950-acre immediate watershed translates to a moderate watershed-to-lake size ratio of 10:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

The lake was monitored 15 times from mid-April to mid-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	37.6	18.0	64.0	C
CLA (µg/l)	12.1	2.4	39.0	B
Secchi (m)	2.6	0.8	4.9	B
TKN (mg/l)	1.03	0.52	1.80	
Overall Grade				B

While the lake's 2005 overall grade (identical to those recorded in 2003-2004) is better than the C's recorded in 1980, 1995, and 1999, it is worse than the recent A's recorded in 2000 and 2002,

A search through the STORET nationwide water quality database for data on the lake resulted in nutrient and Secchi transparency information for 1980, 1990, 1995, and 1999. The 2000 and 2002-2005 water quality years represent the lake's best-monitored water quality. The lake's water quality shows a markable improvement in water quality from 2000 to 2002, before slipping a little in 2003-2005. To better understand the lake's water quality and where it truly may be heading, continued monitoring is suggested.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.5 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 2.5 for recreational suitability (between 2- "minor aesthetic problem" and 3- swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Parkers Lake

Plymouth, Hennepin Co.



Lake ID: 270107

WMO: Bassett Creek

Volunteer: Bob Vdeen





◆ Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	14				6.2	37		1.7	2	3
5/4/05	11				4.3	56		2.5	1	1
5/16/05	13.8				3.2	35.5		3.0		
5/20/05	14.5				3.9	51		3.2	1	1
5/27/05	19				4.1	64		4.9	1	1
6/10/05	24				4	20		4.0	2	2
6/24/05	26.9					18		3.7	2	2
7/1/05	23				8	32		2.9	3	3
7/15/05	29				2.4	20		2.7	3	3
7/29/05	26				2.4	37		1.7	4	3
8/13/05	25.6				4.7	36		1.2	4	4
8/28/05	23.3				39	37		0.8	4	4
9/16/05	21.7				9.2	47		2.6	3	3
9/26/05	20.5				30	38		1.4	2	3
10/9/05	15				18	45		1.7	4	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C												
Chlorophyll a	C										B		
Secchi Depth	C										B		
Overall	C												

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			C				C	A		A	B	B	C
Chlorophyll a			B				B	A		A	B	A	B
Secchi Depth			C				C	B		A	B	C	B
Overall			C				C	A		A	B	B	B

Source: Metropolitan Council and STORET data

Peltier Lake (2-0004) Rice Creek Watershed District

Lake Peltier, with a surface area of 465 acres, is located one mile north of the City of Centerville (Anoka County). The maximum and mean depths of the lake are 4.9 and 2.1 m (16 and seven feet), respectively. The approximate volume of the lake is 3,255 ac-ft. The lake has a drainage area of roughly 68,082 acres, which translates to a extremely large watershed-to-lake size ratio of 391:1. The greater the ratio, the greater the potential stress on the lake from surface runoff. Public access is possible on the southwestern end of the lake through the Rice Creek Chain of Lakes Regional Park.

Peltier Lake is managed by the St. Paul Water Utility as a back-up water supply, and due to its multi-recreational uses, is considered a “Priority Lake” in the area by the Metropolitan Council. One aspect which may hinder recreational uses on the lake is the recent discovery of Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM]. Additionally, the lake, which is managed by the MDNR as a gamefish lake, experiences frequent winterkills.


Lake Peltier has been involved in CAMP since 1993 and was monitored 16 times from mid-April to mid-October, 2005. Results are presented in graphs and data tables on the following page.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP (µg/l)	203.5	62.0	399.0	F
CLA (µg/l)	104.1	2.4	380.0	F
Secchi (m)	1.6	0.5	3.2	C
TKN (mg/l)	2.09	0.85	4.60	
Overall Grade				D

The 2005 overall grade of D is similar to those recorded in 1993, 1995-1997, 1999-2000, and 2003-2004. This is better than the overall grade of F recorded in 1994, and 2001-2002.

Of the 13 years of CAMP data the best water quality was in 1995 and 1999, while the worst was 2001 (1994 was the second worst). Other than the 1993-2005 CAMP data, the only other data found through a search of the STORET database was from 1983. While statistical analysis on the lake's water quality



database revealed no “statistically significant” trends, and grades seems to promote the idea that the lake’s overall quality has remained fairly constant over the past decade [fluctuating between a low D and F], a simple trend line calculated from the annual summer means shows a slight degradation in the lake’s Secchi and chlorophyll-a means (see graph).

The average user perception rankings, on a 1-to-5 scale, was 3.0 for physical condition (3- “definite algae present”), and 2.6 for recreational suitability (between 2- “minor aesthetic problem” and 3- “swimming slightly impaired”).



The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Peltier Lake Centerville/Lino Lakes, Anoka Co.




Lake ID: 20004
WD: Rice Creek
Volunteer: Wayne LeBlanc

◆ Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/17/05	14				13	60.5		1.3	1	1
4/29/05	11				27	77.5		1.7	1	1
5/17/05	14.2				5.2	74		2.8		
5/18/05	13				2.4	88.5		3	1	1
5/28/05	17				4.2	63.5		3.2	1	1
6/4/05	19.6				2.8	320		3	1	1
6/26/05	26				380	242		0.6	5	5
7/10/05	26				330	234		0.6	4	4
7/22/05	29				90	138		0.6	4	3
8/2/05	26				87	214		0.6	4	3
8/12/05	26				68	274		0.6	4	3
8/21/05	23				120	399		0.5	4	4
9/2/05	23				14	305		1	3	3
9/19/05	22				47	345		0.6	4	3
10/1/05	16				65	304		0.8	4	3
10/9/05	12				25	206		0.9	4	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus				F									
Chlorophyll a				D									
Secchi Depth				D									
Overall				D									

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	F	F	D	F	D	F	F	F	F	F	D	F	F
Chlorophyll a	C	D	C	D	C	F	C	D	F	F	D	D	F
Secchi Depth	D	F	D	D	F	C	C	D	F	D	D	C	C
Overall	D	F	D	D	D	D	D	D	F	F	D	D	D

Source: Metropolitan Council and STORET data



Pike Lake [Ramsey Co.] (62-0069) Rice Creek Watershed District

Pike Lake is a 35-acre lake located within the City of New Brighton (Ramsey County). The mean and maximum depths of the lake are 2.1 m (7 feet) and 4.9 m (16 feet). The lake's mean depth and surface area translate to a lake volume of 245 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

This was the seventh year that Pike Lake has been involved in CAMP. A search through the STORET nationwide water quality database for data on the lake resulted in nutrient and Secchi transparency information for 1981-1983, 1985-1991, and 1999-2004, as well as just Secchi data for 1992-1993.

As part of the watershed district's involvement in CAMP in 2005, the lake was monitored 15 times between mid-April and mid-October. During each sampling event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	80.9	56.0	101.0	D
CLA (µg/l)	49.9	8.8	82.0	D
Secchi (m)	1.0	0.5	1.6	D
TKN (mg/l)	2.00	1.60	2.40	
Overall Grade				D

The lake's 2005 overall grade was similar to that of 1981-1982, 1987-1990, and 1999-2004, better than 1991 (F), and worse than that of 1983, and 1985-1986 (all of which were B's). Thus, the lake's quality seems fluctuate quite a bit, but mostly falls with the overall grade range of low-C/high-D.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the following page.

The mean physical condition ranking was 2.9 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.3 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Pike Lake



New Brighton, Ramsey Co.

Lake ID: 620069

W/D: Rice Creek




Volunteer: Philip Goodrich

◆ Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/19/05	16				66	103		0.7	2	2
5/7/05	14				18	74		1	3	2
5/15/05	12				16	77		1.1	2	2
5/21/05	16				88	96		1.6	3	2
6/1/05	24				39	79		1.2	3	2
6/13/05	25				42	68		1.3		
6/24/05	27				76	100		0.7	3	2
7/9/05	28				51	63		0.5	3	2
7/23/05	28				82	81		0.5	3	3
8/1/05	28				73	88		0.6	3	3
8/14/05	26				59	94		1.5	3	3
8/31/05	22				59	101		0.8	3	2
9/17/05	20				67	90		0.7	3	2
10/1/05	15				30	76		1.2	3	3
10/15/05	14				29	71		1.2	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus		C	C	D		C	C	D	D	D	D	D	
Chlorophyll a		C	D	A		A	C	C	C	D	C	F	
Secchi Depth		F	D	D		F	D	D	D	D	F	F	D
Overall		D	D	C		C	C	D	D	D	D	F	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	D	D	D	D	D	D
Chlorophyll a							C	C	D	C	D	C	D
Secchi Depth		D					D	D	C	D	D	D	D
Overall							D	D	D	D	D	D	D

Source: Metropolitan Council and STORET data



Pike Lake [Scott County] [Site-1] (70-0076) Prior Lake - Spring Lake Watershed District

Pike Lake is a 57-acre lake located within the City of Prior Lake (Scott County). The maximum depth of the lake is 2.7 m (roughly 9 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The majority of the land within the 1,991-acre watershed is undeveloped. The watershed-to-lake size ratio is 35:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This was the seventh year that Pike Lake has been involved in CAMP (the lake was also involved in 1997 and 1999-2004). Other than for the CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty. Thus, 1997 and 1999-2004 are the only years of available data.


Pike Lake was monitored 13 times from late-April to mid-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP (µg/l)	273.7	98.0	518.0	F
CLA (µg/l)	101.4	25.0	210.0	F
Secchi (m)	0.5	0.1	1.1	F
TKN (mg/l)	3.10	1.10	6.90	
Overall Grade				F

The lake's 2005 overall water quality grade was identical to that recorded in 2002-2004, and worse than the D's recorded in 1999-2001 and C recorded in 1997.

As mentioned earlier, there are no water quality data available for Pike Lake other than the 1997, and 1999-2005 CAMP data. While there is no "statistically significant" trend evident from the lake's database, the compilation of lake's grades, and a simple trend line calculated from the annual summer means, does seem to show that the lake's overall quality is decreasing. To better understand the lakes water quality and in which way it may be heading, continued monitoring is suggested.



Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The mean perceived physical condition of Pike Lake was 3.4 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability was 4.1 (between 4- "no swimming - boating ok" and 5- "no aesthetics possible").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



Pike Lake, Site 1

Prior Lake, Scott Co.

Lake ID: 700076





WD: Prior Lake-Spring Lake

Volunteers: David and Mona Hanson



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/26/05	13.1				26	12		0.7	2	4
5/15/05	13.4				28	98		0.6	3	4
5/19/05	18.7				26	105		0.8		
5/24/05	22				25	191		1.1	3	4
5/30/05	22.9				33	193		0.8	4	5
6/12/05	25.8				27	242		0.5	4	4
6/29/05	28				140	235		0.4	4	4
7/15/05	29.7				200	219		0.2	4	4
8/3/05	30.2				210	512		0.1	5	4
9/6/05	26.3				120	364		0.2	2	4
9/23/05	20.2				130	518		0.2	2	4
10/10/05	14.6				46	443		0.2	2	4
10/19/05	14.6				28	424		0.2	2	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus					D		D	D	D	F	F	F	F
Chlorophyll a					B		C	C	F	D	F	F	F
Secchi Depth					D		F	F	C	F	D	D	F
Overall					C		D	D	D	F	F	F	F

Source: Metropolitan Council and STORET data

Pine Tree Lake (82-0122) Rice Creek Watershed District

Pine Tree Lake, located on the eastern edge of the City of Dellwood (Washington County), covers an area of 174 acres and has a maximum depth of 7.9 m (26 feet). The mean depth of the lake, 3.0 m (10 feet), and its surface area translate to an approximate lake volume of 1,740 ac-ft. Because of its multi-recreational uses, it is considered a “Priority Lake” in the Metropolitan Area.

Pine Tree Lake has been a part of CAMP since 1993. In 2005, the lake was monitored 12 times between mid-May and mid-October. On each outing, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake’s perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	24.8	15.0	35.0	B
CLA (µg/l)	10.5	2.6	24.0	B
Secchi (m)	2.4	1.3	4.0	B
TKN (mg/l)	0.80	0.58	0.99	
Overall Grade				B

The lake’s 2005 overall is identical to those recorded in 1993-1994, 1997-2001 and 2004 and better than the C’s of 1995-1996 and 2002-2003 and the D recorded in 1985. No statistically significant long-term trend is evident from the lake’s overall water quality database (including TP, CLA, and Secchi data), in the short-term however, it seems that the lake’s overall water quality is well represented by a B/C grade

The physical and recreational conditions of the lake, as perceived by the volunteer(s), were ranked on a 1-to-5 scale. These rankings are shown in both table and graphic form on the following page. The mean physical condition ranking was 1.9 (between 1-“ crystal clear” and 2- “some algae present”), while the mean recreational suitability ranking was 2.5 (between 2- “minor aesthetic problem” and 3- “swimming slightly impaired”).

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.


If you notice any errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



	Surf. Tmp	Bot Tmp	Surf. DO	Bot DO	CLA	Surf. TP	Bot TP	Secchl	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	µm S	µm S
5/15/05	13				4.1	24		3	1	1
5/30/05	17				4.2	15		3	1	1
6/12/05	26.1				2.6	17		4	2	3
6/26/05	26.6				3.8	19		3	2	3
7/12/05	27.7				4.3	21		3	2	3
7/24/05	27.9				14	27		1.4	2	3
8/6/05	26.8				16	28		1.5	3	3
8/21/05	22.8				18	35			2	3
9/4/05	23				24	33		1.3	2	3
9/18/05	22.8				14	29		1.3	2	2
10/2/05	19.4				16	26		1.4	2	3
10/16/05	13.9				5	25		2.1	2	3

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus						C							
Chlorophyll <i>a</i>						D							
Secchi Depth						D							
Overall						D							

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B	B	C	C	B	B	B	C	C	C	C	B	B
Chlorophyll <i>a</i>	A	A	C	B	A	B	B	A	A	B	C	A	B
Secchi Depth	C	B	C	C	B	C	C	A	B	C	C	B	B
Overall	B	B	C	C	B	B	B	B	B	C	C	B	B



Powers Lake (82-0092) City of Woodbury

Powers Lake, located within the City of Woodbury (Washington County), has a surface area of approximately 57 acres (a shoreline length of 1.75 miles), and maximum depth of 12.5 m (41.0 feet). Approximately 50 percent of the lake's surface area is considered littoral, the shallow (0-15 feet) area dominated by aquatic vegetation. There is a public (canoe only) access on the northwest end of the lake near one of its two inlets. The lake has no outlet. Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has been reported on the lake.


Currently, about 47 percent of the lake's 1,238-acre watershed is open/undeveloped land with the rest either residential or open water/wetlands. Eventually nearly 84 percent of the lake's watershed will be developed as single-family and multi-family residential units. The lake's watershed-to-lake size ratio is 22:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

Powers Lake has been involved in CAMP since 1994. Between mid-April and mid-October, 2005, the lake was monitored 14 times. Similar to past years, the lake was monitored on each sampling date for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP (µg/l)	49.4	28.0	149.0	C
CLA (µg/l)	34.7	7.2	72.0	C
Secchi (m)	1.5	0.6	3.4	C
TKN (mg/l)	1.44	0.77	2.20	
Overall Grade				C

The lake's water quality in 2005 continues to be inferior to those recorded in 1994-1997 and 1999-2000. The lake has received overall grades of A in 1994, 1996, and 1999, B in 1995, 1997, 2000 and 2003, and C in 1998, 2001-2002 and 2004-2005.






Because of the wide fluctuation in the available data, no “statistically significant” long-term trend was determined. In the short-term however, the lake’s recent overall grades of C are worse than the A/B recorded in the 1990’s. Additionally the earlier graph reveals that the lake has experienced an increase in TP and CLA means over the past 10 years. More data are needed, however, to determine if this potential decrease in water quality falls within the lake’s normal range, or if the increased development around the lake has added to the lake’s nutrient load resulting in an increase in algal abundance and reduced clarity. Continued monitoring is suggested.

The physical and recreational conditions of the lake, as perceived by the volunteer, were ranked on a 1-to-5 scale and are displayed on the next page. The mean physical condition ranking was 3.3 (between 3- “definite algae present” and 4- “high algal color”), while the mean recreational suitability ranking was 4.2 (between 4- “no swimming – boating ok” and 5- “no aesthetics possible”).


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/1/05	9.8	4.7	6.73	0.47	9.3	53		2.1	2	3
4/22/05	15.2	5.7	7.37	0.19	7.8	40		2.6	2	3
5/9/05	15.1	6.4	7.43	0.86	7.2	38		3.4	3	3
5/23/05	14.5	7	5.9	0.27	13	32		2.6	3	4
6/7/05	22.8	7.8	5.69	0.32	35	38		1.2	3	4
6/20/05	26.2	9.1	9.42	0.58	32	37		1.1	2	4
7/5/05	23.3	10	8.21	0.13	47	45		1.1	4	4
7/18/05	28.3	10.5	9.5	0.79	43	40		1.4	3	4
8/3/05	27.5	9.9	8.23	0.5	56	47		0.8	5	5
8/17/05	25.5	10.2	12.39	0.41		149		0.6	4	5
8/29/05	24.1	10.1	5.88	0.43	39	47		0.9	4	5
9/12/05	23.5	10.2	5.95	0.55	21	42		2.0	2	4
9/30/05	18.5	10.4	6.3	0.58	16	28		1.7	3	4
10/13/05	15.3	10.3	8.72	0.52	42	165		1.4	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	B	B	A	A	C	A	B	C	B	C	C	C	C
Chlorophyll a	A	B	A	B	C	B	B	C	C	B	C	C	C
Secchi Depth	A	B	A	C	C	A	B	C	C	B	C	C	C
Overall	A	B	A	B	C	A	B	C	C	B	C	C	C

Source: Metropolitan Council and STORET data



Prior Lake [Lower Basin] [Site-1] (70-0026) Prior Lake - Spring Lake Watershed District

Prior Lake is divided into two distinct basins (the results of the 2005 monitoring on Prior Lake will be discussed as individual basins, Lower Prior and Upper Prior). Because of the lake's multi-recreational uses it is considered a "Priority Lake" in the Metropolitan Area.

The entire 1,167-acre lake is located within the City of Prior Lake (Scott County). The acreage of each basin is as follows: lower basin= 827 acres, and upper basin= 340 acres. The maximum and mean depths of the lower basin are 18.3 and 4.1 m (60 and 13 feet), which along with the surface area, translate to a lower basin volume of approximately 11,120 ac-ft. Roughly 46 percent of the lake's surface area is considered littoral, (the shallow [0-15 feet] area dominated by aquatic plants). The lower basin's 2,090-acre watershed translates to a rather small watershed-to-lake area ratio of 2.5:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff).

The lower basin's public access is located at the southern end of the lake. The lower basin of Prior Lake has one inlet (that from the upper basin of Prior Lake), and one outlet. The outlet structure, located on the southwestern portion of the basin, is a man-made structure that was installed to regulate surface water elevations. Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has been reported on the lower basin of the lake.

In an attempt to address issues either contributing to the eutrophication of Prior Lake or the symptoms from the resulting eutrophication, the Prior Lake - Spring Lake Watershed District has recently completed a Sustainable Water Quality Management Plan for its lakes (including Spring and Prior lakes). The Plan sets goals addressing the lakes' biological and chemical make-up and developed implementation strategies enabling the lakes' goals to be met (PLSLWD 2004).

While the Metropolitan Council has monitored the lower and upper basins of Prior Lake in the past, both basins have been a part of CAMP since 1997.

Lower Prior was monitored 12 times from mid-May to mid-October, 2005. On each sampling date the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	21.4	8.0	59.0	A
CLA (µg/l)	7.8	1.5	21.0	A
Secchi (m)	4.0	1.8	7.0	A
TKN (mg/l)	0.67	0.51	1.0	
Overall Grade				A

The 2005 overall grade is similar to those recorded in 1997 and 1999, and better than the B's recorded in 1996, 1998 and 2000-2004. The 2005 Secchi mean represents the lakes best recorded water clarity to date.


Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The mean perceived physical condition of Lower Prior Lake was 1.9 (roughly equal to 2- "some algae present"), while the mean recreational suitability was 1.3 (falling between 1- "beautiful" and 2- "minor aesthetics problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you detect any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Prior Lake, Lower Basin, Site 1

Prior Lake, Scott Co.



2005 Data

Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	CLA ug/L	Surf. TP ug/L	Bot. TP ug/L	Secchi M	PC 1 to 5	RS 1 to 5
5/15/05	12.2				2.2	9		7	1	1
5/19/05	14.7				2.7	10		6.5		
5/28/05	15.9				3	49		6	1	1
6/8/05	21.6				2.9	8		4	1	1
6/26/05	26.1				1.5	11		4.2	2	1
7/13/05	28.8				6.7	10		4.8	1	1
7/30/05	26.7				3.1	59		2.9	2	1
8/14/05	27.6				15	16		1.9	3	2
9/2/05	23.3				15	20		1.8	3	2
9/17/05	22.2				21	22		1.8	3	2
10/1/05	19.2				24	22		1.7	3	2
10/15/05	15.1				16	27		2.2	2	2





Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C	A			B								C
Chlorophyll a	B				B					A	B		B
Secchi Depth	C	C	B	C	B	C	B	C	C	B	B	C	C
Overall	C				B								C

Year	1993	1994	1995	1996	1997	1998 Site 1	1998 Site 2	1999 Site 1	1999 Site 2	2000 Site 1	2000 Site 2	2001 Site 1	2001 Site 2
Total Phosphorus				C	A	A	B	A	C	B	B	A	B
Chlorophyll a				A	A	B	C	A	B	B	B	B	C
Secchi Depth	B	B	B	B	B	C	C	B	C	B	C	B	C
Overall				B	A	B	C	A	C	B	B	B	C

Year	2002 Site 1	2002 Site 2	2003 Site 1	2003 Site 2	2004 Site 1	2004 Site 2	2005 Site 1	2005 Site 2
Total Phosphorus	B	C	C		B		A	
Chlorophyll a	B	C	A		B		A	
Secchi Depth	B	C	A		B		A	
Overall	B	C	B		B		A	

Source: Metropolitan Council and STORET data



Prior Lake [Upper Basin] [Site-1] (70-0072) Prior Lake - Spring Lake Watershed District

The maximum and mean depths of the upper basin of Prior Lake are 15.2 and 3.1 m (50 and 10 feet), respectively. The resulting water volume of the 340-acre upper basin is 3,460 ac-ft. About 93 percent of the lake's surface area is considered littoral, (the shallow [0-15 feet] area dominated by aquatic plants). The upper basin's 3,430-acre watershed translates to a watershed-to-lake area ratio of 10:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). The upper basin's public access is located at the northwestern end of the lake.

The upper basin of Prior Lake has two natural inlets, inflow from Spring Lake and the inlet from Rice and Crystal Lake drainage. Agriculturally derived non-point source nutrient loading released through the Spring Lake outlet heavily impacts water quality of the upper basin of Prior Lake.

The upper basin of Prior Lake was monitored nine times from mid-April to mid-September, 2005. Results are presented on graphs and data tables on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	54.7	24.0	79.0	C
CLA (µg/l)	45.5	3.7	110.0	C
Secchi (m)	1.6	0.6	4.0	C
TKN (mg/l)	1.27	0.62	2.40	
Overall Grade				C

Historical data for the upper basin of Prior Lake indicate that the water quality of the basin has remained fairly constant over the past decade fluctuating between overall grades of C and D. Lake quality grades (see the lake's information sheet on the following page) show that when nutrient data were collected on the upper basin of Prior Lake, overall grades ranged from C in 1981, 1990, 1996-1997, 2003 and 2005, and a D in 1980, 1984, 1989, 1998-2002, and 2004.

As apparent by the historic lake water quality grades, the lower basin of Prior Lake has better water quality than the upper basin. The reason being that the upper basin actually acts as a sort of detention basin for the lower basin. That is, the majority of the water entering the lakes goes through the upper basin first, allowing the settlement of sediments and associated nutrients before it enters the lower basin of the lake. The result is better quality water entering the lower basin of Prior Lake than is entering the upper basin.


Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The mean perceived physical condition of Upper Prior Lake was 3.5 (ranking between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability was 2.6 (falling between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you recognize any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Prior Lake, Upper Basin, Site 1

Prior Lake, Scott Co.



Regional Park Lake (82-0087) South Washington Watershed District

Regional Park Lake is a 16-acre lake located within the City of Cottage Grove (Washington County). The maximum depth of the lake is 5.8 m (roughly 19 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The majority of the land within the 600-acre watershed is undeveloped. The watershed-to-lake size ratio is 38:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). There is no formal boat access point on the lake.

This was the eighth year that Regional Park Lake has been involved in CAMP. Other than the 1998-2005 CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty. The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page. On each sampling date the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	48.7	26.0	82.0	C
CLA (µg/l)	23.7	3.2	67.0	C
Secchi (m)	1.9	0.9	3.1	C
TKN (mg/l)	0.88	0.58	1.20	
<i>Overall Grade</i>				C

The lake's 2005 overall grade is identical to that recorded in 1999 and 2004, and better than the D's of 1998 and 2000-2003.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's water quality seems well represented by an overall grade of D+/C. To better understand the lake's water quality and where it may be heading, continued monitoring is suggested.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.4 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.3 for recreational suitability (between 4- "no swimming - boating ok" and 5- "no aesthetics possible").



The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Regional Park Lake Cottage Grove, Washington Co.




Lake ID: 820086
WD: South Washington
Volunteer: City of Cottage Grove

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/11/05	13.8	6	7.08	1.2	19	58		1.2	3	4
4/22/05	15.9	8	0.16		19	62		1.2	2	3
5/9/05	16.6	8	7.63	1.59	6.7	37		2.4	2	4
5/23/05	16.5	9.3	5.89	0.27	4.6	41		2.4	3	4
6/7/05	23.3		4.63		3.2	39		2.7	3	4
6/20/05	24.6	13.4	11.58	0.5	26	41		1.7	3	4
7/5/05	22.8	15	10.08	0.28	50	69		0.9	4	4
7/18/05	27	15	8.23	0.13	67	82		1.1	3	4
8/3/05	26.7	15.4	7.7	0.61	25	57		1.5	5	5
8/17/05	23.9	15.9	8.45	0.31	31	45		1.8	4	5
8/29/05	23.8	22.3	4.78	0.36	11	53		1.7	5	5
9/12/05	22.3	17.7	6.01	0.57	9.8	46		3.0	2	4
9/30/05	16.6	15.1	8.1	6	26	26		1.8	3	4
10/13/05	14.9	12.9	6.84	2.87	6.2	45		1.4	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus						F	C	D	D	D	D	C	C
Chlorophyll a						B	B	C	C	D	C	C	C
Secchi Depth						F	D	F	F	F	F	D	C
Overall						D	C	D	D	D	D	C	C

Source: Metropolitan Council and STORET data



Riley Lake (10-0002) City of Chanhassen

While Riley Lake has previously been monitored by Council staff, 2005 marks the third year the lake has been monitored through CAMP. Riley Lake, with a surface area of 297 acres (2.9 miles in circumference), is located with the cities of Chanhassen and Eden Prairie (Carver and Hennepin counties). The maximum and mean depths of the lake are 15.0 and 6.6 m (49 and 21.6 feet), respectively. Roughly 34 percent of the lake's surface area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The approximate volume of the lake is 6,429 ac-ft.

The lake has a 4,796-acre immediate watershed, which translates to a watershed-to-lake area ratio of 16:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff). Public access is possible on the southeastern end of the lake. The lake is considered a "Priority Lake" by the Metropolitan Council" because of its multi-recreational uses. Eurasian Water Milfoil (*Myriophyllum spicatum*) [EWM] has been reported on the lake.

In 2005, Riley Lake was monitored 14 times from late-April to mid-October. On each outing, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary




<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	49.4	30.0	125.0	C
CLA (µg/l)	25.2	1.3	54.0	C
Secchi (m)	2.1	0.7	6.4	C
TKN (mg/l)	1.31	0.77	2.00	
Overall Grade				C

The lake's 2005 overall grade of C is identical to those recorded in each of the past years of monitoring. Therefore, the lake seems well represent by an overall grade of C.

The lake's average user perception rankings for 2005, on a 1-to-5 scale, were 3.6 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 2.3 for recreational suitability (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.



2005 Data

Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	CLA ug/L	Surf. TP ug/L	Bot. TP ug/L	Secchi M	PC 1 to 5	RS 1 to 5
5/3/05	9				1.5	56		6.4	2	1
5/16/05	11.8				1.3	125		5.1	3	2
5/20/05	18.6				25	55		2	5	4
6/10/05	22.5				17	37		2.2	5	4
6/23/05	25.1				12	30		1.3	4	3
7/8/05	25.1				20	35		1.5	3	2
7/11/05	25				24	33		1.5		
7/21/05	27				48	44		1	4	2
8/3/05	26.9				54	33		0.7	4	2
8/20/05	22.8					34		0.8	4	2
8/30/05	23.7				54	35		0.8	3	2
9/14/05	21.6				33	67		1.1	4	2
9/30/05	17				11	58		1.8	2	2
10/14/05	14.8				11	81		2.5	3	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C	B	C	C	C	C	C	C				C	
Chlorophyll a	C	C	C	C	C	C	C	D			C	C	
Secchi Depth	C	C	C	C	C	C	C	C	C		C	C	
Overall	C	C	C	C	C	C	C	C				C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C				C			C		C	C	C	C
Chlorophyll a	C				C			C		C	D	C	C
Secchi Depth	C				C			C		C	C	B	C
Overall	C				C			C		C	C	C	C

Source: Metropolitan Council and STORET data



Sand Lake (82-0067) *Marine on St. Croix Watershed Management Organization*

Sand Lake is a 46-acre lake located within New Scandia Township (Washington County). The lake has a surface area of 46 acres (1.8 miles in circumference) and a mean and maximum depth of 2.4 m (8 feet) and 5.5 m (18 feet), respectively. The lake, which has two inlets has an approximate volume of 368 ac-ft. Approximately 46 percent of the lake's surface area is considered littoral, the shallow (0-15 foot) area dominated by aquatic vegetation.

This was the eighth year that Sand Lake has been involved in CAMP (the lake was previously enrolled in 1993-1996 and 2002-2004). The 1993-1996 and 2002-2004 CAMP data were the only historic water quality data found for the lake. In 2005, the lake was monitored seven times from mid-April to early-October. During each event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP ($\mu\text{g/l}$)	59.8	24.0	135.0	C
CLA ($\mu\text{g/l}$)	44.6	3.1	90.0	C
Secchi (m)	1.6	0.3	3.1	C
TKN (mg/l)	1.42	0.79	1.90	
Overall Grade				C

The lake's 2005 overall grade is identical to those recorded in 1993-1996 and 2002-2003 and worse than the B recorded in 2004. While 2004 represents the lakes best-recorded water quality year, 1993 represents the worst (as determined by the individual parameter means).

The perceived conditions of the lake (both physical and recreational) were ranked on a 1-to-5 scale by the volunteer monitors. These user perception rankings are shown on the lake's information sheet. The mean physical condition ranking was 3.0 (3- "definite algae present"), while the mean recreational suitability ranking was 3.2 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

The Sand Lake water quality database consists of eight years of CAMP monitoring information for 1993-1996, and 2002-2005. Statistical analysis of the lake's water quality database failed to produce any statistically significant long-term trends. To better understand the lake's current water quality condition, and which direction it may be heading, continued monitoring is suggested. In the short-term, however, the lake's quality seems best described by a grade of C.



The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Sand Lake **New Scandia Twp., Washington Co.**




Lake ID: 820067
WMO: Marine-on-St. Croix
Volunteer: Wash. Co. SWCD

◆ Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 thru 5	1 thru 5
4/18/05	16.1	8	5.66	5.98	2.4	30		3.4	1	1
5/16/05	13.2	12.7	6	6.19	3.1	26		3.0	2	3
6/14/05	23.9	14.6	8.24	0.46	4	24		3.0	2	2
7/11/05	22		10.61	0.13	90	49		0.3	4	4
8/9/05	26.3	22.3	5.96	0.47	67	135		0.9	3	3
9/7/05	23	17.9	10.16	0.65	59	65		0.8	4	4
10/3/05	18.6	16.1	7.43	0.56	26	56		1.7	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C	C	C	C						C	C	B	C
Chlorophyll a	C	C	B	C						B	C	B	C
Secchi Depth	D	D	C	C						C	C	C	C
Overall	C	C	C	C						C	C	B	C

Source: Metropolitan Council and STORET data



Schmidt Lake (27-0102) Shingle Creek Watershed Management Commission

This was the sixth year that the 47-acre lake has been enrolled in CAMP. In fact, a search through Metropolitan Council and STORET databases indicated that the only other year besides 1995, 1998, 2000-2001 and 2004-2005 (CAMP data) for which data are available was 1994, when Secchi transparencies were taken as part of the MPCA's volunteer lake monitoring program.

The land uses within the lake's 190-acre immediate watershed are 77 percent low density residential and 23 percent wetland/water. The lake's watershed area to surface area ratio is 4:1. An area of concern and need for future management is the recent detection of Eurasian Water Milfoil (*Myriophyllum spicatum*) in the lake.

In an attempt to reduce the lake's algal population and improve the lake's water quality, an experimental bacterial treatment took place on Schmidt Lake in 2004 and 2005.

Schmidt Lake was monitored nine times between mid-May and late-September, 2005. During each event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The data and related graphs are presented on the information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	46.0	27.0	71.0	C
CLA (µg/l)	15.5	1.6	30.0	B
Secchi (m)	2.1	1.0	3.7	C
TKN (mg/l)	0.95	0.68	1.20	
Overall Grade				C

The lakes overall grade for 2005 of C is identical to those for 1995, 1998, 2000-2001, and 2004. Further analysis of the annual parameter means shows that the water quality in 2004 and 2005 (years with the in-lake bacterial treatments) was slightly better than those recorded in 1998 and 2000-2001, and very similar to that of 1995. The lake's two best water clarity means were recorded in 2004 and 2005.

Throughout the monitoring period, the volunteers ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The resulting user perception rankings are shown on the information sheet. The mean physical condition ranking was 2.4 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.1 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

Because of the limited size of the database (just Secchi data in 1994, and CAMP data in 1995, 1998, 2000-2001, and 2004-2005), no long- or short-term trends can realistically be determined. To better understand the quality of the lake and what direction it may be heading, more years of data collection are needed. In the very short-term however, the lake's current condition seems to be well represented by an overall lake water quality grade of C.


The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Schmidt Lake Plymouth, Hennepin Co.

Lake ID: 270102
WMO: Shingle Creek
Volunteer: Dale Wahlstrom

• Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/21/05	17.4				1.6	27		3.7	1	1
6/4/05	25				3.1	33		2.9	1	1
6/19/05	24.9				3.4	55		2.9	1	1
7/4/05	25.3				26	56		2	3	3
7/19/05	27.9				30	71		1	4	4
7/25/05	27				27	46		1		
7/31/05	27.9				25	60		1	4	3
8/28/05	23.8				13	39		1.7	3	2
9/25/05	19.7				22	27		2.1	2	2

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			C			C		C	C			C	C
Chlorophyll a			B			C		C	C			B	B
Secchi Depth		C	C			C		C	D			C	C
Overall			C			C		C	C			C	C

Source : Metropolitan Council and STORM data



School Lake (13-0057) *Comfort Lake-Forest Lake Watershed District*

School Lake is a 48-acre lake located near Chisago City (Chisago County). There is very little known morphological data available for the lake.

This marks the first year in which School Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake was unsuccessful. Therefore, 2005 is the only known year of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.




2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	64.5	40.0	99.0	C
CLA (µg/l)	42.1	2.6	73.0	C
Secchi (m)	1.4	0.8	3.1	C
TKN (mg/l)	1.47	0.80	2.00	
Overall Grade				C

As mentioned earlier, there are no water quality data available for School Lake other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.3 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 3.5 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/18/05	14.3	5.7	7.4	0.18	26	70		1.1	3	2
4/25/05	11.8	6.4	5.87	0.08	26	77		1.4	2	3
5/13/05	12.2	7.5	6.12	0.19	29	69		1.5	2	2
5/24/05	16.1	8.3	4.99	0.07	7	40		3.0	3	3
6/9/05	22.1	9.6	5.41	0.32	26	55		1.5	4	4
6/21/05	24.6	10.3	10.67	0.57	47	47		1.5	3	3
7/1/05	23.5	11.2	12.26	0.55	64	87		0.9	3	3
7/22/05	27	11	7.52	0.59	27	63		1.4	3	4
8/2/05	27.2	11.5	10.54	0.51	73	70		0.9	4	4
8/16/05	24.4	11.8	9.46	0.38	70	74		1.1	5	4
9/1/05	21.9	12.1	7.91	0.48	2.6	74		1.2	3	3
9/14/05	21.4	12.8	6.1	0.4	61	99		1.1	3	4
9/29/05	17.2	13.4	5.37	0.62	57	84		0.8	3	4
10/11/05	13.7	12.8	4.68	3.05	33	74		1.2	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													C
Chlorophyll a													C
Secchi Depth													C
Overall													C

Source: Metropolitan Council and STORET data



Schroeder's Pond (82-0301) *Carnelian - Marine Watershed District*

Schroeder's Pond is a small land-locked lake located within May Township (Washington County). The maximum depth of the lake is 3.0 m (roughly 10 feet). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column

This was the second year that Schroeder's Pond has been involved in CAMP. On each of the sampling days the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The lake was monitored 14 times between mid-April and mid-October, 2005.

The following are the averages for each of the parameters tested.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	104.6	25.0	216.0	D
CLA (µg/l)	50.2	7.0	150.0	D
Secchi (m)	1.4	0.9	2.1	C
TKN (mg/l)	1.04	0.41	1.80	
Overall Grade				D

Other than for the 2004-2005 CAMP data, there are no known water quality data available for Schroeder's Pond. Therefore it is not possible to determine any long-term or short-term trends. The lake's water quality in 2005 however, was dramatically worse than that recorded in 2004 (overall grade of B). The lakes extremely poor water quality from mid-August to mid-September, 2005 was the main reason for the poor overall grade. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


The perceived physical and recreational conditions (ranked on a 1-to-5 scale) are shown on the lake's information sheet on the next page. The average user perception rankings, were 2.6 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 4.2 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible"). Similar to that shown in the lakes 2005 grades, the 2005 user rankings are also considerably worse than those of 2004.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Schroeder's Pond May Twp., Washington Co.

Lake ID: 820301
W/D: Carnelian - Marine
Volunteer: Wash. Co. SWCD

• Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	13	8.7	4.54	0.25	12	54		1.7	3	4
4/25/05	11.5	10.1	4.34	3.03	7	55		1.8	2	4
5/10/05	17.3	14.2	4.05	2.38	7.3	52		2.1	3	4
5/24/05	17.6	10.6	4.31	0.07	7	50		1.8	3	4
6/6/05			4.09	0.24	18	25		1.5	2	4
6/23/05	23.1	14.5	5.53	0.43	24	83		0.9	2	4
7/6/05	21.2	15.6	6.03	0.69	21	75		1.7	2	4
7/19/05	25.1	16	2.12	0.36	36	130		1.1	3	5
8/6/05	23.1	17.9	1.36	0.52	14	81		1.4	3	5
8/19/05	20.8	19.3	1.32	0.31	150	172		1.2	3	4
8/30/05	21.3	16.7	2.11	0.36	130	216		1.4	3	4
9/13/05	21.1	16.7	7.54	0.35	120	210		1.4	2	4
9/28/05	15.8	15.7	7.52	5.06	25	96		1.4	3	4
10/10/05	10.9	10.2	6.03	5.9	9	43		2.4	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												C	D
Chlorophyll a												A	D
Secchi Depth												C	C
Overall												B	D

Source: Metropolitan Council and STORET data



Schutz Lake (10-0018) Minnehaha Creek Watershed District

Schutz Lake is a 105-acre lake located within the City of Victoria (Carver County). The maximum and mean depths of the lake are 15.0 m (roughly 49.2 feet) and 6.0 m (19.5 feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 2,100 ac-ft. Approximately 27 percent of the lake's area is considered littoral zone (area of aquatic plant dominance).

The lake's 943-acre immediate watershed and surface area translates to a watershed-to-lake size ratio of 9:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). There is no formal boat access point on the lake. An area of concern and need for future management is the presence of Eurasian Water Milfoil (*Myriophyllum spicatum*) in the lake. The MNDNR conducted a 2001-2003 study on the use of a herbicide (fluridone) for selective control of EWM on Schutz Lake (as well as five other area lakes). While the results of the study indicate that the fluridone treatment controlled the EWM (immediately after treatment in mid-summer 2002 through 2003), the abundance of native submersed vegetation also declined dramatically (MNDNR 2004). On the flip side, however, the frequency of curly-leaf pondweed increased (MNDNR 2004).

This was the sixth year that Schutz Lake has been involved in CAMP (the lake was also monitored by Council staff back in 1984). Other than the 1984 and 2000-2005 data, a search through the STORET nationwide water quality database solely includes Secchi data (1981-1988 and 1990-1991).

The lake was monitored 11 times from early-May to mid-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	46.2	23.0	74.0	C
CLA (µg/l)	35.2	10.0	58.0	C
Secchi (m)	1.9	1.4	2.7	C
TKN (mg/l)	1.10	0.63	1.50	
Overall Grade				C

The lake's 2005 overall grade is identical to that recorded in 2003 and 2004 and worse than the B's recorded in 1984 and 2000-2002.

As mentioned earlier, other than the 1984 and 2000-2005 Council data, the lake's water quality database includes Secchi readings from 1981-1988 and 1990-1991. The lake does not seem to show any long-term water clarity trends and seems to be represented with a water quality grade of C. The limited TP and CLA data however, reveals a recent (2003-2005) decline in TP and CLA grades as compared to the earlier grades (2000-2002). The reason for this is unknown. To better understand the lake's TP and CLA conditions and to determine any possible trends, continued monitoring is suggested.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.8 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 1.3 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries



Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Schutz Lake Victoria, Carver Co.




Lake ID: 100018
WD: Minnehaha Creek
Volunteer: Mike Shouldice

• Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/8/05	11.7				10	65		2.1	1	1
5/30/05	20.2				46	72		1.4	4	3
7/5/05	24.2				54	38		1.5	3	1
7/10/05	26.2				12	74		2.1	3	1
7/24/05	27.8				19	35		1.7	3	1
8/7/05	26.7				26	23		1.5	3	1
8/28/05	25				58	30		2.2	3	1
9/10/05	22.5				53	36		2.7	2	
9/25/05	19.8				39	43		1.8	3	1
10/2/05	18.2				23	47		1.7	3	1
10/16/05	13.9				13	48		2.3	4	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					C								
Chlorophyll a					A								
Secchi Depth		C	C	C	C	C	C	C	C	C	C	D	
Overall					B								

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								B	B	B	C	C	C
Chlorophyll a								A	B	B	B	C	C
Secchi Depth								C	C	B	C	C	C
Overall								B	B	B	C	C	C

Source: Metropolitan Council and STORET data



Seidl's Lake (19-0095) Cities of *Inver Grove Heights and South St. Paul*


Seidl's Lake is a 14-acre lake located in the City of Inver Grove Heights (Dakota County) which receives inflow from five inlets. Other than the fact that the maximum depth of the lake is approximately 5.0 m (17 feet), there is very little known morphological data available. The lake has been enrolled in CAMP since 1995. In 2005 it was monitored 14 times from mid-April to mid-October. On each sampling date the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	65.7	49.0	88.0	D
CLA (µg/l)	28.5	9.0	46.0	B
Secchi (m)	0.8	0.5	1.4	C
TKN (mg/l)	1.03	0.89	1.20	
Overall Grade				C

Similar to that reported in past lake reports (and noticed again in 2005), the difference between the TP, CLA and Secchi grades in current and past years (see report grade on the lake's information page), may indicate that suspended sediments play a large role in the inner workings of the lake. This scenario can be fairly typical for small shallow lakes where wind action and storm sewer inflow either increase the influx of sediments to the system or cause the re-suspension of existing bottom sediments. That is, the suspended sediments influence the lake's phosphorus make-up (a larger portion of the in-lake phosphorus in particulate form rather than a soluble form more readily available for algal uptake), reduce water clarity, and could actually be limiting the amount of light available for algal growth, thus keeping the CLA concentrations down (resulting in a better than expected grade).

The water quality database for Seidl's Lake consists of nutrients and Secchi data in 1991, Secchi data in 1993-1994, and CAMP data in 1995-2005. While the lake's database is expanding, it is lacking in pre-1995 data. Statistical analysis on the lake's water quality database revealed no "statistically significant" long-trends. A simple regression on the lake's available water quality data, however, does reveal a slight decrease in the lakes quality. Over this time span, The lake's overall water quality grades fluctuated between an overall C and low-B grade in 1991-1998, 2000-2001 and 2003-2005, and a low grade of D in 1999 and 2002.



In an attempt to address the lake's possible degradation concerns and watershed influences on said degradation, lake area homeowners have been trying to work with the local communities to address areas of concern to the lake's future management. They are currently working on garnering city involvement and outside funding to further initiate the lake planning/improvement process.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The summertime mean recorded physical condition was 3.6 (between 3- "definite algae present" and 4- "high algal color"), while the mean suitability for recreation ranking was 4.0 (4- "no swimming - boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Seidl Lake


Iver Grove Heights,
Dakota Co.

Lake ID: 190095

WMO: Lower Mississippi River

Volunteers: Janet and Harv Bartz

- Sampling site
- Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 to 5	1 to 5
4/10/05	14.3				57	136		0.6	3	↓
4/24/05	13.9				21	75		0.9	2	↓
5/10/05	17.1				9	71		1.4	2	↓
5/23/05	18.4				20	58		1.1	3	↓
6/9/05	24.8				17	69		0.8	3	↓
6/30/05	23.1				34	84		0.5	↓	↓
7/6/05	25.8				46	76		0.6	↓	↓
7/18/05	28.6				43	59		0.6	↓	↓
8/2/05	28.7				28	58		0.8	↓	↓
8/16/05	25.1				29	49		0.9	↓	↓
8/27/05	25				30	88		0.5	↓	↓
9/14/05	22.4				23	50		0.9	↓	↓
9/27/05	19.4				34	61		0.8	↓	↓
10/10/05	15.2				9.6	107		1.2	3	↓

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus												C	
Chlorophyll a												C	
Secchi Depth												D	
Overall												C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			C	C	C	C	D	C	C	D	C	D	C
Chlorophyll a			A	B	B	C	C	C	C	C	B	B	C
Secchi Depth	D	D	B	B	C	D	D	C	C	D	D	C	D
Overall			B	B	C	C	D	C	C	D	C	C	C

Source: Metropolitan Council and STORET data



Shaver Lake (27-0086) City of Minnetonka

Shaver Lake is a small 11-acre lake located within the cities of Deephaven, Minnetonka, and Woodland (Hennepin County). There is very little known morphological data available for the lake.

This was the first year that Shaver Lake has been involved in CAMP. On each of the sampling days the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The lake was monitored 14 times between mid-April and mid-October, 2005. The following are the averages of the three events for each of the parameters tested.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	54.9	41.0	70.0	C
CLA (µg/l)	7.9	1.0	22.0	A
Secchi (m)	1.17	1.00	1.30	D
TKN (mg/l)	0.77	0.48	1.10	
Overall Grade				C

When comparing the lakes TP (nutrient), CLA (algal biomass estimator), and Secchi (water clarity) grades, it is apparent that the TP and Secchi grades (and summer means) are quite a bit worse than the CLA grade. In a most cases, the three should be fairly comparable. One possible explanation for the lake's 2005 findings may be that the majority of the lake's TP comes from either in-lake suspended sediments (re-suspension), or the intrusion of sediment-laden runoff to the lake, which in turn lessens the clarity of the water and inhibits algal growth.

Because 2005 is the only year of known water quality data for the lake, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


The perceived physical and recreational conditions (ranked on a 1-to-5 scale) are shown on the lake's information sheet on the next page. The average user perception rankings, were 4.0 for physical condition (4- "high algal color"), and 4.1 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Shaver Lake

Deephaven, Minnetonka and Woodland
Hennepin Co.

Lake ID: 270086
WD: Minnehaha Creek
Volunteers: The Davis Family



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/18/05	17.5				4	32		1.3	5	4
4/28/05	10.9				12	34		1.3	5	4
5/15/05	14.3				2.6	50		1.3	3	3
5/28/05	18.3				2.2	70		1.3	3	3
6/13/05	25.2				1	69		1.3	3	4
6/28/05						42			5	4
7/12/05					2.4	50		1.2	5	5
7/28/05					3.4					
8/1/05	29.1				22	65		1.1	5	5
8/22/05	24.1				14	54		1.0	5	5
9/3/05	20.7				5.4	41		1.0	4	5
9/16/05	23.5				18	53		1.2	3	3
10/2/05	18.5				12	40		1.3	3	4
10/16/05	15.1				12	69		1.4	3	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus													C
Chlorophyll a													A
Secchi Depth													D
Overall													C

Source: Metropolitan Council and STORET data



Shields Lake (82-0162) *Comfort Lake-Forest Lake Watershed District*

Shields Lake is located in the City of Forest Lake in Washington County. The lake has a surface area of 27 acres (0.8 miles in circumference) and a maximum depth of 8.2 m (27 feet). About 85 percent of the lake's area is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation.

Shields Lake has been involved in CAMP since 1993. The lake was monitored 14 times between mid-April and mid-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary


<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	177.0	60.0	357.0	F
CLA (µg/l)	50.7	4.4	91.0	D
Secchi (m)	1.1	0.5	3.1	D
TKN (mg/l)	2.06	1.10	2.60	
Overall Grade				D

The 2005 CLA and Secchi means were the worst recorded to date. The lakes best recorded water quality was in 1997 (TP mean of 159.0 µg/l, CLA mean of 7.0 µg/l, and a Secchi mean of 2.8 m).

The Shield Lake water quality database includes 14 years (1991, 1993-2005) where TP, CLA, and Secchi transparency data are available to calculate annual grades. The overall grades range from C's in 1991, and 1994-1997, to D's in 1993 and 1998-2005. Additional data found for 1988-1989 had only TP and CLA concentrations, and 1990 had only Secchi depth information.

A quick look at the lake's database seems to show that the its TP concentrations have remained consistently high (between D and F). The CLA and Secchi numbers, which improved slightly after the biomanipulation of 1994, degraded in 1998 and 1999 and, until 2002-2005, had shown some improvement in 2000-2001. However, because of the absence of historic data and the great variability of existing data, statistically accurate long-trend analysis is difficult. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

While the lake's CLA concentrations (corresponding to algal abundance) and Secchi transparencies had shown some improvement in 2000-2001, the recent 2002-2005 data show conditions similar to those of 1998-1999. It was mentioned in the 1998 and 1999 lake reports that TP, CLA, and Secchi transparency conditions in 1998 were similar to that of 1994 and 1995 until early-July, and after the big storms (extreme winds, heavy rains) of late-June, when the lake started to experience below normal water quality conditions. It was further mentioned that, during the July 7, 1998 monitoring event 12 dead turtles were seen at the lake's inlet, no "freshwater shrimp" (zooplankton) were seen in the water samples, and the lake had no oxygen in the water below one meter (approximately three feet). At this point the lake's TP concentrations rose, but more abnormally, the lake's CLA and Secchi readings dramatically worsened. The lake's CLA and Secchi readings did not start to rebound until again until late-September of that year. It is thought that the 1998 storms started the degradation of the lake's water quality by increasing runoff from the surrounding watershed, and riling up the lake's sediments. This potentially resulted in an increased TP and sediment load to the lake, increasing turbidity, TP concentrations, algal populations, and reducing water clarity and in-lake oxygen levels (Anhorn 1999).



The perceived physical and recreational conditions of the lake, recorded by the volunteer(s), were ranked on a 1-to-5 scale. The rankings are shown in both tabular and graphical form on the lake's associated information sheet. The mean physical condition ranking was 3.3 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability ranking was 4.0 (4- "no swimming - boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Shields Lake

Forest Lake,
Washington Co.



Lake ID: 820162

WMO: Comfort Lake - Forest Lake

Volunteer: Wash. Co. SWCD

● Sampling site

Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/14/05	13.3	4.2	9.36	0.62	50	130		1.1	3	4
4/26/05	11.1	4.3	5.95	0.15	31	93		1.4	3	4
5/13/05	12.3	4.6	5.46	0.19	4.4	76		3.0	2	4
5/24/05	19.1	5.1	5.68	0.04	6.6	60		2.9	2	4
6/9/05	25.3	5.9	4.76	0.36	35	169		1.2	3	4
6/22/05	25	7.2	17.4	0.67	91	214		0.6	3	3
7/7/05	25.8	6.8	9.45	0.65	49	290		0.8	4	4
7/22/05	29.9	7.1	9.43	0.67	55	152		0.6	4	5
8/2/05	26.9	7.8	8.53	0.56	90	357		0.5	3	4
8/16/05	27.3	7.5	11.75	0.43	49	160		0.6	4	4
9/1/05	22.5	8	6.8	0.53	49	156		0.9	4	4
9/14/05	22.7	8.4	7.74	0.42	51	135		0.8	3	4
9/30/05	15.7	8.7	7.26	0.73	78	178		0.6	4	4
10/14/05	13.2	9.2	9.96	0.49	74	187		0.8	3	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus									F	D		D	
Chlorophyll a									D	D		C	
Secchi Depth											F	C	
Overall												C	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	F	D	F	F	F	F	F	F	F	F	F	F	F
Chlorophyll a	C	C	C	B	A	C	C	C	C	C	C	C	D
Secchi Depth	C	C	B	B	B	C	C	C	C	C	C	C	D
Overall	D	C	C	C	C	D	D	D	D	D	D	D	D

Source: Metropolitan Council and STORET data



Silver Lake (82-0016) *Carnelian - Marine Watershed District*

Silver Lake is a 98-acre lake located within Stillwater Township (Washington County). The maximum and mean depths of the lake are 3.4 m (roughly 11 feet) and 1.7 m (five-and-a-half feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 549 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's 455-acre watershed and surface area translates to a watershed-to-lake size ratio of 4.6:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). There is no formal boat access point on the lake.

This was the sixth year that Silver Lake has been involved in CAMP (although just Secchi transparencies were collected during three of those years). A search through the STORET nationwide water quality database for data on the lake produced a limited amount of data. The only years in which data were found, other than the 2000-2004 CAMP data, was 1997-1999. The only years of which included TP, CLA and Secchi transparency data are available are 1997-2000 and 2004-2005.

The lake was monitored seven times between mid-April and early-October, 2005. During each event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	36.8	23.0	48.0	C
CLA (µg/l)	9.52	5.8	17.0	A
Secchi (m)	2.2	1.7	2.7	B
TKN (mg/l)	0.80	0.59	0.95	
Overall Grade				B

The lake's 2005 overall grade is identical to that recorded in 2004, and better than the overall grades of C in 1996-1998 and 2000-2001, and D in 1999. When looking at the grades and individual parameter means, it is apparent that 2004 was the lakes best-recorded water quality year.


Because of the nature of the lake's water quality database the determination of any statistically significant long-term trend detection is not possible. In the short-term however, the lake's water quality seems to be well represened by an overall grade of C+/B-. To better understand the lake's overall water quality and where it may be heading, continued monitoring is suggested.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.6 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 3.0 for recreational suitability (3- "swimming slightly impaired").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.




Silver Lake

Stillwater Twp., Washington Co.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1 to 5	1 to 5
4/20/05	16.7	16.4	5.49	5.53	7.6	31		2.3	2	3
5/17/05		12.8	5.93	5.26	6.3	23		2.7	2	2
6/16/05	24.3	22.6	7.88	4.46	11	46		2.4	3	3
7/13/05	31.7	25	9.21	1.1	17	36		1.7	3	4
8/9/05	27	23.3	6.86	0.37	7.5	36		1.8	2	3
9/6/05	20.7	20.7	8.68	0.52	5.8	39		2.4	3	3
10/6/05	15.6	15.6	7.2	4.86	10	37		2.4	2	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				C	C	C	D	C	C			B	C
Chlorophyll a				C	C	C	D	B	B			A	A
Secchi Depth				C	D	D	D	C	C	C	B	B	B
Overall				C	C	C	D	C	C			B	B

Source: Metropolitan Council and STORET data



South Rice Lake (27-0645) Bassett Creek Watershed Management Organization

South Rice Lake is a 3.2-acre lake located within the City of Golden Valley (Hennepin County). The maximum and mean depths of the lake are 2.5 m (roughly 8 feet) and 0.5 m (one-and-a-half feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 5.4 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's 63-acre immediate watershed and surface area translates to a watershed-to-lake size ratio of 20:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). When including the lake's whole contributing watershed (including flow from Grimes Pond and North Rice Lake), however, the size increases to 514 acres (160:1) (Barr 1997).

This was the sixth year that South Rice Lake has been involved in CAMP (it was also involved in 2000-2004). Other than the 2000-2005 CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty. The lake was monitored 15 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	132.8	51.0	304.0	D
CLA (µg/l)	22.6	5.3	50.0	C
Secchi (m)	0.7	0.3	1.1	D
TKN (mg/l)	1.57	0.52	3.8	
Overall Grade				C

Of the six years of monitoring data available for the lake, it is apparent that the lake experienced its best water quality in 2004 and its the worst water quality was recorded in 2000. The lake received overall grades of F in 2000, D in 2001-2003 and 2005, and C in 2004.

A recent in-lake alum treatment (applied at ice-off in mid-April, 2002) was meant to lower phosphorus levels, control algal growth and improve water clarity. It was reported in the 2002 Lake Report that the alum treatment was successful in the reducing of in-lake TP and CLA (indicating a reduction in algal biomass) in 2002. While, the lake's 2002, and 2004-2005 water quality concitions were better than pre-alum treatment, the 2003 water quality was not. In fact, the 2003 summer mean TP concentration was more than double those recorded in 2002 and 2004-2005. Additional years of monitoring are needed to truly determine the effectiveness and long-term efficiency of the alum treatment.




The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.8 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.4 for recreational suitability (between 4- "no swimming - boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

South Rice Pond Golden Valley, Robbinsdale, Hennepin Co.


Lake ID: 270645
WMO: Bassett Creek
Volunteer: Steve Striff

• Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/10/05	14.3				42	124		1	2	4
4/24/05	12.3				13	67		1	2	4
5/8/05	16.8				13	85		0.9	2	4
5/16/05						70		1		
5/29/05	18.5				7	77		1.1	3	4
6/12/05	26.3				5.3	51		0.9	4	4
6/25/05	25.2				20	55		0.7		
7/9/05	25.2				32	55		0.6	4	5
7/23/05	25.8				23	304		0.3	5	5
8/6/05	25.2				50	221		0.4	5	5
8/21/05	20.2				14	243		0.3	5	5
9/5/05	20.2				26	66		0.8	3	4
9/18/05	23.5				36	234		0.7	3	4
10/2/05	17.9				23	63		1	2	4
10/15/05	13.4				13	72		1	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								F	F	D	F	D	D
Chlorophyll a								F	B	B	C	A	C
Secchi Depth								F	F	F	F	D	D
Overall								F	D	D	D	C	D

Source: Metropolitan Council and STORET data



South School Section Lake (82-0151) Browns Creek Watershed District

South School Section Lake, located in southeastern Hugo Township in Washington County, was monitored 7 times between late-April and mid-July 1996. The 125-acre lake has a maximum depth of 8.0 m (26 feet). About 41 percent of the lake's area is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation.

On each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. In 2005, the lake was monitored 14 times between mid-April and mid-October.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	52.7	23.0	98.0	C
CLA (µg/l)	32.9	3.5	80.0	C
Secchi (m)	2.3	0.8	5.2	B
TKN (mg/l)	1.20	0.66	1.90	
Overall Grade				C

While the lake's 2005 overall grade was identical to that of 1995, 1996, and 1998, the 2005 mean Secchi reading is the best recorded to date.

The physical and recreational conditions of South School Section Lake as perceived by the volunteer(s) were ranked on a 1-to-5 scale. These rankings are shown on the lake's information sheet on the next page. The summertime mean physical condition was 3.4 (between 3- "definite algal presence" and 4- "high algal color"). The mean suitability for recreation ranking was 3.6 (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

Because data for South School Section Lake are only available for 1995, part of 1996, 1998 and 2005, no long- or short-term trends can be determined. Additional year of data are needed to better understand the lake's water quality and what direction it may be heading.



The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

South School Section Lake, Hugo, Washington Co.




Lake ID: 820151
WMO: Browns Creek
Volunteer: Wash. Co. SWCD

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	10.5	9.5	6.9	0.78	6	106		5.8	1	2
4/28/05	10.9	10.9	6.47	2.34	3.6	21		4.4	1	1
5/9/05	13.9	12.3	6.49	0.13	3.5	36		5.2	2	2
5/26/05	15.2	14.2	5.1	2.4	4.5	23		4.1	3	3
6/8/05	21.3	16.2	4.71	1.35	4.6	45		4.0	3	4
6/23/05	25.5	16.5	8.4	0.34	6.4	32		4.1	2	3
7/5/05	22.9	18	7.53	0.08	22	37		1.8	3	3
7/25/05	23.5	18.5	7.65	0.15	15	45		1.4	4	4
8/5/05	24.9	22.4	4.64	0.61	80	55		0.9	4	4
8/19/05	23.3	21.2	6.14	0.29	63	72		0.8	4	4
8/30/05	24.6	21.8	11.1	1.54	66	98		0.8	4	4
9/15/05	21.4	21.3	5.93	5.28	44	74		1.1	4	4
9/30/05	17.3	16.8	8.75	0.89	53	62		1.2	4	4
10/14/05	14.6	14.4	7.87	0.37	35	45		1.7	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			C	C		C							C
Chlorophyll a			C	C		C							C
Secchi Depth			C	C		C							B
Overall			C	C		C							C

Source: Metropolitan Council and STORET data



South Twin Lake (82-0019) *Carnelian - Marine Watershed District*

South Twin Lake is a 54-acre lake located within Stillwater Township (Washington County). The maximum and mean depths of the lake are 4.0 m (roughly 13 feet) and 2.0 m (six-and-a-half feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 356 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's 63-acre immediate watershed and surface area translates to a very small watershed-to-lake size ratio of 1.2:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). There is no formal boat access point on the lake.

This was the sixth year that South Twin Lake has been involved in CAMP (although just Secchi transparencies were collected in two of those years). A search through the STORET nationwide water quality database for data on the lake produced a limited amount of data. The only years in which data were found, other than the 2000-2005 CAMP data, was 1996-1999. The years of which included TP, CLA and Secchi transparency data were 1996-2001 and 2004-2005.

The lake was monitored seven times between mid-April and early-October, 2005. During each event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	67.4	37.0	132.0	C
CLA (µg/l)	34.1	7.7	78.0	C
Secchi (m)	1.4	0.6	2.1	C
TKN (mg/l)	1.55	0.94	2.70	
Overall Grade				C

Because of the variability in the lake's water quality database, the determination of any statistically significant long-term trend detection is not possible. In the short-term however, the lake's water quality seems to be well represented by an overall grade of D+/C, with overall grades of F in 1999, D in 1996-1998 and 2001, and C in 2000 and 2004 (the lakes best recorded water quality year) and 2005. To better understand the lake's overall water quality and where it may be heading, continued monitoring is suggested.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 3.4 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.2 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

South Twin Lake Stillwater Twp., Washington Co.



LAKE ID: 820019

WCD: Carnelian-Marine

Volunteer: Wash. Co. SWCD




● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl. a	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/20/05	16.7	14.9	5.6	4.97	8.8	42		2.1	2	4
5/17/05	13.1	12.9	5.98	5.75	7.7	46		2.1	2	3
6/16/05	24.1	22.6	8.43	2.77	16	37		1.8	3	4
7/13/05	23.1	23.1	9.72	4.21	14	37		1.8	3	4
8/12/05	25.2	25	5.18	0.29	95	85		0.6	5	5
9/6/05	25	20.8	12.54	0.47	78	130		0.6	4	5
10/4/05	19.7	17	7.98	1.3	64	69		0.6	4	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				C	C	D	D	C	D			C	C
Chlorophyll a				D	D	D	F	C	D			B	C
Secchi Depth				D	D	F	F	D	F	D	C	C	C
Overall				D	D	D	F	C	D			C	C

Source: Metropolitan Council and STORET data



Spring Lake [Scott County] (70-0054) *Prior Lake - Spring Lake Watershed District*

Spring Lake, located in southeastern Spring Lake Township in Scott County, was monitored 14 times between early-May and mid-October, 2004. The 630-acre lake (5.0 miles in circumference) is considered a “Priority Lake” by the Metropolitan Council because of its multi-recreational uses.

The lake has a large 13,500-acre watershed. The lake and watershed areas translate to a large watershed-to-lake area ratio of 21:1. The larger the ratio, the greater the potential stress on the lake’s quality from surface runoff. The majority of the lake’s watershed is agricultural.

The maximum and mean depths of the lake are 11.3 and 5.6 m (37 and 18 feet), respectively. About 50 percent of the lake’s area is considered littoral (the 0-15 foot depth area dominated by aquatic vegetation). The approximate volume of the lake is approximately 11,500 acre-feet (ac-ft) and a public access to the lake is located on its southwestern shores.

Spring Lake is very fertile, receiving nutrients from runoff and from internal sources. The great fertility causes legendary algal growths. The blue-green algal blooms are a serious nuisance, and purportedly have been the cause of the death of four dogs, which died after drinking the water in 1980.

In an attempt to improve the lake’s water quality, a ferric chloride addition system was constructed at the outlet of the Highway 13 wetland in 1998 with continuous operation starting in 1999. The system, which consists of a dosing station at the outlet of the wetland, followed by a desiltation (settling) basin, meters ferric chloride into stormwater to enhance phosphorus removal prior to entering the lake. The ferric chloride removes nutrients from the water column, thereby reducing their availability to algal growth. As the ferric chloride dosed stormwater enters the desiltation basin the ferric chloride rapidly dissociates to form free iron which reacts with soluble phosphorus to form relatively insoluble iron-phosphorus complex (referred to as floc). The desiltation basin then provides an area where the floc can settle out through the water column and can be eventually removed.

The results from the monitoring of the system in 1999 indicate that there is significant reductions in the ortho-phosphorus load (41 percent) and some reduction in the total phosphorus load (21 percent) from the ditch prior to entering the lake (Prior Lake – Spring Lake Watershed District 2001). The watershed district has continued to monitor the effectiveness of the system

While Spring Lake has been monitored by Metropolitan Council staff in the past, 2005 was the sixth year it has been involved in CAMP (the others being 1997 and 2000-2004). In 2005 the lake was monitored 13 times between early-May and mid-October. On each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the lake’s perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	76.3	40.0	132.0	D
CLA (µg/l)	40.6	5.0	130.0	C
Secchi (m)	1.7	0.5	4.0	C
TKN (mg/l)	1.30	0.72	2.40	
Overall Grade				C

Historical data for the Spring Lake indicates that the water quality of the basin has remained fairly constant over the past decade fluctuating between overall grades of C and D (before the F the lake received this year). Lake quality grades (see the lake's information sheet on the following page) show that when nutrient data were collected on the lake it corresponded to overall grades of C in 1981-1982, 2003 and 2005, and a D in 1980, 1984, 1996-1997, 2000-2001, and 2004, and F in 2002. Because of the fluctuation in the lake's overall grades, no long-term trends are apparent. To better understand all aspects of the lake's water quality and what direction it may be heading, continued monitoring is suggested.

In an attempt to address issues either contributing to the eutrophication of Spring Lake or the symptoms from the resulting eutrophication, the Prior Lake - Spring Lake Watershed District has recently completed a Sustainable Water Quality Management Plan for its lakes (including Spring Lake). The Plan sets goals addressing the lakes' biological and chemical make-up and developed implementation strategies enabling the lakes' goals to be met (PLSLWD 2004).

The physical and recreational conditions of Spring Lake as perceived by the volunteer(s) were ranked on a 1-to-5 scale. These rankings are shown on the lake's information sheet on the next page. The mean summertime physical condition was 3.0 (3- "definite algae present"). The mean suitability for recreation ranking was 1.8 (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.


Spring Lake

Prior Lake/Spring Lake Twp., Scott Co.

Lake ID: 700054

W/D: Prior Lake-Spring Lake

Volunteer: Bill Tisdell



2005 Data





Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	Chl. a µg/L	Surf. TP µg/L	Bot. TP µg/L	Secchi M	PC 1 to 5	RS 1 to 5
5/3/05	10.6				5	69		4	1	1
5/17/05	12.8				17	72		3	1	1
5/19/05	14.8				19	72.5		2.7		
5/31/05	18.5				12	42		2	3	2
6/18/05	26.7				9.7	40		1.3	4	2
7/1/05	24.3				5.7	92		1.2	3	1
7/15/05	32				8.3	40		1.5	3	2
8/1/05	27.8				110	85		0.7	3	1
8/21/05	19				65	77		0.6	4	3
9/4/05	22.4				43	122		0.8	4	
9/16/05	22.8				130	132		0.5	4	3
10/2/05	19.1				66	115		0.4	3	3
10/15/05	15.6				78	160		1	3	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	F	D	D		D								
Chlorophyll a	C	C	C		D						C		
Secchi Depth	C	B	C	C	C	D	D	D	D	C	B	D	C
Overall	D	C	C		D								

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				D	D			F	D	D	D	D	D
Chlorophyll a				C	C			D	D	F	C	D	C
Secchi Depth	C	C	C	D	D			C	D	F	C	D	C
Overall				D	D			D	D	F	C	D	C

Source: Metropolitan Council and STORET data



Square Lake (82-0046) *Marine on St. Croix Watershed Management Organization*

Square Lake, located in eastern May Township (Washington County), is a 193-acre lake (shoreline length of about 2.2 miles) with a maximum and mean depth of 20.7 and 9.0 m (68.0 and 29.5 feet), respectively, for an approximate lake volume of 5,694 ac-ft. About 65 percent of the lake's area is considered littoral (the 0-15 foot depth area dominated by aquatic vegetation). The lake can be accessed through the county park on the southeastern end of the lake. Because of its multi-recreational uses, it is considered a "Priority Lake" in the Metropolitan Area.

The lake is only one of six lakes in the seven-county metropolitan area stocked with trout (rainbows). The lake's level is maintained by a combination of groundwater/ and runoff from the lake's watershed (MDNR 1996).

The lake's watershed is small (about 782 acres) and rural. The watershed and lake size translate to a very small watershed-to-lake size ratio of 4:1 (the smaller the ratio the less the stress on the lake from surface runoff). The watershed is largely undeveloped; wetlands, parks and open spaces, grasslands and woodlands comprise about 70 percent of the watershed's area.

Square Lake, which was involved in CAMP in 1993-1997, and monitored by Council staff in 1998 (as an in-kind contribution to a Clean Water Partnership project on the lake), was a part of CAMP again from 1999-2005. The lake was monitored 14 times from mid-April to mid-October, 2005.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	12.7	8.0	17.0	A
CLA (µg/l)	4.2	2.6	8.5	A
Secchi (m)	5.0	3.4	6.1	A
TKN (mg/l)	0.51	0.42	0.70	
Overall Grade				A

The lake's 2005 overall water quality grade calculated from the TP, CLA, and Secchi grades is identical to those recorded in 1993-2004. That said, a recent trend analysis by the MPCA on the lake's historical (1970-present) Secchi transparency database, revealed a statistically significant decline in recent water clarity.

The water quality graphs show seasonal trends in TP and CLA concentrations, and Secchi transparency for 2005, which closely resemble those of past years. In most metro area lakes, TP, CLA and Secchi transparency generally have a tightly linked relationship such that as TP concentrations increase, algal biomass increases resulting in higher CLA concentrations and lower water clarity. This issue is one that has been addressed as part of the Clean Water Partnership on the lake (Square Lake 2001).

As was mentioned in the previous Council lake reports, the data for Square Lake, shows that the above mentioned relationships are not exclusively dependent on each other. While the graphs show a correlation between CLA and Secchi transparency (clarity increases as CLA decreases and vice versa), TP seemed independent of the other two. An increase or decrease in TP does not automatically result in the same reaction in CLA concentration, which means that phosphorus is not the limiting factor in Square Lake's algal abundance. In fact, earlier Council studies have noted that the lake has lower CLA concentrations than would be expected based on its nutrient levels (Osgood 1981). The reason was discussed in a 1980 Council report and a more recent Clean Water Partnership report on Square Lake which both state that CLA

is limited by the presence of large zooplankton (*Daphnia pulicaria*) which are herbivores that graze on algae and keep the lake's CLA concentrations in check. Therefore, the lake's excellent clarity of Square Lake is due to the presence of *Daphnia* rather than limited by nutrients.

More detailed discussions on the lake, its water chemistry, biological make-up, and hydrologic and nutrient influence the lake's watershed has on the lake can be found in the recent diagnostic-feasibility study completed on the lake as part of a Clean Water Partnership (Square Lake 2001). The complete report highlights the concern of a degrading water clarity trend, the importance of the lake's biological make-up on its overall water quality, the and influence the lake's surface and groundwater watersheds have on the lake's phosphorous load. The Clean Water Partnership report also includes proposed watershed, shoreland, and in-lake projects designed to address issues affecting the lake's quality. An additional resource is an October 2002 report summarizing the lakes recent zooplankton population from monitoring conducted from August 2001-July 2002 (Washington Conservation District 2002)

On each monitoring date, volunteers ranked their opinions of physical and recreational conditions of the lake on a 1-to-5 scale, which are graphed on the lake information sheet. The summertime mean recorded physical condition was 1.9 (between 1- "crystal clear" and 2- "some algae present"). The mean suitability for recreation ranking was 1.3 (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.




Square Lake

May Twp,
Washington Co.

Lake ID: 820046


WMO: Marine-on-St. Croix

Volunteer: Wash. Co. SWCD



2005 Data

Date	Surf Temp	Bot Temp	Surf DO	Bot DO	CIA	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ig/L	ig/L	ig/L	M	1 thru 5	1 thru 5
4/13/05	7.6	4.6	6.27	0.15	3.4	10		7.6	1	1
4/25/05	11.4	5.2	7.01	0.33	3.1	15		6.7	1	1
5/10/05	14.1	5.7	7.03	0.18	4.4	13		5.8	1	1
5/20/05	12.9	5.8	6.58	0.18	8.5	17		4.9	2	1
6/6/05	20.6	6.1	5.25	0.35	3.6	13		4.6	2	2
6/21/05	25.3	6.5	8.44	0.7	2.9	10		5.5	2	1
7/6/05	24	6.5	8.3	0.64	3	14		4.9	2	1
7/19/05	25.7	6.6	8.23	0.11	2.6	13		5.2	2	2
8/1/05	25.9	6.6	6.6	0.66	3.3	12		5.3	2	1
8/15/05	24.7	6.7	7.21	0.65	3	8		6.1	2	2
8/29/05	23.2	6.7	5.55	0.94	5	13		5.2	2	1
9/12/05	23.6	6.8	6.56	0.45	3.9	16		4.4	2	1
9/29/05	18.6	7	7.66	0.81	6	11		3.4	2	1
10/10/05	15.7	6.9	7.69	0.58	4.2	9		5.8	2	1




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	B	A	A	A	A	A				A			
Chlorophyll a	A	A	A	A	A	A				A			
Secchi Depth	A	A	A	A	A	A	A	A	A	A	A		
Overall	A	A	A	A	A	A				A			

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	A	A	A	A	A	A	A	A	A	A	A	A	A
Chlorophyll a	A	A	A	A	A	A	A	A	A	A	A	A	A
Secchi Depth	A	A	A	A	A	A	A	A	A	A	A	A	A
Overall	A	A	A	A	A	A	A	A	A	A	A	A	A

Source: Metropolitan Council and STORET data



Staples Lake (82-0028) *Carnelian - Marine Watershed District*

Staples Lake is a 24-acre lake located within May Township (Washington County). The maximum and mean depths of the lake are 4.3 m (roughly 14 feet) and 2.1 m (seven feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 165 ac-ft. Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

The lake's 127-acre watershed and surface area translates to a watershed-to-lake size ratio of 5.3:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff). There is no formal boat access point on the lake.

This was the sixth year that Staples Lake has been involved in CAMP (although just Secchi transparencies were collected in two of those years). A search through the STORET nationwide water quality database for data on the lake produced a limited amount of data. The only years in which data were found, other than the 2000-2005 CAMP data, was 1997-1999. The years of which included TP, CLA and Secchi transparency data were 1997-2001 and 2004-2005.

The lake was monitored seven times between mid-April and early-October, 2005. During each event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	21.8	18.0	26.0	A
CLA (µg/l)	4.4	2.9	5.8	A
Secchi (m)	2.96	2.44	3.35	B
TKN (mg/l)	0.65	0.54	0.76	
Overall Grade				B

The lake's 2005 overall water quality grade of A, is its best to date. Previous years of monitoring resulted in overall grades of B.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's quality seems well represented by an overall grade of B. To better understand the lake's overall water quality and where it may be heading, continued monitoring is suggested.




The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.0 for physical condition (between 2- "some algae present"), and 3.4 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Staples Lake May Twp., Washington Co.


LAKE ID: 820028
WD: Carnelian-Marine
Volunteer: Wash. Co. SWCD

● Sampling site
Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/20/05	16.4	11.8	5.14	6.02	6.2	18		2.4	2	3
5/19/05	13.8	13.6	5.11	2.33	5.7	20		3.4	2	4
6/15/05	22.9	17.7	5.3	0.3	5.8	25		3.0	2	4
7/13/05	23.5	23.5	7.09	0.53	3.4	26		3.0	2	3
8/11/05	25.1	24.9	3.67	1.99	4.3	20		2.4	2	3
9/8/05	22.3	20.6	7.06	1.3	2.9	18		2.9	2	3
10/6/05	15.8	16	6.23	5.6	2.8	13		3.2	2	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus					B	A	A	C	B			C	A
Chlorophyll a					C	B	B	B	B			A	A
Secchi Depth					B	B	B	B	B	B	C	B	B
Overall					B	B	B	B	B			B	A

Source: Metropolitan Council and STORET data



St. Croix Lake [Whole Lake] (82-0001) *St. Croix Basin Planning Team*

Lake St. Croix is divided into four distinct pools (Bayport Pool, Troy Beach Pool, Black Bass Pool, and Kinnickinnic Pool). There were five monitoring sites amongst the four pools in 2005. The results will be discussed for the entire lake, as well as individually for each of the five sites.

Lake St. Croix (approximately 8,600 acres) is considered by the MNDNR to extend from Stillwater, Minnesota to Prescott, Wisconsin, a distance of approximately 23 miles. The morphometry of each of the pools is shown in the table below.

Lake St. Croix Morphometry

<i>Pool Name</i>	<i>Length (miles)</i>	<i>Area (ac)</i>	<i>Volume (ac-ft)</i>	<i>Mean depth range (dry vs. wet years) (meters)</i>
Bayport Pool	6.0	2,800	62,500	6.2-7.3
Troy Beach Pool	6.0	3,100	107,800	9.9-11.0
Black Bass Pool	7.0	1,300	59,600	12.9-14.0
Kinnickinnic Pool	5.0	1,400	46,274	9.2-10.3

(USGS 2002)

This marks the first year in which any of the Lake St. Croix sites have been formally involved in CAMP. A citizen-monitoring program conducted by the St. Croix Basin Team produced water quality data for four sites (Bayport Pool- Site 2; Troy Beach Pool-Site 3; Troy Beach Pool-Site 5; and Black Bass Pool-Site 6) during the 1999-2002 period, and for one site (Kinnickinnic Pool-Site 7) during the 2000-2001 period. All data are available in STORET.

As part of this report, the lake will first be discussed as a combined “whole” lake system, and then will be followed by sections on each of the five sites individually.

On each sampling event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake’s perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	41.0	17.0	103.0	C
CLA (µg/l)	19.7	<1.0	44.0	B
Secchi (m)	1.5	0.9	2.3	C
TKN (mg/l)	0.81	0.40	2.30	
Overall Grade				C

The whole lake’s 2005 overall grade of “C” (the lake grading system is detailed on page 9 of this report), is identical to those recorded in 1999-2001. That said, the individual parameter means indicate that 2005 was the lake’s best water quality year since the inception of the volunteer monitoring program. The whole lake’s TP and Secchi means of 41.0 µg/l and 1.5 m respectively, are the best recorded to date. The 2005 whole lake CLA mean of 19.7 µg/l is the second best whole lake CLA mean recorded. Because of the limited nature of the lake’s database however, it is not possible to determine any long-term or short-term trends. To better understand the lake’s water quality and where it may be heading, additional years of data collection are needed.

Lake water quality grades based on the whole lakes summer means

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	D	C	C			C
Chlorophyll <i>a</i>							B	C	C	C			B
Secchi Depth							C	C	C	C			C
Overall							C	C	C	C			C

Source: Metropolitan Council and STORET data

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.0 for physical condition (2- "some algae present"), and 2.0 for recreational suitability (2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

St. Croix Lake [Bayport Pool-Site 2] (82-0001) *St. Croix Basin Planning Team*

Lake St. Croix [Bayport Pool-Site 2] was monitored 10 times between early-May and late-September, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	50.0	33.0	72.0	C
CLA (µg/l)	22.0	9.5	44.0	C
Secchi (m)	1.3	0.9	1.6	C
TKN (mg/l)	0.79	0.62	1.00	
Overall Grade				C

The site's 2005 overall grade (C), is identical to those recorded in 1999-2001, and better than the D recorded in 2002. Because of the limited nature of the sites database however, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.5 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 1.8 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.



If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Lake St. Croix, Bayport Pool, Site 2 Minnesota/Wisconsin

Lake ID: 820001-02


Volunteers: Jim and Roberta Harper

- Sampling site
- Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
5/6/05	13.9				21	38		1.4	1	1
5/23/05	16.6				14	39		1.4		
6/5/05	23.7				9.5	45		1.4		1
6/20/05	27.4				13	64		1.4	2	2
7/1/05	25.2				14	72		1.1		
7/17/05	26.7							1.0		
8/2/05	28.7				35	65		0.9	3	
8/12/05	26.3				25	40		1.2	3	2
8/29/05	25.3				21	33		1.6	3	2
9/25/05	20.1				44	52		1.1	3	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							C	D	D	D			C
Chlorophyll a							B	C	C	C			C
Secchi Depth							C	C	C	D			C
Overall							C	C	C	D			C

Source: Metropolitan Council and STORET data



St. Croix Lake [Troy Beach Pool-Site 3] (82-0001) *St. Croix Basin Planning Team*

Lake St. Croix [Troy Beach Pool-Site 3] was monitored 11 times between late-May and late-September, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	45.0	30.0	66.0	C
CLA (µg/l)	17.0	6.8	32.0	B
Secchi (m)	1.4	1.1	1.8	C
TKN (mg/l)	0.68	0.49	0.87	
<i>Overall Grade</i>				C

The site's 2005 overall grade (C), is identical to those recorded in 1999-2001, and better than the D recorded in 2002.

Because of the limited nature of the site's database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 1.7 for physical condition (between 1- "crystal clear" and 2- "some algae present"), and 1.7 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.



If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Lake St. Croix, Troy Beach Pool, Site 3 Minnesota/Wisconsin

Lake ID: 820001-03


Volunteers: Cecilia and Harry Martin

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
5/28/05	17.4				17	40		1.3	3	2
6/7/05	21.9				13	50		1.5	2	2
6/19/05	22.2				7.1	51		1.2	2	2
7/3/05	23.3				6.9	66		1.2	1	2
7/17/05	28.3				6.8	36		1.6	2	2
7/31/05	24.4				28	46		1.2	2	2
8/14/05	26.7				18	38		1.1	2	2
8/17/05	25.0				15	42		1.6	2	1
8/28/05	24.4				16	30		1.8	1	2
9/11/05	23.3				24	43		1.3	1	1
9/26/05	20.0				32	50		1.2	1	1




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	D	D	D			C
Chlorophyll a							B	C	C	C			B
Secchi Depth							D	C	C	D			C
Overall							C	C	C	D			C

Source: Metropolitan Council and STORET data



St. Croix Lake [Troy Beach Pool-Site 5] (82-0001) *St. Croix Basin Planning Team*

Lake St. Croix [Troy Beach Pool-Site 5] was monitored eight times between late-May and late-September, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	33.0	17.0	61.0	C
CLA (µg/l)	23.0	6.9	44.0	C
Secchi (m)	1.6	1.0	2.2	C
TKN (mg/l)	0.69	0.40	1.10	
Overall Grade				C

The lake's 2005 overall grade (C), is identical to those recorded in 1999-2002.

Because of the limited nature of the site's database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 1.4 for physical condition (between 1- "crystal clear" and 2- "some algae present"), and 1.5 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Lake St. Croix, Troy Beach Pool, Site 5 Minnesota/Wisconsin

Lake ID: 820001-05

Volunteers: Richard and Sheryl Lindholm




● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/22/05	14.6				17	22		1.7	1	1
6/19/05	23.0				14	53		1.6	1	1
7/3/05	23.6				6.9	61		1.8	1	2
7/17/05	29.5				35	35		1.0	3	2
7/31/05	26.4				48	35		1.2	2	2
8/21/05	24.3				19	17			1	1
8/28/05	23.7				21	21		1.6	1	1
9/25/05	20.7				23	23		2.2	1	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D	D	C	C			C
Chlorophyll a							B	C	C	C			C
Secchi Depth							C	C	C	C			C
Overall							C	C	C	C			C

Source : Metropolitan Council and STORET data



St. Croix Lake [Black Bass Pool-Site 6] (82-0001) *St. Croix Basin Planning Team*

Lake St. Croix [Black Bass Pool-Site 6] was monitored 11 times between early-May and late-September, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	46.0	27.0	103.0	C
CLA (µg/l)	18.0	<1.0	41.0	B
Secchi (m)	1.6	1.0	2.0	C
TKN (mg/l)	1.10	0.64	2.30	
Overall Grade				C

The lake's 2005 overall grade (C), is identical to those recorded in 1999-2002.

Because of the limited nature of the site's database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.3 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 2.0 for recreational suitability (2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us




Lake St. Croix, Black Bass Pool, Site 6 Minnesota/Wisconsin

Lake ID: 820001-06

Volunteer: Rick Meierotto


● Sampling site

Contours in meters



2005 Data

Date	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	Chl a	Surf. TP	Bot. TP	Secchi	PC	RS
	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1-5	1-5
5/6/05					16	48		1.4	2	2
5/22/05	14.3				8.1	35		1.6	2	2
6/5/05	20.3				5.2	28		1.6		2
6/20/05	23.4				6.6	66		1.9	2	2
7/3/05	23.8				6.7	54		1.8	2	2
7/17/05	29.3				47	103		1.0	2.5	2
7/31/05	27.1				40	39		1.2	3	2.5
8/14/05	25.8				24	27		2.0	2	2
8/27/05	24.4				14	27		1.7	2.5	2
9/11/05	23.5				14	30		1.8	3	2
9/25/05	20.8				16	34		2.0	2	1




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								C	C	C	C		C
Chlorophyll a								B	C	B	C		B
Secchi Depth								C	C	C	C		C
Overall								C	C	C	C		C

Source: Metropolitan Council and STORET data



St. Croix Lake [Kinnickinnic Pool-Site 7] (82-0001) *St. Croix Basin Planning Team*

Lake St. Croix [Kinnickinnic Pool-Site 7] was monitored eight times between late-May and late-September, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	27.0	15.0	40.0	B
CLA (µg/l)	20.0	5.2	33.0	B
Secchi (m)	1.5	1.1	2.3	C
TKN (mg/l)	0.80	0.42	1.30	
Overall Grade				B

The lake's 2005 overall grade of B, is better than the C recorded in 2000.

Because of the limited nature of the site's database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.5 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 2.4 for recreational suitability (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Minnesota/Wisconsin

Lake ID: 820001-07

Volunteer: Carpenter Nature Center





● Sampling site
Contours in meters



2005 Data

	Surf Temp	Bot Temp	Surf DO	Bot DO	Chl a	Surf TP	Bot TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1000S	1000S
5/23/05	15.0				20	40		15	1	1
6/6/05	23.2				5.2	23		23	2	2
7/18/05	28.6				25	36		14	3	3
8/1/05	30.5				24	24		1.1	3	3
8/15/05	27.9				33	26		1.2	3	3
8/29/05	26.2				25	24		1.5	3	3
9/12/05	23.4				17	15		2.0	2	2
9/21/05	21.8				11	31		1.2	3	2.5

Lake Water Quality Grades Based on Summer Averages

[illegible]

Source: Metropolitan Council and STORRET data

St. Joe Lake (10-0011) *City of Chanhassen*

St. Joe Lake is a 14-acre lake located within the City of Chanhassen (Carver County), with a maximum depth of 15.9 m (roughly 52 feet). There is very little other known morphological data available for the lake.

This marks the second year in which St. Joe Lake has been involved in CAMP. A search through the STORET nationwide water quality database for historic data on the lake was provided only two years of Secchi transparency data (1994 and 1996). Nutrient data are only available for 2004-2005 are the only years of nutrient data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored eight times between mid-May and late-September, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	17.3	13.0	22.0	A
CLA (µg/l)	7.4	4.2	18.0	A
Secchi (m)	3.1	1.6	4.2	A
TKN (mg/l)	0.58	0.42	0.94	
Overall Grade				A

As mentioned earlier, there is very little water quality data available for other than the two years of mid-1990's Secchi data and the 2004-2005 CAMP data (both received an overall grade of A). Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.1 for physical condition (between 3- "definite algae present" and 4- "no swimming – boating ok"), and 1.9 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

St Joe's Lake Chanhassen, Carver Co.

LAKE ID: 100011


W/D: Minnehaha Creek

Volunteers: Sue Morgan and

Linda Scott

◆ Sampling site

Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
5/22/05	15				4.5	15		4.2	2	1
5/25/05	17.5				6.2	16		4.0		
6/6/05	23				5.1	16		4.2	2	1
7/10/05	25.7				6	19		2.3	3	1
7/23/05	27.5				4.2	18		2.7	4	3
8/1/05	25.3				6.7	13		3.0	4	2
8/21/05	23.3				7.6	19		2.8	3	1
9/25/05	19.8				18	22		1.6	4	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												A	A
Chlorophyll a												A	A
Secchi Depth		C		B								B	A
Overall												A	A

Source: Metropolitan Council and STORET data



Sunfish Lake (82-0107) Valley Branch Watershed District

Sunfish Lake is a 50-acre lake located within Lake Elmo (Washington County). The lake has a 526-acre immediate drainage area, which results in a watershed-to-lake area ratio of approximately 11:1. The greater the ratio, the greater the potential stress on the lake from surface runoff. There is little else morphological data available for the lake.

This marks the second year in which Sunfish Lake has been involved in CAMP (2000 being the other). A search through the STORET nationwide water quality database for historic data on the lake produced only the forementioned CAMP data. Therefore, 2004-2005 are the only known years of available data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The lake was monitored 13 times between early-May and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	59.3	40.0	97.0	C
CLA (µg/l)	35.3	6.1	74.0	C
Secchi (m)	0.68	0.40	0.90	F
TKN (mg/l)	1.59	1.20	2.30	
<i>Overall Grade</i>				D

As mentioned earlier, there are no water quality data available for Sunfish Lake other than the 2000-2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.


Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.0 for physical condition (2- "some algae present"), and 3.6 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Sunfish Lake Lake Elmo, Washington Co.

Lake ID: 820107
WD: Valley Branch
Volunteer: Tom Hilpish

● Sampling site
Contours in meters



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/3/05	8.9				49	71		0.7		
5/18/05	15.2				35	52		0.8		
5/19/05	14.5				30	63		0.7		4
6/6/05	25				17	40		0.8	2	
6/19/05	25.2				6.1	42		0.7	1	3
7/3/05	24				23	58		0.9	2	
7/21/05	30.2				15	41		0.8	2	3
8/7/05	25.8				23	47		0.8	3	4
8/20/05	24.4				35	55		0.6	2	
9/4/05	21.8				57	65		0.5	2	4
9/18/05	22				59	97		0.5	2	
9/25/05	19.3				74	80		0.4	2	
10/9/05	13.7				40	72		0.4	3	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								C					C
Chlorophyll a								C					C
Secchi Depth								D					F
Overall								C					D

Source: Metropolitan Council and STORM data



Sunnybrook Lake (82-0133) Valley Branch Watershed District

Sunnybrook Lake is a 16-acre lake located within Grant Township (Washington County). The maximum and mean depths of the lake are 6.1 and 2.0 m (20.0 and 6.5 feet), respectively, and the approximate volume of the lake is 104 ac-ft. The majority of the lake's area is considered littoral zone (the area of aquatic vegetation dominance). The lake has a 630-acre immediate watershed, which translates to a watershed-to-lake area ratio of 39:1 (the larger the ratio the greater the potential stress put on the lake from surface runoff).

This was the sixth year in which Sunnybrook Lake has been involved in CAMP (1999 and 2001-2004 being the others). The lake was monitored 16 times between mid-April and mid-October, 2005. Other than for the 1999 and 2001-2004 CAMP data, a search through the STORET nationwide water quality database for data on the lake came up empty. Thus, 1999 and 2001-2005 is the only year of available data.

During each monitoring event, the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. Results are presented on graphs and data tables on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	32.2	19.0	63.0	C
CLA (µg/l)	11.5	3.1	29.0	B
Secchi (m)	2.3	0.9	3.9	B
TKN (mg/l)	1.00	0.80	1.50	
Overall Grade				B

The lake's 2005 overall lake quality grade is identical to those recorded in 2001-2004, and better than the C in 1999.

As mentioned earlier, there are no water quality data available for Sunnybrook Lake other than the 1999 and 2001-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term however, the lakes water quality seems well represented by an overall grade of A. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

The average user perception rankings, on a 1-to-5 scale, were 2.1 for physical condition (roughly equal to 2- "some algae present"), and 1.8 for recreational suitability (between 1- "beautiful" and 2- "minor aesthetic problem").




If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Sunnybrook Lake

Grant, Washington Co.


Lake ID: 820133
W/D: Valley Branch
Volunteer: Arnie Johnson

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/17/05	15.4				8.2	34		1.6	1	1
5/1/05	10.3				3.5	26		2.6	1	1
5/15/05	12.6				3.1	26		3.5	1	1
5/18/05	14.5				5.7	21		3.3		
5/29/05	19.4				4.8	32		3.9	3	2
6/12/05	25.6				9.8	28		2.7	2	2
6/26/05	25.9				5.5	19		2.7	2	2
7/9/05	27.6				3.8	19		2.8	2	1
7/24/05	28.4				5.7	26		2.2	2	2
7/25/05	28				6.2	29		2.1		
8/6/05	28.2				6.3	33		1.9	2	2
8/20/05	23.6				21	58		1.6	3	2
9/4/05	21.7				29	63		1.5	2	2
9/15/05	21.5				19	34		1.3	3	2
9/30/05	16.6				26	37		0.9	2	2
10/11/05	13.3				25	50		1.1	2	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							C		B	B	C	B	C
Chlorophyll a							B		A	A	A	A	B
Secchi Depth							C		B	B	C	B	B
Overall							C		B	B	B	B	B

Source: Metropolitan Council and STORET data



Sunset Lake (82-0153) Rice Creek Watershed District

Sunset Lake, with a surface area of about 124 acres (2.3 miles in circumference), is located in the southern portion of the City of Hugo (Washington County). The lake is considered a “Priority Lake” by the Metropolitan Council due to its multi-recreational uses. One problem that may possibly hinder future recreational activity on the lake, however, is Eurasian Water Milfoil (*Myriophyllum spicatum*), which has been reported in the lake. Its deepest point is approximately 5.2 m (17 feet).

Sunset Lake has been involved in CAMP since 1993. The lake was monitored 17 times from mid-April to mid-October, 2005. The data and resulting graphs showing seasonal variability in TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake information sheet.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	15.1	9.0	24.0	A
CLA (µg/l)	3.7	1.9	6.3	A
Secchi (m)	3.6	2.5	4.9	A
TKN (mg/l)	0.52	0.34	0.72	
Overall Grade				A

When comparing the 2005 overall grade to those of previously monitored years it becomes apparent that the lake’s 2001-2005 overall water quality grade (A) were the best monitored years to date (compared to B’s in 1994 and 2000, and C’s in 1993 and 1995-1999).

Besides the lake’s CAMP data, Secchi transparencies had been measured throughout the mid- and late-1980’s as part of the MPCA’s volunteer program. The lake’s historic individual parameter and overall water quality grades (shown on the following information sheet) indicate that the lake’s water quality has fluctuated over the years. Because of the range in the lake’s quality, a baseline quality for the lake as well as an overall water quality trend is difficult to determine. With this in mind, however, a primitive interpretation of the data seems to show that recently the lake has maintained an “A” grade average (with normal fluctuations). In fact, a recent trend analysis on the lake’s Secchi transparency data by the MPCA, revealed a statistically significant improvement in recent water clarity.

The average user perception rankings on a 1-to-5 scale were 2.0 for physical condition (2- “some algae present”), and 2.7 for recreational suitability (between 2- “minor aesthetic problem” and 3- “swimming slightly impaired”).


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of errors in the lake’s data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Sunset Lake Hugo, Washington Co.

Lake ID: 820153
WD: Rice Creek
Volunteers:
Diane and Bob Coderre

● Sampling site
Contours in meters



2005 Data





Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	CLA ug/L	Surf. TP ug/L	Bot. TP ug/L	Secchi M	PC 1 thru 5	RS 1 thru 5
4/17/05	15				4.1	17		4.0	1	1
4/24/05	13				3.5	14		4.0	2	1
5/6/05	15.2				6.3	11		4.0	2	2
5/23/05	17				2.5	11		4.0	2	2
5/30/05	19				2.7	9		4.0	2	2
6/2/05	21				4.5	14		4.2		
6/8/05	23.3				2.1	17		4.8	2	
6/13/05	25				1.9	11		4.9	2	3
7/1/05	24.4				3.1	16		3.0	2	3
7/18/05	28				3.2	15		3.0	2	3
8/1/05	29.1				5	16		3.6	2	3
8/15/05	25.2				4.9	24		2.5	2	3
8/29/05	24.5				5.6	17		2.6	2	3
9/19/05	22				3.6	20		3.3	2	3
10/1/05	17.9				4	16		4.2	1	4
10/10/05	18				4.7	14		4.0	1	1
10/16/05	9				5.2	19		3.2	1	3

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus					D								
Chlorophyll a					C								
Secchi Depth					C	D	C	D	D	C	C		
Overall					C								

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C	B	C	C	C	C	C	B	A	A	A	A	A
Chlorophyll a	B	B	B	C	C	B	B	A	A	A	A	A	A
Secchi Depth	C	B	C	B	C	C	C	B	A	A	A	A	A
Overall	C	B	C	C	C	C	C	B	A	A	A	A	A

Source: Metropolitan Council and STORET data



Swede Lake (10-0095) Carver County Environmental Services

Swede Lake is a 376-acre lake located in Watertown Township (Carver County) with a maximum depth of approximately 4.0 m (13.1 feet). Because of the shallowness of the lake, its entire surface area is considered littoral (the shallow [0-15 foot depth] area dominated by aquatic vegetation).

The year 2005 marks the fourth year that Swede Lake has been involved in CAMP (2002 being the first). Additionally, Metropolitan Council staff has monitored the lake in 1996 and 2001. The 1996, and 2001-2005 data are the only water quality data found for the lake.

On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The lake was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	254.7	125.0	518.0	F
CLA (µg/l)	60.2	7.3	190.0	D
Secchi (m)	1.0	0.2	1.8	D
TKN (mg/l)	3.85	2.70	6.00	
Overall Grade				D

The lake's 2005 overall grade is similar to that of 2001-2002 and better than 1996 and 2003 (F). A review of past and present parameter means, revealed that 2003 represented the lake's worst monitored water quality to date and 2000 the best monitored water quality year.

As mentioned earlier, there is a limited amount of water quality data available for Swede Lake. Therefore it is not possible to determine any long-term trend. In the short-term however, the lake's quality seems well represented by an overall grade of D/F. To better understand the lake's water quality and where it may be heading, continued monitoring is suggested.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.5 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 4.0 for recreational suitability (4- "no swimming - boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.





If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Swede Lake

Watertown Twp., Carver Co.


Lake ID: 100095
WMO: Carver County
Volunteer:
Wayne Hubin

◆ Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	13.9				4.1	740		2.0	2	4
5/1/05	13.3				7.3	178		1.8	2	4
5/15/05	13.3				29	149		1.5	2	4
5/29/05	21				8	125		1.7	2	4
6/15/05	22				18	224		1.6	2	4
6/29/05	25				190	263		0.8	3	4
7/13/05	27				28	143		1.1	3	4
7/24/05	26				17	283		0.5	2	4
8/7/05	27				51	343		0.4	3	4
8/21/05	22				64	278		0.4	3	4
9/4/05	22.5				100	298		0.4	3	4
9/18/05	21				150	518		0.2		
10/2/05	16				180	527		0.2	3	4
10/15/05	15				110	329		0.3	3	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				D					D	F	F	F	F
Chlorophyll a				F					D	C	F	D	D
Secchi Depth				F					D	C	F	F	D
Overall				F					D	D	F	F	D

Source: Metropolitan Council and STORET data



Sweeney Lake (27-0035) Bassett Creek Watershed Management Organization

The 66-acre lake has a mean and maximum depth of 3.6 m (11.8 feet) and 8.0 m (26.0 feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 790 ac-ft. Because of the shallowness of the lake, and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column) it is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation),

This was the sixth year of CAMP monitoring in Sweeney Lake, which is located in the City of Golden Valley (Henepin County). The lake has two separate depressions each reaching a maximum depth of approximately 8 meters (26 feet). Roughly 52 percent of the lake's area is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation). Additionally, the lake's surface area and 2,400-acre watershed translates to a rather large 36:1 watershed-to-lake size ratio. The greater the ratio, the greater the potential stress on the lake from surface runoff.

The Sweeney Lake branch of the Bassett Creek flows into the lake on the south and outlets at the north over a dam. Sweeney Lake is connected to Twin Lake during periods of high lake levels by a meandering channel through a cattail marsh between the northeast shore of Sweeney and the north shore of Twin Lake. The surface elevations of the two lakes are about the same, indicating a minimal flow between the two lakes except during periods of heavy runoff when transfer of water between the two lakes increases. The west and south shoreline of Sweeney Lake consists of privately owned single family homes. The east shore is bordered by the Glenwood Hills Hospital and park consisting of a lawn, a golf course, and a wooded area (Barr, 1994).

While the lake has been monitored at two separate sites (north end and south end) in the past, only one site (the southern site) was monitored in 2005. The lake was monitored 11 times between mid-April and mid-October, 2005. Results are presented on graphs and data tables on the following page. During each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	49.5	30.0	81.0	C
CLA (µg/l)	21.2	3.3	60.0	C
Secchi (m)	1.5	1.1	2.0	C
TKN (mg/l)	1.08	0.70	1.50	
<i>Overall Grade</i>				C

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's quality seems well represented by an overall grade of C (recorded in 2000-2005). To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. The 2004 mean perceived physical condition of the lake was 1.4 (between 1- "crystal clear" and 2- "some algae present"), while the mean recreational suitability was 1.0 (1- "beautiful").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Sweeney (Twin) Lake Site 1 Golden Valley, Hennepin Co.

Lake ID: 270035
WMO: Bassett Creek
Volunteer: Dave Hanson

• Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	11.9		11.9		24	52		1.2	1	1
5/3/05	11.5		12.9		18	39		1.3	1	1
5/22/05	15.6		9.9		3.4	45		2	1	1
6/5/05	20		9.1		3.3			1.8	1	1
6/21/05	23.7		5.6		12	49		1.5	2	1
7/4/05	24.2		7.8		33	54		1.4	2	1
7/21/05	29.5		8.1		10	30		1.9	1	1
8/6/05	26.7		6.1		28	54		1.2	1	1
8/26/05	24.3		6.7		23	44		1.1	2	1
9/16/05	22.4		6.4		60	81		1.4	1	1
10/13/05	15.2		8.15		55	41		1.2	1	1





Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000 Site 1	2000 Site 2	2001 Site 1	2001 Site 2	2002 Site 1	2002 Site 2
Total Phosphorus								C	C	C	C	C	
Chlorophyll a								C	C	B	C	B	
Secchi Depth								D	D	C	C	C	
Overall								C	C	C	C	C	

Year	2003 Site 1	2003 Site 2	2004 Site 1	2004 Site 2	2005 Site 1	2005 Site 2
Total Phosphorus	C		C		C	
Chlorophyll a	B		B		C	
Secchi Depth	C		C		C	
Overall	C		C		C	

Source: Metropolitan Council and STORET data



Tamarack Lake (10-0010) Minnehaha Creek Watershed District

This was the fifth year that Tamarack Lake has been involved in CAMP (the lake was initially enrolled in 2001). While the 24-acre lake has an unexpected maximum depth of roughly 20.0 m (66 feet), the majority of the lake surface area is considered littoral zone (the shallow 0-15 foot area dominated by aquatic plants). A search through the STORET nationwide water quality database for data on the lake provided limited data (just Secchi data in 1985 and Secchi and nutrient data for 2000-2004).

The lake was monitored 11 times from mid-April to mid-October, 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	33.8	19.0	47.0	C
CLA (µg/l)	24.8	5.1	64.0	C
Secchi (m)	2.1	1.1	3.1	C
TKN (mg/l)	1.11	0.68	1.60	
Overall Grade				C

The lake's 2005 overall grade is identical to that of 2000 and 2003-2004, and worse than the overall grade of B recorded in 2001-2002.

As mentioned earlier, there are very limited amounts of water quality data available for Tamarack Lake. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

The last two graphs show seasonal variation in the lake's perceived physical condition and recreational suitability. The average user perception rankings, on a 1-to-5 scale, were 2.5 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 1.3 for recreational suitability (roughly equal to 1- "beautiful").


The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Tamarack Lake Victoria, Carver Co.





Lake ID: 100010
W/D: Minnehaha Creek
Volunteer: Mike Shouldice

◆ Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	13.2				18	99		1.8	2	1
5/30/05	16.8				16	34		3.1	4	3
6/18/05	23.3				25	40		3.1	3	1
7/10/05	25.8				18	39		2.0	3	1
7/24/05	26.5				49	47		1.5	3	1
8/6/05	23.8				64	43		1.1	4	1
8/20/05	22				16	19		1.5	1	1
9/5/05	21				5.3	22		1.9	1	1
9/17/05	20.6				5.1	26		2.4	1	1
10/1/05	16.2				2	32		2.7	1	1
10/15/05	13.4				4.5	23		3.7	3	1



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus								C	B	B	C	D	C
Chlorophyll a								C	A	B	B	C	C
Secchi Depth				A				C	B	C	C	C	C
Overall								C	B	B	C	C	C

Source: Metropolitan Council and STORET data

Terrapin Lake (82-0031) *Marine on St. Croix Watershed Management Organization*

Terrapin Lake is an 86-acre lake located within the May Township (Washington County), with a maximum depth of 4.6 m (roughly 15 feet). Because of the shallowness of the lake, its entire surface area is considered littoral (the shallow [0-15 foot depth] area dominated by aquatic vegetation). There is very little other known morphological data available for the lake.

This marks the second year in which Terrapin Lake has been involved in CAMP (2004 being the first). A search through the STORET nationwide water quality database for historic data on the lake produced only the forementioned CAMP data. Therefore, 2004-2005 are the only complete years of water quality data available for the lake. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored seven times between mid-April and early-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP ($\mu\text{g/l}$)	22.4	18.0	28.0	A
CLA ($\mu\text{g/l}$)	4.0	1.0	5.9	A
Secchi (m)	3.1	2.6	4.1	A
TKN (mg/l)	0.72	0.49	0.95	
<i>Overall Grade</i>				A

The lake's 2005 overall grade of A is identical to that recorded in 2004.

Because of the limitedness of the lake's water quality database, it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteers ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The resulting user perception rankings are shown on the information sheet. The mean physical condition ranking was 2.0 (2- "some algae present"), while the mean recreational suitability ranking was 3.0 (3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MNDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MNDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Terrapin Lake May Twp., Washington Co.

LAKEID: 820031


WMO: Marine

Volunteer: Wash. Co. SWCD

◆ Sampling site




Contours in meters

0 200 400 600 Meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/20/05	15.9	10.8	5.84	3.3	4.2	21		3.7	3	4
5/19/05		13.3	5.34	5.28	4.4	20		4.1	2	4
6/16/05	23.4	20	7.32	0.31	4.6	26		3.4	2	2
7/13/05		22.7	9.96	0.47	4	20		2.7	2	3
8/11/05	25.5	22.2	6.94	7.8	1	18		2.6	2	3
9/1/05	22.2	20.6	7.8	0.68	5.9	28		2.6	2	3
10/10/05	14.6	13.7	6.71	6.34	7.2	14		3.2	2	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												B	A
Chlorophyll a												A	A
Secchi Depth												A	A
Overall												A	A

Source: Metropolitan Council and STORET data



Turtle Lake (82-0036) *Carnelian - Marine Watershed District*

This was the sixth year of CAMP monitoring in Turtle Lake which is located in the May Township (Washington County). A search through the STORET nationwide water quality database revealed a moderate amount of recent data on the lake. Other than for the 2000-2005 CAMP data (only Secchi transparencies collected in 2002 and 2004), data were found for 1991-1999 (just Secchi data) and nutrient data from 1991-1992 and 1996-1999.

The 44-acre lake has a mean and maximum depth of 2.4 m (eight feet) and 1.2 m (four feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 172 ac-ft. Because of the shallowness of the lake, it is entirely considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation), and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The lake does not have a public access and its 699-acre watershed translates to a 16:1 watershed-to-lake size ratio (the greater the ratio, the greater the potential stress on the lake from surface runoff).

The lake's Secchi transparency was monitored seven times from late-April to early-October 2005. Results are presented in both graphs and data tables on the lake's information sheet on the following page.

Water samples to be analyzed for TP, TKN and chlorophyll were not collected for the lake in 2005. Because Secchi transparency was the only data collected there are no nutrient or chlorophyll concentration means to compare to previous years. The lake's 2005 summertime (May through September) mean Secchi transparency was 1.3 m (minimum of 0.6 m and a maximum of 1.8 m). This translates to a grade of C for water clarity. The lake's overall water clarity grade in 2005 is identical to those recorded in 1999-2004.


As mentioned earlier, there is a moderate amount of historic data available for Turtle Lake recent data collected in the 1990's and early-to-mid-2000's. While no "statistically significant" long-term trends were determined through statistical analysis, a glance at the lake's overall grades from 1991-2005 seems to indicate that the lake's water quality has improved. In the short-term, the lake seems well represented by an overall grade of C. To better understand the lake's water quality and where it may be heading, continued monitoring is suggested.

Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 2.6 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 4.4 (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Turtle Lake

May Twp., Washington Co.



2005 Data





	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/21/05	15.6		3.58	3.31				1.5	2	4
5/19/05	13.6	13	5.2	5.22				1.8	2	4
6/14/05	24.3	23.6	9.38	0.32				0.6	2	4
7/13/05			11.34	11.12				1.8	3	5
8/12/05	23.5	23.3	6.44	4.97				1.2	3	5
9/9/05	21.7	21.7	9.64	9				1.1	3	4
10/6/05	14.2	14.3	7.69	7.43				1.2	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus												F	F
Chlorophyll a												F	F
Secchi Depth												F	F
Overall												F	F

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				C	C	C	B	D	C		D		
Chlorophyll a				D	D	D	C	B	B		B		
Secchi Depth	D	C	D	D	D	D	C	C	C	C	C	C	C
Overall				D	D	D	C	C	C		C		

Source: Metropolitan Council and STORET data



Twin Lake [Burnsville] (19-0028) Black Dog Watershed Management Commission

Twin Lake, an 11-acre lake located in the City of Burnsville (Dakota County). Because of the shallowness of the lake, its entire area is considered littoral zone (the area of aquatic vegetation dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). There is very little known morphological data available for the lake. An area of concern and need for future management is the recent detection of Eurasian Water Milfoil (*Myriophyllum spicatum*) in the lake.

This was the sixth year in which Twin Lake has been involved in CAMP (1999 and 2001-2004 being the others [although the lake was only monitored twice in 2004]). As part of the lake's involvement in CAMP in 2005, the lake was monitored 11 times between late-April and late-September. During each sampling event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake's 2005 raw data and resulting graphs are presented on the associated lake information page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	64.5	28.0	186.0	C
CLA (µg/l)	8.9	1.8	26.0	A
Secchi (m)	1.5	1.0	2.4	C
TKN (mg/l)	0.92	0.51	2.20	
Overall Grade				B

In an attempt to reduce algal blooms and improve the lake's water quality, crushed cornmeal was used in 2005 as an in-lake organic carbon amendment. A recent study on Valley Lake-Lakeville, Minnesota (discussed later in Valley Lake section of this report), has suggested that carbon from the decaying barley straw inhibits algal populations via microbial competition for phosphorus (McComas and Anhorn 2004). The use of the cornmeal on Twin Lake did seem to result in the anticipated improvements in the lake's water quality (McComas 2005).

In fact, the lake's Secchi transparency in 2005 would have been greater except on many monitoring events the lake's excessive submergent macrophyte growth got in the way. Therefore, the lake's 2005 water clarity was actually better than that represented by the summer mean.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's overall water quality seems to be well represented by a water quality grade of high-C/low-B

Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 3.5 (between 3- "definite algae present" and 4- "high algal color"), while the mean recreational suitability ranking was 4.0 (4- "no swimming – boating ok").




If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Twin Lake Burnsville, Dakota Co.

Lake ID: 190028


WMO: Black Dog

Volunteer: Bernie DeMaster



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/24/05	14.9				3.1	27		2.6	1	1
5/7/05	15.7				3	36		2.4	2	4
5/30/05	21.1				1.8	35		1.9	5	4
6/21/05	26.7				4.6	34		1.7	4	4
7/5/05	24.3				6.4	130		1.5	5	4
7/18/05	26.3				7.4	34		1.2	5	4
8/1/05	26.5				7.9	55		1.3	3	4
8/16/05	25.2				8.1	38		1.0	3	4
8/30/05	24.3				26	186		1.0	3	4
9/11/05	25.5				9.2	28		1.2	3	4
9/25/05	20.7				15	69		1.5	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus							D		C	C	C		C
Chlorophyll a							B		A	A	A		A
Secchi Depth							D		C	C	C		C
Overall							C		B	B	B		B

Source: Metropolitan Council and STORET data



Twin Lake (Robbinsdale) [Lower Basin] (27-0042) Shingle Creek Watershed Man. Comm.

This was the fourth year that the lower basin of Twin Lake, which is located in the City of Robbinsdale (Hennepin County), was monitored as part of CAMP. The lake has also been monitored by Council-staff in the past. As part of the volunteer monitoring program the lower basin of Twin Lake was sampled eight times from late-April to early-August, 2005.

The entire 212-acre lake has maximum and mean depth of 14.0 and 2.1 m (46 and 7 feet), respectively. The acreage of each basin is as follows: lower basin= 46 acres, middle basin= 69 acres, and the upper basin= 137 acres.

The lower basin itself has a mean and maximum depth of 7.0 m and 1.2 m (23 and 4 feet). The total volume of the whole lake is approximately 1,490 ac-ft (175 ac-ft of which is contained within the lower basin). About 81 percent of the whole lake's area is considered littoral (the 0-15 foot depth area dominated by aquatic vegetation). Access to the lake can be obtained at two locations, the southern end of the lake and the lake's eastern shoreline.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP ($\mu\text{g/l}$)	42.3	29.0	60.0	C
CLA ($\mu\text{g/l}$)	39.0	7.0	160.0	C
Secchi (m)	1.7	0.9	3.3	C
TKN (mg/l)	1.46	0.83	2.20	
Overall Grade				C

The lake's 2005 overall grade of C is identical to those recorded in 1993, 1996, 1998, 2000, and 2003, and better than the D recorded in 1991.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's quality seems well represented by an overall grade of C grade. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 2.2 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.0 (2- "minor aesthetic problem").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Twin Lake, Lower Basin, Robbinsdale, Hennepin Co.


Lake ID: 270042-03

WMO: Shingle Creek

Volunteers: Roni Brunner and Bob Hill

● Sampling site




Contours in meters



Lower
Basin

2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/24/05	14				6.2	40		2.7	1	1
5/15/05	12.6				7	44		2.5	2	1
5/29/05	17.5				9.2	29		3.3	1	1
6/21/05	27.9				25	40		1.8	2	2
7/6/05	27.3				24	49		0.9	4	3
7/20/05	29.5				8.8	35		1	2	3
7/25/05	29				11	60		1.1		
8/3/05	28.3				160	39		1	2	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus												D	
Chlorophyll a												D	
Secchi Depth												D	
Overall												D	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C			C		C		D			C		C
Chlorophyll a	C			C		B		C			B		C
Secchi Depth	D			C		C		C			C		C
Overall	C			C		C		C			C		C

Source: Metropolitan Council and STORET data



Twin Lake (Crystal) [Middle Basin] (27-0042) *Shingle Creek Watershed Man. Comm*

The middle basin itself has a mean and maximum depth of 14.0 m and 4.9 m (46 and 16 feet). The total volume of the whole lake is approximately 1,490 ac-ft (918 ac-ft of which is contained within the middle basin).

Two thousand and five was the fourth year that the lower basin of Twin Lake, which is located in the City of Crystal (Hennepin County), was monitored as part of CAMP. Between late-April and early-August, 2005, the middle basin of Twin Lake was monitored eight times. On each monitoring event the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the lake's perceived physical condition and recreational suitability. The lake was also enrolled in CAMP in 1997, 1999 and 2003.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	57.8	37.0	70.0	C
CLA (µg/l)	14.9	5.3	34.0	B
Secchi (m)	1.8	0.9	3.20	C
TKN (mg/l)	1.53	1.30	1.90	
Overall Grade				C

The lake's 2005 overall grade of C is identical to those recorded in 1996, 1999, and 2000, better than the D recorded in 1991, and worse than the B recorded in 1985 and 1997.

Because of the sporadic and limited nature of lake's database, no statistically significant long-term or short-term trend can be determined. To better understand the quality of the lake and what direction it may be heading, continued monitoring is suggested.

Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 2.4 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.2 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Twin Lake, Middle Basin, Crystal, Hennepin Co.



Lake ID: 270042-02

WMO: Shingle Creek

Volunteers: Roni Brunner and Bob Hill




● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/24/05	13.4				6.9	63		3.1	1	1
5/15/05	13.2				5.3	65		3.2	1	1
5/29/05	17.6				10	57		3.2	1	1
6/21/05	28.2				34	70		1.2	4	2
7/6/05	26.4				18	48		0.9	4	4
7/20/05	30.2				11	70		1.1	2	3
8/3/05	29.2				11	37		1.1	2	2
8/7/05					11	65				




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus						C						C	
Chlorophyll a						B						D	
Secchi Depth						A						D	
Overall						B						D	

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				C	C		C	C					C
Chlorophyll a				C	A		B	C					B
Secchi Depth				C	C		C	C					C
Overall				C	B		C	C					

Source: Metropolitan Council and STORET data



Twin Lake [St. Louis Park] (27-0656) City of St. Louis Park

Twin Lake is a small shallow lake located within City of St. Louis Park (Hennepin County). There is very little known morphological data available for the lake.

This marks the fourth year in which Twin Lake has been involved in CAMP (2002-2004 being the others). A search through the STORET nationwide water quality database for historic data on the lake provided only the forementioned 2002-2004 CAMP data. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 11 times between mid-May and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	160.6	108.0	237.0	F
CLA (µg/l)	44.0	14.0	120.0	C
Secchi (m)	0.6	0.5	0.9	D
TKN (mg/l)	1.41	0.81	2.50	
Overall Grade				D

The lake's 2005 overall grade of D is identical to those recorded in 2002-2004.


As mentioned earlier, there are no water quality data available for Twin Lake other than the 2002-2005 CAMP data. Therefore it is not possible to determine any long-term trend. In the short-term however, the lake's water quality seems well represented by an overall grade of D. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 2.2 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 3.8 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you notice any errors in the lake data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us





Twin Lake St. Louis Park, Hennepin Co.

Lake ID: 270666
W/D: Minnehaha Creek
Volunteer: Ed Voyles



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
5/16/05	17.2				39	148		0.6		
5/21/05	18.7				34	139.5		0.5	1	4
6/5/05	23.1				34	166		0.6	2	4
6/18/05	28.9				18	160		0.5	2	4
7/7/05	26.9				14	146		0.7	2	4
7/24/05	30.2				120	179		0.5	3	4
8/14/05	23.2				64	237		0.5	2	4
8/28/05	24.1				44	183		0.7	4	2
9/11/05	23.6				24	108		0.9	2	4
10/9/05	13.3				6.5	89		1.1	2	4
10/16/05	13.3				22	85		1	2	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus										F	F	F	F
Chlorophyll a										B	C	D	C
Secchi Depth										D	D	D	F
Overall										D	D	D	D

Source: Metropolitan Council and STORET data

Valentine Lake (62-0071) Rice Creek Watershed District

Valentine Lake is located within the City of Arden Hills in Ramsey County. The lake has a surface area of 60-acres, and a maximum and mean depth of 4.0 m (13.1 feet) and 1.5 m (4.9 feet), respectively. Because of the shallowness of the lake, its entire surface area is considered littoral, the shallow (0-15 foot depth) area dominated by aquatic vegetation, and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). The mean depth and surface area of the lake translates to an approximate volume of 300 ac-ft. The result of comparing the lake's surface area to its 2,237-acre drainage area (watershed) is a rather large 37:1 watershed-to-lake size ratio (the greater the ratio, the greater the potential stress on the lake from surface runoff).

This was the fifth year that Valentine Lake has been involved in CAMP (2001-2004 being the others). In fact, the 2001-2004 CAMP data were the only data found through STORET nationwide water quality database search. Therefore 2001-2005 represents the only water quality data readily available for the lake.

The lake was monitored nine times between mid-April and mid-October, 2005. On each sampling day the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	62.7	29.0	117.0	C
CLA (µg/l)	30.3	2.8	82.0	C
Secchi (m)	1.5	0.7	2.8	C
TKN (mg/l)	1.75	1.40	2.20	
<i>Overall Grade</i>				C

While the resulting overall grade for 2005 (C) is identical to those of 2001-2004, the individual grades of 2001-2002 were better. The lake's 2005 nutrient concentrations and Secchi transparencies are graphed on the following page.

Because of the limitedness of the lake's water quality database, the determination of any only long- or short-term trends are not possible to determine. It is reported on the MPCA website, however, that a recently conducted trend analysis on the lake's Secchi transparency data revealed a statistically significant improvement in recent water clarity. To better understand the lake's water quality and what direction it may be heading, more years of data collection are needed.



The perceived physical and recreational conditions of the lake, recorded by the volunteers, were ranked on a 1-to-5 scale. The rankings are shown in both tabular and graphical form on the lake's associated information sheet. The mean physical condition ranking was 1.8 (between 1- "crystal clear" and 2- "some algae present"), while the mean recreational suitability ranking was 2.8 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Valentine Lake Arden Hills, Ramsey Co.




Lake ID: 620071
WD: Rice Creek
Volunteer: Bob Kistler

● Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/17/05	14.7				19	54		0.9	3	2
4/29/05	10.9				9.2	35		1.3	2	1
5/14/05	12.6				5.6	30		1.6	2	1
5/30/05	21				2.8	34		2.8	1	2
6/10/05	24.8				3.2	29		1.6	1	2
8/15/05	25.3				7.3	55		1.4	2	4
9/2/05	21.7				82	117		0.7	2	4
9/27/05	18.6				81	111		0.8	3	4
10/15/05	14.2				20	100		1.2	3	2




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus									C	C	C	C	C
Chlorophyll a									B	B	C	C	C
Secchi Depth									C	C	D	C	C
Overall									C	C	C	C	C

Source: Metropolitan Council and STORM data



Valley Lake (19-0348) City of Lakeville

This was the tenth year that Valley Lake, located in the City of Lakeville (Dakota County), has been involved in CAMP. The lake has been monitored through CAMP in 1995-1997 and 1999-2005. A search through the nationwide water quality database (STORET) found no water quality data on the lake prior to the 1995 CAMP data.

The 15-acre lake has a maximum depth of 3.2 m (10 feet). Because of the shallowness of the lake, the entire lake is considered littoral (the area of dominated by aquatic vegetation). The majority of the land within the lake's 117-acre watershed is parkland or single-family residential homes. The watershed-to-lake size ratio is 8:1 (the greater the ratio, the greater the potential stress on the lake from surface runoff).


The lake has been involved in CAMP from 1999-2004 as part of a barley straw project on the lake (where barley straw has been added to the lake in order to inhibit algal populations). While, barley straw has been used for algal control in the United Kingdom for many years, the controlling mechanism had not truly known. Therefore, the Valley Lake study was trying to determine two questions; 1) the success of the barley straw treatment in Valley Lake; and 2) to clarify the controlling mechanism.

In 2005, TP, TKN, CLA, Secchi transparency as well as total and dissolved carbon were tested 13 times between late-April and mid-October.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	51.0	25.0	88.0	C
CLA (µg/l)	37.6	5.7	86.0	C
Secchi (m)	1.3	0.6	2.2	C
TKN (mg/l)	0.94	0.34	1.70	
Overall Grade				C

The lake's 2005 water quality grades, although identical to those recorded in 2004, were worse than those recorded in 2000-2003 (2002 was the lake's best-recorded water quality year).



While the above graph shows a definite improvement in lake water quality since the use of barley straw first began in 1999, the lake's quality has shown some degradation over the last three years (2003-2005). This is especially evident in 2005.

One explanation for the recent dip in water quality could be directly linked to an escalating panfish population. Similar to that mentioned for Alimagnet and Lee lakes, where recent fish surveys suggested that unusually high fish densities might be negating the benefits of the barley treatment on the lakes. The Alimagnet and Lee lake surveys revealed the lake's fishery being dominated by bluegill sunfish, black crappies, and black bullheads (with as much as 30+ times the areas average number of bluegills per net in Laee Lake) (McComas 2004). The recent removal of roughly 80 pounds of fish per acre on Lee Lake, coinciding with its improved water quality, suggests not only that the barley straw treatment was successful in improving Lee Lake's 2005 water quality, but that the unbalanced fishery may have indeed been negating the benefits of previous barley straw treatments.

For this reason an in-lake fisheries survey was conducted for Valley Lake in 2005. Results of the survey were not available at the printing of this report.

While, prior to the change in the lake's fishery, barley straw seemed to inhibit algal growth, which in turn resulted in improved water clarity, the reason was not truly understood. In an attempt to identify the algal inhibitor released by the decaying barley straw, additional in-lake water samples were taken in 2001-2005 in Valley Lake along with monthly samples in a nearby sediment basin where barley straw was not used (viewed as a control). These additional samples, which in the past had centered around the break down of phenols concentrations (one of the theories behind the barley straw inhibitor) as a part of 57 base neutral acids organic compounds (BNAs), as well as total and dissolved carbon. Because the breakdown of BNA compounds for each of the collected samples in 2002 came back below detection limit ($< 2.0 \mu\text{g/l}$), it is not thought that chemical compounds (such as phenols) released from the decomposing straw is the mechanism inhibiting the algal growth (McComas 2003) (McComas and Anhorn 2004).

The examination of the Valley Lake and sedimentation basin carbon data was the focus of the additional monitoring in 2002-2005. The thought is that the carbon may aid another potential algal inhibiting process, that is the microbial competition for phosphorus (McComas 2003) (McComas and Anhorn 2004). Therefore, the presence of decaying barley straw results in the lake's algal biomass actually being phosphorus-limited not inhibited by a released chemical compound. That is, that the barley straw serves as an organic carbon source for increased heterotrophic production which results in lowering the water column phosphorus and thus reduces algal growth (McComas and Anhorn 2004). Based on the results of the four-year Valley Lake study, heterotrophic production enhanced by barley straw is the most likely algal inhibiting mechanism. For this reason, another organic carbon source, crushed corneal (which is less expensive than barley straw yet has a similar C:P ratio), was used in 2005 as the in-lake organic carbon amendment rather than barley straw.

While initial analysis of the 2002-2004 carbon and chlorophyll data seems to strongly support the organic carbon amendment theory, the 2005 data seems to indicate a change in the inner-lake workings. Could this be the result of a shift in the lake's fishery, similar to Alimagnet and Lee lakes, or are there other reasons that the in-lake cornmeal treatment did not work? The results of the fishery survey and additional monitoring will hopefully shed some light on this question.

In addition to the collection of TP, TKN, CLA and Secchi information during each monitoring event in 2005, the volunteer(s) ranked their perception of the lake's physical and recreational conditions on a 1-to-5 scale as shown on the attached information sheet. The summertime mean recorded physical condition was 2.7 (between 2- "some algae present" and 3- "definite algae present"). The mean suitability for recreation ranking was 2.8 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you know of any errors in the lake's data, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Valley Lake

Lakeville, Dakota Co.


Lake ID: 190348

WMO: Dakota County

Volunteer: City of Lakeville





● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/26/05	10.1				12	45		2.1	2	2
5/9/05	15.7				8.1	40		1.6	2	2
5/24/05	19.6				7.8	25		1.6	2	2
6/6/05					8.9	38		2.2	3	3
6/21/05	25.8				42	57		1.4	3	3
7/5/05	23				68	43		1.1	4	4
7/20/05	26.9				26	53		1.6	2	2
8/2/05	27.4				73	70		0.6	3	3
8/16/05	26.9				86	60		0.7	3	3
8/29/05	26.9				35	47		1.1	3	3
9/13/05	22.4				57	40		0.9	2	3
9/27/05	19.6				53	88		0.9	3	3
10/10/05	15.7				28	21		1.1	3	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			D	D	C			C	C	C	C	C	C
Chlorophyll a			C	C	C		C	B	A	A	B	C	C
Secchi Depth			D	D	D		D	C	C	B	B	C	C
Overall			D	D	C		C	B	B	B	B	C	C

Source: Metropolitan Council and STORET data

Virginia Lake (10-0015) Minnehaha Creek Watershed District

This was the sixth year of CAMP monitoring in Virginia Lake, which is located in the City of Victoria (Carver County). A search through the STORET nationwide water quality database revealed a limited amount of historic data on the lake (1988-1991 [consisting of just a single monitoring event] and 2000-2004). The 110-acre lake has a mean and maximum depth of 3.3 m (11 feet) and 10.4 m (34 feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 1,210 ac-ft. Roughly 88 percent of the lake's area is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation). Additionally, the lake's 450-acre immediate watershed translates to a 7:1 watershed-to-lake size ratio (the greater the ratio, the greater the potential stress on the lake from surface runoff). An area of concern and need for future management is the presence of Eurasian Water Milfoil (*Myriophyllum spicatum*) in the lake.

The lake was monitored 13 times between early-May and early-October, 2005. Results are presented on graphs and data tables on the following page. During each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the perceived physical condition and recreational suitability. The collected data and resulting graphs showing TP and CLA concentrations, Secchi transparency, and user perception (physical condition and recreational suitability) are presented on the lake's information sheet on the following page.

2005 summer (May-September) data summary

Parameter	Mean	Minimum	Maximum	Grade
TP (µg/l)	48.7	36.0	94.0	C
CLA (µg/l)	18.5	4.9	45.0	B
Secchi (m)	1.1	0.5	1.7	D
TKN (mg/l)	1.05	0.70	1.60	
Overall Grade				C

While the lake's 2005 overall grade is identical to those recorded in 2000-2004, the 2005 Secchi mean is its worst recorded to date.

Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 2.8 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.6 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

As mentioned earlier, there is a limited amount of historic data available for Virginia Lake other than the single event monitoring of 1988-1991 (no grades are shown for these data due to the limited number of events) and 2000-2005 CAMP data. Therefore it is not possible to determine any long-term trends. In the short-term however, the lake seems well represented by an overall water quality grade of C. To better understand the lake's water quality and where it may be heading, continued monitoring is suggested.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Valley Lake

Lakeville, Dakota Co.


Lake ID: 190348

WMO: Dakota County

Volunteer: City of Lakeville





● Sampling site

Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/26/05	10.1				12	45		2.1	2	2
5/9/05	15.7				8.1	40		1.6	2	2
5/24/05	19.6				7.8	25		1.6	2	2
6/6/05					8.9	38		2.2	3	3
6/21/05	25.8				42	57		1.4	3	3
7/5/05	23				68	43		1.1	4	4
7/20/05	26.9				26	53		1.6	2	2
8/2/05	27.4				73	70		0.6	3	3
8/16/05	26.9				86	60		0.7	3	3
8/29/05	26.9				35	47		1.1	3	3
9/13/05	22.4				57	40		0.9	2	3
9/27/05	19.6				53	88		0.9	3	3
10/10/05	15.7				28	21		1.1	3	4



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			D	D	C			C	C	C	C	C	C
Chlorophyll a			C	C	C		C	B	A	A	B	C	C
Secchi Depth			D	D	D		D	C	C	B	B	C	C
Overall			D	D	C		C	B	B	B	B	C	C

Source: Metropolitan Council and STORET data

Waconia Lake (10-0059) Carver County Environmental Services

Lake Waconia, located next to the City of Waconia in Carver County, is considered a Metropolitan Area “Priority Lake” because of its multi-recreational uses. Lake Waconia is one of the largest bodies of water in the region. It has a surface area of approximately 3,000 acres (6.8 miles in circumference), and mean and maximum depths of 4.0 and 11.3 m (13.1 and 47.1 feet), respectively. The lake has an approximate volume of 38,632 ac-ft (resulting in a retention time of about 10 years) and an approximate watershed-to-lake size ratio of 4:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

There are a couple of public accesses on the lake to support its high-volume recreational use. One access is a city-owned access on the southern end of the lake while the other is a state-owned access on the northeastern shore. In the future, the recreational use on and around the lake may increase if a proposed regional park on the lake’s southeastern shores becomes a reality. One problem that may possibly hinder future recreational activity on the lake, however, is Eurasian Water Milfoil (*Myriophyllum spicatum*) which has been reported in the lake.

A variety of land uses around Lake Waconia may contribute to the lake’s nutrient load. There are residential areas along the lake’s shoreline (25.9 homes/shoreline mile), wetlands, commercial/ industrial areas, and rural/agricultural uses. The predominant uses associated with rural/agricultural areas are livestock and crop farming (51 percent), while those affiliated with the City of Waconia include: single and multi-family residential, commercial/industrial, and paved areas such as parking lots and city streets. All the land uses around the lake pose a potential runoff and pollution problem to the lake. Shoreline homes provide the possibility of lawn runoff of herbicides and fertilizers. Rural/agricultural uses, if not properly managed, can result in herbicides, fertilizers, and eroded soils ending up in the lake. City uses, where a majority of the area is paved, can result in large amounts of nutrient rich debris entering the lake after a rainstorm. These non-point pollution problems can hasten the lake’s natural eutrophication process, resulting in a lake that cannot support all of its recreational uses.

Lake Waconia has been involved in CAMP since 1994 (and monitored by Council-staff in 2004). In 2005, the lake was monitored 14 times between mid-April and mid-October. During each monitoring event the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake’s perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	28.2	11.0	58.0	B
CLA (µg/l)	16.1	1.0	36.0	B
Secchi (m)	4.5	1.2	7.4	A
TKN (mg/l)	0.83	0.49	1.20	
Overall Grade				B

The lake’s 2005 water clarity mean of 4.5 meters, was the best recorded to date. The lake’s best overall water quality year recorded through CAMP was 1994 (TP= 21.0 µg/l, CLA= 6.3 µg/l, and Secchi= 3.1 m resulting in an overall grade of A). The worst was 1998 (TP= 39.1 µg/l, CLA= 18.2 µg/l, and Secchi= 2.0 m resulting in an overall grade of C).

A search of Council, MPCA, and STORET databases revealed nutrient water quality data for 1980, 1981, 1985, 1990, 1994-2004 (all as a part of the Council’s lake monitoring programs). Additionally, Secchi transparencies have been collected through the MPCA’s Citizen Lake Monitoring Program since 1974. The

lake received C's in 1980,, and B's and C's in 1981, 1985 and 1990, an A in 1994 a strong B in 1995 and 2002, moderate B in 1996-1997, 2001, and 2004-2005, and a C in 1998-2000 and 2003. Supplemental Secchi data from 1980-1993 has resulted in annual grades of C or D. The lake's overall water quality grade seems to be well represent by an overall grade of C+/B. Additionally, the MPCA recently conducted a trend analysis on the lake's Secchi transparency data, which revealed a statistically significant improvement in recent water clarity

The volunteer monitor's perception of the lake's physical and recreational conditions were ranked on a 1-to-5 scale during each monitoring event. The rankings are shown on the information sheet on the next page. The mean physical condition ranking was 1.7 (between 1- "crystal clear" and 2- "some algae present"), while the mean recreational suitability ranking was 1.0 (1- "beautiful").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.


If you know of any errors in the lake's data or physical information, or are aware of any additional information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Lake Waconia

Laketown Twp./Waconia Twp.,
Carver Co.





Lake ID: 100059
WMO: Carver Creek
Volunteer: Carver Co.
Env. Services

● Sampling site
Contours in meters



2005 Data

Date	Surf. Temp C	Bot. Temp C	Surf. DO mg/L	Bot. DO mg/L	Chl. a µg/L	Surf. TP µg/L	Bot. TP µg/L	Secchi M	PC 1-5	RS 1-5
4/15/05	11.9		10.6		2.8	14.5		4.9	2	2
4/27/05	10.6		10.9		3.3	18.5		4.8	1	1
5/11/05	13.5		10.4		3.3	20		6.6	1	1
5/24/05	19.4		8.9		2.5	12		7.4	2	1
6/5/05	19.7		9.1		8.1	17		7	2	1
6/20/05	23		8.82		1	11		7	1	1
7/1/05	23.4		9.33		5.8	18		2.9	1	1
7/18/05	24.9		8.12		9.2	20				
8/1/05	23.7				27	55		1.5	2	1
8/18/05	23.8		8.2		36	49			2	1
8/31/05	22.7				33	37		1.2	2	1
9/13/05	22		7.33		34	37		1.4	2	1
9/23/05	20.3		15.6		17	58		2	2	1
10/10/05	14.2		10.6		10	41		2.4	1	1



Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus	C	B				B							
Chlorophyll a	C	B				B					C		
Secchi Depth	C	C	C	C	D	C	C	C	D	C	C	C	C
Overall	C	B				B							

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus		A	A	B	B	C	C	C	C	B	C	B	B
Chlorophyll a		A	B	B	B	B	B	B	B	B	B	B	B
Secchi Depth	C	A	B	C	C	C	C	C	B	B	C	C	A
Overall		A	B	B	B	C	C	C	B	B	C	B	B

Source: Metropolitan Council and STORET data

West Boot Lake (82-0044) *Carnelian - Marine Watershed District*

This was the fifth year of CAMP monitoring in West Boot Lake which is located in May Township (Washington County). A search through the STORET nationwide water quality database revealed a moderate amount of data on the lake over the past 10+ years (1991 and 1996-1999 and the 2000-2004 CAMP data). The 110-acre lake has a mean and maximum depth of 5.9 m (19 feet) and 11.9 m (39 feet), respectively. The mean depth of the lake and its surface area translate to an approximate lake volume of 2,090 ac-ft. Roughly 56 percent of the lake's area is considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation). The lake's 209-acre immediate watershed translates to a 2:1 watershed-to-lake size ratio (the greater the ratio, the greater the potential stress on the lake from surface runoff).

The lake was monitored 14 times between mid-April and mid-October, 2005. Results are presented on graphs and data tables on the following page. During each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the perceived physical condition and recreational suitability.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	19.5	15.0	25.0	A
CLA (µg/l)	4.2	2.2	9.2	A
Secchi (m)	4.3	2.4	6.4	A
TKN (mg/l)	0.70	0.50	0.97	
Overall Grade				A

The lake's 2005 overall grade is identical to those recorded in 1999-2004.

Throughout the summer, the volunteer ranked the lake's perceived physical condition on a 1-to-5 scale (see lake information sheet). The mean physical condition ranking was 2.2 (between 2- "some algae present" and 3- "definite algae present"), while the mean recreational suitability ranking was 2.2 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

No statistically significant long-term trend is evident from the lake's overall water quality database, in the short-term however, the lake seems to have a wide range of fluctuation (overall grade of B in 1996, C in 1997, B in 1998 and A's in 1999-2004). A recent MPCA conducted trend analysis on the lake's Secchi transparency data, however, revealed a statistically significant improvement in recent water clarity. To better understand the lake's overall water quality and where it may be heading, continued monitoring is suggested.

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.




West Boot Lake

May Twp., Washington Co.

LAKE ID: 820044


W/D: Camelian-Marine

Volunteer: Wash. Co. SWCD



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
4/14/05	12.8	4.3	5.87	0.3	2.7	24		5.0	2	2
4/25/05	13	4.4	5.36	0.04	2	28		7.6	2	2
5/10/05	16	4.5	6.02	0.14	9.2	24		3.2	2	3
5/24/05	19.8	4.7	5	0.03	8.1	15		2.4	3	3
6/6/05	22.6	4.7	3.8	0.35	2.7	19		6.4	2	2
6/22/05	26.9	4.8	7.96	0.57	3	22		5.0	2	1
7/8/05	24.7	4.9	8.02	0.29	4	19		3.8	2	2
7/19/05	27.6	4.9	6.1	0.61				3.7	2	2
8/1/05	28.8	5.1	6.21	0.77	2.2	16		4.3	2	2
8/15/05	25.2	5.2	6	0.51	3.5	15		3.5	3	3
8/29/05	23.6	5.3	4.72	0.74	2.7	19		5.8	2	2
9/13/05	23.3	5.5	7	0.54	3.4	25		4.4	2	2
9/29/05	17.5	6	6.47	0.82	3.5	21		4.3	2	2
10/10/05	14.5	5.8	7.26	0.54	3.1	7		4.0	2	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				B	C	A	A	A	A	A	B	A	A
Chlorophyll a				A	B	C	A	A	A	A	A	A	A
Secchi Depth				B	C	B	A	A	A	A	A	A	A
Overall				B	C	B	A	A	A	A	A	A	A

Source: Metropolitan Council and STORET data



Westwood Lake (27-0711) Bassett Creek Watershed Management Organization

This was the seventh year of CAMP monitoring in Westwood Lake (1993 and 2000-2004 being the others), which is located in the City of St. Louis Park (Washington County). The 41-acre lake has a maximum depth of 2.0 m (six-and-a half feet). Because of the shallowness of the lake, it is entirely considered littoral zone (the 0-15 foot depth area dominated by aquatic vegetation), and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column).

Westwood Lake was monitored seven times between mid-May and late-September, 2005. Results from the monitoring are presented on the information sheet on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	100.0	28.0	321.0	D
CLA (µg/l)	44.3	9.7	100.0	C
Secchi (m)	1.4	1.1	1.7	C
TKN (mg/l)	1.75	0.94	3.0	
Overall Grade				C


Because there is a limited amount of historic data available for Westwood Lake, it is not possible to determine any long-term trends. In the short-term however, the lake's water quality shows a wide range of fluctuation (overall grade of D in 1982, C in 1993, 2001-2002, and 2005, and B in 2000 and 2003-2004). To better understand the lake's water quality and where it may be heading, continued monitoring is suggested.

Throughout the monitoring period, the volunteers' opinion of the lake's physical and recreational conditions were ranked on a 1-to-5 scale. These user perception rankings are shown on the lake information sheet. The average user perception rankings, on a 1-to-5 scale, were 2.8 for physical condition (between 2- "some algae present" and 3- "definite algae present"), and 3.6 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming – boating ok").

If you know of any errors in the lake's data or physical information, or are aware of any additional information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.




Lake ID: 270711
WMO: Bassett Creek
Volunteers: The Carlson Family

Westwood Lake, St. Louis Park, Hennepin Co.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 to 5	1 to 5
5/11/05	16.8				15	49		1.5	2	3
5/16/05	18.2				17	38		1.4		
7/6/05	27.8				9.7	37		1.2	2	2
7/20/05	31.5				28	158		1.1	5	4
9/2/05	22				100	69		1.4	4	5
9/16/05	26.3				100	321		1.7	2	4
9/30/05	16.8				13	28		1.6	2	




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus			F										
Chlorophyll a			C										
Secchi Depth			D										
Overall			D										

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C							B	B	C	C	C	D
Chlorophyll a	C							B	C	B	A	A	C
Secchi Depth	C							C	C	C	C	C	C
Overall	C							B	C	C	B	B	C

Source: Metropolitan Council and STORET data



Wilmes Lake (82-0090) City of Woodbury

Wilmes Lake, located in the City of Woodbury (Washington County) is classified as a minnow lake that experiences frequent fish kills. The lake has a surface area of 41 acres (1.3 miles around) and a maximum depth of 5.5 m (18 feet).

While there is currently no public access to the lake, one is planned at the northern end of the lake. The lake's 2,247-acre watershed translates to a large watershed-to-lake size ratio of 55:1. The larger the ratio, the greater the potential stress on the lake quality from surface runoff.

The future "ultimate" land uses within the lake's contributing watershed are expected to be: 16.4 percent single-family residential, 4.5 percent multi-family residential, 10.5 percent commercial/retail, 3.7 percent parks/open space, 1.0 percent ponds/wetlands, and 63.9 percent indirect drainage (City of Woodbury 1994).

Wilmes Lake has been involved in CAMP since 1994. In 2005, the lake was monitored 10 times between mid-April and mid-October. During each monitoring event, the lake was monitored for TP, CLA, TKN, Secchi transparency, as well as the perceived physical condition and recreational suitability. Results are presented on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	75.6	43.0	110.0	D
CLA (µg/l)	38.3	25.0	60.0	C
Secchi (m)	1.0	0.8	1.1	D
TKN (mg/l)	1.43	1.10	2.00	
Overall Grade				C

The lake's 2005 overall water quality grade of D is identical to those of 1997-1998 and 2001, and worse than the C's recorded in 1995-1996, 1999-2000 and 2003-2004, and the B recorded in 1994.

That said, the 1994 and 1995 CAMP data were actually collected in the northern basin of Wilmes Lake, while the 1996-2004 data were collected in the lake's south basin. For this reason, comparisons between the 1994-1995 database and the 1996-2004 should not be made.

When strictly looking at the lake's 1996-2005 water quality database (south basin), it is apparent that the lake's 2004 water quality was the best recorded to date.

No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the overall lake grade in the north basin seems to be C/B, while the overall grade of the south basin seems to be C/D+. To better understand the lake's water quality and where it may be heading, continued monitoring is suggested.


Throughout the monitoring period, the perceived physical condition and recreational suitability of the lake were ranked on a 1-to-5 scale by the volunteer monitors. These user perception rankings are presented in data tables and graphs on the information sheet. The mean physical condition ranking was 2.7 (between 2- "some algae present" and 3- "definite algae present"). The mean recreational suitability ranking was 2.7 (between 2- "minor aesthetic problem" and 3- "swimming slightly impaired").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Wilmes Lake, South Basin Woodbury, Washington Co.




Lake ID: 820090
W/D: South Washington
Volunteer: Bill Aamodt

◆ Sampling site
Contours in meters



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/17/05	16				84	127		0.9	2	2
5/15/05	13				42	95		1	2	2
5/17/05	14.5				32	110		1.1		
5/29/05	18.5				25	77		1.1	2	2
7/1/05	24.5				32	66		1	3	3
7/14/05	31				39	47		0.9	3	3
8/7/05	28				25	43		1.1	3	3
9/2/05	23				60	84		0.9	3	3
9/18/05	22.5				45	83		0.8	3	3
10/16/05	15.5				30	143		0.8	2	3




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus	C	D	D	D	D	D	D	D	D	D	C	C	D
Chlorophyll a	B	B	C	C	C	C	C	C	D	C	C	C	C
Secchi Depth	B	C	C	D	D	C	C	D	D	C	C	C	D
Overall	B	C	C	D	D	C	C	D	D	C	C	C	D

Source: Metropolitan Council and STORET data



Windsor Lake (27-0082) City of Minnetonka

Windsor Lake is a shallow 14-acre land-locked lake located within the City of Minnetonka (Hennepin County). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). There is very little other known morphological data available for the lake.

This marks the second year that Windsor Lake has been involved in CAMP (2004 being the first). On each of the sampling days the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between late-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	118.6	71.0	167.0	F
CLA (µg/l)	37.6	8.0	73.0	D
Secchi (m)	0.69	0.50	1.10	F
TKN (mg/l)	1.50	1.10	2.50	
Overall Grade				F

The lake's 2005 overall grade of D is better than the F recorded in 2004.

As mentioned earlier, there are no water quality data available for Windsor Lake other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.2 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.6 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Windsor Lake

Minnetonka, Hennepin Co.

Lake ID: 270082

WID: Minnehaha Creek

Volunteers: Sue Proudfoot
and Karen Mueller

● Sampling site

Contours in meters



2005 Data





Date	Surf Temp	Bot Temp	Surf DO	Bot DO	Chl a	Surf TP	Bot TP	Secchi	PC	RS
	C	C	mg/L	mg/L	µg/L	µg/L	µg/L	M	1-5	1-5
4/24/05	14				19	127		0.9	2	4
5/15/05	13				73	158		0.5	3	4
5/16/05	14				67	142		0.6		
5/28/05	16				57	167		0.6	5	4
6/9/05	26.9				12	79		1.0	3	4
6/24/05	27.4				8	71		1.1	4	4
7/1/05	28.5				26	81		0.8	3	5
7/23/05	26.5				54	124		0.6	4	5
8/7/05	26.2				34	119		0.7	3	5
8/28/05	23.1				26	104		0.5	2	5
9/16/05	22.7				44	142		0.6	2	5
9/29/05	15.8					117		0.7	3	5
9/29/05					42					
10/16/05	14.8				25	90		0.8	2	4

Lake Water Quality Grades Based on Summertime Averages

Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus												F	D
Chlorophyll a												D	C
Secchi Depth												F	F
Overall												F	D

Source: Metropolitan Council and STORET data



Windsor Lake (27-0082) City of Minnetonka

Windsor Lake is a shallow 14-acre land-locked lake located within the City of Minnetonka (Hennepin County). Because of the shallowness of the lake, the entire area is considered littoral zone (area of aquatic plant dominance) and it does not maintain a thermocline (a density gradient owed to changing water temperatures throughout the lake's water column). There is very little other known morphological data available for the lake.

This marks the second year that Windsor Lake has been involved in CAMP (2004 being the first). On each of the sampling days the lake was monitored for TP, CLA, TKN, and Secchi transparency, as well as the lake's perceived physical condition and recreational suitability.

The lake was monitored 14 times between late-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2005 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	118.6	71.0	167.0	D
CLA (µg/l)	37.6	8.0	73.0	C
Secchi (m)	0.69	0.50	1.10	F
TKN (mg/l)	1.50	1.10	2.50	
<i>Overall Grade</i>				D




The lake's 2005 overall grade of D is better than the F recorded in 2004.

As mentioned earlier, there are no water quality data available for Windsor Lake other than the 2005 CAMP data. Therefore it is not possible to determine any long-term or short-term trends. To better understand the lake's water quality and where it may be heading, additional years of data collection are needed.

Throughout the monitoring period, the volunteer(s) ranked their opinions of the lake's physical and recreational conditions on a 1-to-5 scale. The average user perception rankings were 3.2 for physical condition (between 3- "definite algae present" and 4- "high algal color"), and 4.6 for recreational suitability (between 4- "no swimming – boating ok" and 5- "no aesthetics possible").


If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us

Winkler Lake Benton Twp., Carver Co.



2005 Data

	Surf. Temp	Bot. Temp	Surf. DO	Bot. DO	CLA	Surf. TP	Bot. TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	ug/L	ug/L	ug/L	M	1 thru 5	1 thru 5
4/15/05	14.3		16.9		54	171		0.2	5	5
4/28/05	7.1		14.7		35	121		0.7	3	4
5/10/05	17.1		16.7		120	228		0.4	3	4
5/26/05	15.1				26	83		0.7	2	4
6/7/05	28.1		12.76		49	54		1	2	2
6/23/05	27		4.76		14	192		1	2	2
7/6/05	23.5		12.1		4.7	134		0.4	2	2
7/21/05	27.1		9.56		53	267		0.4	3	4
8/3/05	27.5		6.2		140	227			3	4
8/17/05	24.3		4.61		130	327		0.4	3	4
8/30/05	23.8				140	366		0.3	4	4
9/15/05	20.8		8.57		64	497		0.4	3	4
9/26/05	16.1		13.4		130	485		0.3	3	4
10/12/05	12.1				35	260		0.8	2	4




Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus			F				F	F	F		F		F
Chlorophyll a			A				D	F	C		F		F
Secchi Depth			C				F	F	F		F		F
Overall			C				F	F	D		F		F

Source: Metropolitan Council and STORET data



Wood Lake [Burnsville] (19-0024) *Black Dog Watershed Management Commission*

Wood Lake is a 9-acre lake located within the City of Burnsville (Dakota County). The maximum depth of the lake is 4.5 m (14.7 feet). Because the maximum depth is only 4.5 m (almost 15 feet), the entire lake area is considered littoral zone (the area of aquatic vegetation dominance). The majority of the land within the lake's 157-acre immediate watershed is urban/developed. The resulting watershed-to-lake size ratio is 17:1. The greater the ratio, the greater the potential stress on the lake from surface runoff.

This was the tenth year that Wood Lake has been involved in CAMP. The lake (which has been enrolled in CAMP since 1996) was monitored 14 times between mid-April and mid-October, 2005. The resulting data and graphs appear on the next page.

2004 summer (May-September) data summary

<i>Parameter</i>	<i>Mean</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Grade</i>
TP (µg/l)	53.5	32.0	102.0	C
CLA (µg/l)	26.2	5.0	55.0	C
Secchi (m)	1.4	0.5	2.7	C
TKN (mg/l)	1.54	0.98	2.30	
Overall Grade				C

The 2005 overall lake quality grade for Wood Lake, calculated from the individual parameter grades, is C (similar to 1996, 1997, and 1999-2004, but worse than the B of 1998).

As mentioned in previous reports, an alum sulfate (alum) treatment in October of 1997 resulted in the lake's best water quality year in 1998. An alum treatment to a lake involves adding the chemical to bind and precipitate phosphorus, removing it from the water column, and sealing the bound phosphorus in the sediment rendering it inactive for release to the overlying water. By removing the phosphorus from the water column and locking it in the sediments, its availability for plant growth is reduced. The success of this treatment depends on the lake's residence time (the time it would take to entirely refill the lake basin with water if it were empty) and external phosphorus load. The shorter the residence time and the larger the external phosphorus load, the quicker new sources of phosphorus will replenish the water column. Since 1998, however, the lake's water quality has been more comparable to that of the pre-alum treatment years of 1996 and 1997 as opposed to that of 1998. For this reason it seems that the alum treatment has not been wholly successful.

Other than the data collected through CAMP, there are no historical water quality data available for Wood Lake. A search through STORET (EPA's nationwide water quality database) came up empty. Therefore the only summertime data available are those from 1996-2005. No statistically significant long-term trend is evident from the lake's water quality database, in the short-term however, the lake's water quality seems well represented by an overall grade of C. To better understand the water quality of the lake and determine in what direction the water quality is heading, additional years of data collection are needed.

The volunteer monitor's perception of the lake's physical and recreational conditions were ranked on a 1-to-5 scale during each monitoring event. The rankings are shown on the information sheet on the next page. The average user perception rankings were 3.6 for physical condition (which falls between 3- "definite algae present" and 4- "high algal color"), and 3.6 for recreational suitability (between 3- "swimming slightly impaired" and 4- "no swimming: boating ok").

The Fisheries Section of the Minnesota Department of Natural Resources (MDNR) has conducted a fisheries survey on the lake. Information on the survey can be obtained through the MDNR Fisheries Section by calling (651) 297-4916 or by downloading the information off the Internet at <http://www.dnr.state.mn.us/lakefind/>.

If you notice any errors in the lake's data or physical information, or are aware of any additional or missing information, please contact Randy Anhorn of the Metropolitan Council at (651) 602-8743 or randy.anhorn@metc.state.mn.us.

Wood Lake

Burnsville, Dakota Co.

Lake ID: 190024

WMO: Black Dog

Volunteer: David Bess

● Sampling site

Contours in meters



2005 Data





	Surf Temp	Bot Temp	Surf DO	Bot DO	CIA	Surf TP	Bot TP	Secchi	PC	RS
Date	C	C	mg/L	mg/L	sq/L	ug/L	ug/L	M	1-5	1-5
4/18/05	18				22	49		1.4	2	2
5/5/05	13.5				11	46		1.6	2	2
5/17/05	13				10	54		1.7	2	2
5/18/05	16.2				12	47		1.9		
6/2/05	21.6				26	82		2.3	2	4
6/6/05	24.6				14	32		2.3	3	4
6/23/05	26.3				43	35		1.3	4	4
7/6/05	25.5				5	102		0.6	4	4
7/20/05	26.9				26	48		0.5	5	4
8/2/05	30.5				30	41		0.3	5	4
8/23/05	24.6				43	35		0.3	5	4
9/12/05	23.8				55	66		1.1	4	4
10/6/05	15.8				20	68		1.2	3	4
10/18/05	14.9				14	70		1.9	3	4

Lake Water Quality Grades Based on Summertime Averages

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Total Phosphorus													
Chlorophyll a													
Secchi Depth													
Overall													

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Phosphorus				C	C	B	C	C	C	C	C	C	C
Chlorophyll a				B	B	B	B	B	C	C	B	B	C
Secchi Depth				C	C	C	C	C	C	C	C	C	C
Overall				C	C	B	C	C	C	C	C	C	C

Source: Metropolitan Council and STORET data



CONCLUSIONS

To date, the Council's lake monitoring programs (including the staff- and volunteer- monitoring programs) and have provided an important tool for making informed lake management decisions. Data from our regional lake monitoring programs are frequently used to determine possible trends in in-lake water quality, estimate expected ranges in water quality of unmonitored lakes, examine intra-and inter-regional differences, determine potential impairments due to water quality, and investigate the relationships between landscape and water quality. The Council's lake program collected data on 172 lake sites in 2005, including 12 lake sites on 10 lakes monitored by the Council and 160 lakes monitored by CAMP volunteers.

Eighty-six of the 172 lakes monitored in 2005, are listed by the MPCA as impaired waters due to excessive nutrients (phosphorus) affecting the lakes' ability to support their designated recreational uses. Eighty of those lakes were monitored through CAMP, and six were monitored by Council-staff. To learn more about the listing and potential next steps <http://www.pca.state.mn.us/water/tmdl/index.html>.

The year 2005 marked the thirteenth year that the Council-sponsored volunteer monitoring program, entitled "The Citizen-Assisted Monitoring Program" (CAMP), was used to increase our knowledge of the water quality of area lakes. Once again volunteers measured surface water temperature and transparency, and collected surface water samples that were analyzed for total phosphorus, total Kjeldahl nitrogen, and chlorophyll-a on a biweekly basis from mid-April to mid-October (approximately 14 sampling events).


This year's CAMP monitoring included 17 lakes never before monitored by the Council (Birch, Cobblestone, East, Edith, Langton [sites 1, 2, and North Basin], Legion Pond, Moody, O'Connor, School, Shaver, and St. Croix (five sites) lakes), and 129 lake sites returning from 2004. The 2005 lakes program included lake data from all of the 26 watersheds/municipalities/counties represented in the 2004 program. Additionally, the 2005 CAMP program added three new enrolling groups to its growing list of monitoring partners.

The greatest percentage of the lakes monitored through CAMP in 2004 received an overall water quality grade of "C" (38.7 percent). When combining the CAMP and Council-staff monitored lake grades 40.1 percent of the lakes received an overall grade of "C". The water quality of these lakes is considered average as compared to others in the seven-county metropolitan area. When comparing the percentage of above-average lakes (those receiving grades of "A" or "B") to below-average lakes (those receiving "D" or "F"), more lakes were below average (34 percent to 27 percent). The complete 2005 CAMP lake report card grade tally (for those lakes with sufficient data) assigned "A's" to 22 lakes (14.1 percent) and "B's" to 20 lakes (12.9 percent). Sixty lakes acquired "C's" (38.7 percent), 34 received "D's" (21.9 percent), and 19 obtained an "F" (12.3 percent).

The 22 lakes that received "A" grades, within the top 10-percentile range for Metro Area lakes include: Big Carnelian, Big Marine, Brickyard, Cenaiko, Cloverdale, Courthouse, DeMontreville, Edith, Fireman's, Half Breed (Sylvan), Jane, Kingsley, Lac Lavon, Little Carnelian, Long (Mahtomedi), MacDonald's Pond, Olson, Square, St. Joe, Sunset, Terrapin, and West Boot lakes.

The lakes receiving the lowest water quality grade "F" include: Benton, Benz, Cobblecrest, Campbell, Cornelia, East, Farquhar, Gaystock, Goose (Waconia), Hart, Hazeltine, Highland, Langton (North Basin), Long (Apple Valley), Loon, Maria, Meadow, Pike (Scott County), and Winkler lakes.

The worst summertime TP and CLA means recorded in CAMP 2005 were; a TP mean of **327.8** µg/l on Campbell Lake (Scott County), and a CLA mean of **223.4** µg/l on Highland Lake (Anoka County). The lowest Secchi transparency mean recorded in 2005 was **0.28** m, recorded on Benton, and Dean lakes (Carver and Scott counties).



Similar to past years, there is no distinct pattern as to where lakes with specific water quality were located. As was observed in the past, the only similarity between the majority of the D and F grade lakes is their size and mean depth. These lakes are generally shallow with small surface areas. In some cases, the lakes are nothing more than deep marshes with an excess of emergent and submergent vegetation. As was mentioned in past reports, this distinction is important for three reasons: 1) deeper lakes have a greater ability to incorporate nutrients and trap them in the sediments, where they are not available for plant growth (macrophyte and/or algae), 2) shallow lakes tend not to stratify during the summer months, allowing the potential release of phosphorus from bottom sediments to rise through the water column and become available for plant growth, and 3) the small surface areas of these lakes generally result in larger watershed-to-lake ratios. Lakes with large watershed-to-lake ratios, have to handle larger runoff loads for their size than do larger lakes in a similar-sized watershed.

The lakes with above-average water quality (grades of “A” and “B”) similarly were not area specific. They were located in all seven of the region’s counties (lakes receiving an “A” grade were found in all seven Metro counties). Common characteristics of the above-average lakes were: they have deeper maximum and mean depths, they develop and maintain a thermocline, they have small contributing watersheds relative to the lakes’ surface area, and there was little construction within the lakes’ watershed.

Similar to that mentioned in past reports, analysis of The 2005 CAMP lake water quality nutrient data (TP and CLA) produced no “statistically significant” long-term water quality trends. The major reasons for not being able to determine trends are: 1) the majority of lakes in the Metropolitan Area have limited and/or fluctuating databases, or 2) if a sufficient database does exist, analysis revealed no “statistically significant” trend.

In many cases, however, lakes’ Secchi transparency databases are much more extensive than their related nutrient database. The reasons being that: 1) it is much less expensive to do Secchi readings than it is to have water samples analyzed at a laboratory, and 2) the development of the MPCA coordinated a volunteer Secchi transparency monitoring program in the early-1970’s. For these reasons, a few CAMP lakes do have

sufficient information to determine statistically significant trends in Secchi transparency. Of the CAMP 2005 lakes assessed (those with sufficient data), 14 showed an improving trend in water clarity (Big Marine, DeMontreville, Elmo, Halfbreed/Sylvan, Langton (south basin), Little Carnelian, Long (May Township), McKusick, Olson, Sunset, Valentine, Valley, Waconia, and West Boot lakes) and four showed a negative trend (Farquhar, Lac Lavon, Markgrafs, and Square lakes) (MPCA 2005).

Of the 129 repeat lakes which a sufficient database from 2004, 24 had a worse overall water quality grade in 2005 (Bald Eagle [Site-1], Barkers, Bavaria, DeMontreville, Forest [West Basin], Golden, Goose [Waconia], Hart, Herber's Pond, Island, Kismet, Long [Stillwater], Long [Mahtomedi], Louise, MacDonald's Pond, McDonald, McKusick, Mud, Oak, Olson, Sand, Spring, Schroeder's Pond, and Westwood lakes), and 14 had better overall water quality grades in 2005 (Big Comfort, Cloverdale, Downs, Eagle (Carver County), George Watch, Hay, McCarrons, Miller, Prior [Lower], Prior [Upper], Staples, Swede, Wilmes, and Windsor lakes), and 91 had the same overall water quality grade for both years. By further breaking down the 91 lakes that had identical overall grades in 2004 and 2005, 44 had similar summertime mean conditions in both 2004 and 2005 (mean TP, CLA and Secchi transparency), 29 had worse means in 2005, and 18 had better means in 2005.

The location breakdown of the 24 lakes with worse overall water quality grades in 2005 as compared to 2004 was: three in Anoka County, three in Carver County, one in Dakota County, one in Ramsey County, 15 in Washington County, and one in Scott County. The 14 lakes with better water quality in 2005 were located in Anoka County (one), Carver County (three), Hennepin County (one), Ramsey County (one), Scott County (two), and Washington County (six).

Water quality data from the 129 repeat lakes seem to indicate that the Metro Area lakes experienced slightly worse water quality conditions in 2005 as compared to 2004. This after two years where the lake monitoring program reveal slightly better water quality as that recorded during the previously monitored years (2003 better than 2002, and 2004 better than 2003). A recently conducted trend analysis by MPCA on lakes with extensive Secchi transparency databases however, revealed that while the majority of statistically assessed lakes showed no trends in water clarity (either negative or improving), more lakes showed an improving trend than a negative trend (MPCA 2005).

Since 1980, 297 area lakes have been monitored through the Council's Lake Program (including Council-staff monitoring and CAMP). Some of the lakes have multiple monitoring sites [316 sites]. The list of lakes in the Council's monitoring database is shown in Appendix C. The resulting data from the Council's lake monitoring program are permanently stored in the U.S. EPA's national water quality data bank, STORET (stands for STORage and RETrieval). The majority of the 316 lake sites have been revisited on a rotating schedule throughout the past 26 years to develop a working baseline to help determine possible trends and to aid lake and watershed managers in their decision making. While the Council has done its best to enhance and expand the region's lake water quality database, it is apparent that one of the most economical and efficient method to expand knowledge of our lakes has been with the assistance of volunteers and cooperation and financial support of watershed management organizations, counties, and cities. So while the first 13 years of CAMP have been very successful, our future goal is to continue to expand the coverage of our lake monitoring program in order to better understand and manage the areas water resources.

The Council's lake monitoring program, especially the use of volunteer monitors through CAMP, has played a key role in the Council's recent efforts to use satellite images to assess annual lake water clarity for the region as a whole. The monitoring program provides "ground-based" measurements used to calibrate mathematical models, which in turn are used to interpret the satellite images. The use of satellite technology

provides a cost-effective way to extend the analysis of the region's lake water quality from just the lake's involved in our ground-based programs to all the lakes in the region. Over time, the satellite-based information can be used to detect how lake trophic conditions (especially water clarity) have changed over time and space in relation to changes in land-use and land-cover conditions.

Results of the 2005 satellite assessment of the region revealed similar results to that found through the 2004 ground-based monitoring programs, that the region experienced slightly worse lake water quality in 2005 than that recorded in 2004. The complete results of the 2005 satellite analysis can be at <http://www.metrocouncil.org/planning/environment/TCWaterClarity2005.pdf>.

If you have questions pertaining to the lake data or descriptions contained in this report, inquiries about CAMP, or suggestions of lakes the Council should consider monitoring in the future, please contact Randy Anhorn at the Metropolitan Council (651) 602-8743 or randy.anhorn@metc.state.mn.us.

REFERENCES

- American Public Health Association. 1992. *Standard Methods for the Examination of Water and Wastewater*. 18th ed.
- Anhorn, R.J. 1993. *Handbook for the Citizen-Assisted Lake Monitoring Program*. Metropolitan Council. St. Paul, MN.
- Anhorn, R.J. 1994. *A 1993 Study of the Water Quality of 43 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 32-94-012.
- Anhorn, R.J. 1995. *A 1994 Study of the Water Quality of 51 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 32-95-011.
- Anhorn, R.J. 1996. *A 1995 Study of the Water Quality of 59 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 32-96-014.
- Anhorn, R.J. 1997. *A 1996 Study of the Water Quality of 66 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 32-97-004.
- Anhorn, R.J. 1998. *A 1997 Study of the Water Quality of 71 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 32-98-007.
- Anhorn, R.J. 1999. *A 1998 Study of the Water Quality of 70 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 32-99-008.
- Anhorn, R.J. 2000. *A 1999 Study of the Water Quality of 113 Metropolitan Area Lakes*. Metropolitan Council Publ. No. EPE-00-479.
- Anhorn, R.J. 2001. *A 2000 Study of the Water Quality of 124 Metropolitan Area Lakes*. Metropolitan Council Publ. No. EPE-01-502.
- Anhorn, R.J. 2002. *A 2001 Study of the Water Quality of 132 Metropolitan Area Lakes*. Metropolitan Council Publ. No. EPE-02-516.
- Anhorn, R.J. 2003. *A 2002 Study of the Water Quality of 137 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 32-03-019.
- Barr Engineering. 1993. *1993 Marcott Lakes Water Quality Study: EP-78 and EP-80*. Barr Engineering. Minneapolis, MN.
- Barr Engineering. 1993. *1990 and 1991 Lake water Quality Inventory and Historical Water Quality Trend Analysis: Lakes Ann, Duck, Hyland, Lotus, Lucy, Mitchell, Red Rock, Rice Marsh, Riley, Round, Staring and Susan*. Barr Engineering. Minneapolis, MN.
- Barr Engineering. 1994. *Sweeney Lake: Watershed and Lake Management Plan*. Barr Engineering. Minneapolis, MN.
- Barr Engineering. 1997. *Project Assessment and Evaluation for the Crystal/Keller Lake Water Quality Improvement Project*. Barr Engineering. Minneapolis, MN.

- Barr Engineering. 1997. *North Rice, South Rice, and Grimes Ponds Watershed and Lake Management Plan*. Barr Engineering. Minneapolis, MN.
- Barr Engineering. 2001. *2000 Lake Water Quality Study: Northwood Lake, Parkers Lake, Sweeney Lake, and Twin Lake*. Barr Engineering. Minneapolis, MN.
- Barr Engineering. *Personal Communication*. Barr Engineering. Minneapolis, MN.
- Beduhn, R.J. 1993. *LCMR Lake Monitoring Study: The Effects of Aeration/Circulation on Lake and Reservoirs*. Prepared for the Legislative Commission on Minnesota Resources.
- Black Dog Watershed Management Commission. 1987. *Watershed Management Plan*. Orr, Schelen, Mayeron and Associates. St. Paul, MN.
- Blue Water Science and Bonesreoo, Rosene, Anderlik and Assoc. 2005. *Lake Management Plan for Alimagnet Lake, Dakota County, Minnesota*. Blue Water Science, St. Paul, MN.
- Bokenmeier, D. 1989. *Staying Ahead of the Game*. North American Lake Management Society: Lake Line. 9(8): 2-16.
- Browns Creek Watershed Management Organization . 1990. *Watershed Management Plan*. Washington County Soil and Water Conservation District.
- Carlson, R.E. 1977. *Trophic Status Index Indicator of Lakes*. Limnology Oceanography 22:361-369.
- Capital Region Watershed District. 2003. *Lake McCarrons Management Plan*. The Osgood Group. Shorewood, MN.
- Carver County Planning and Zoning Department. 1999. *Carver County 1999 Water Quality Report*. Carver County Planning and Zoning Department.
- Carver Creek Watershed Management Organization. 1990. *Water Management Plan*. Carver County Soil and Water Conservation District.
- City of Inver Grove Heights. 1993. *Simley Lake Water Quality Study*. Orr, Schelen, Mayeron and Associates.
- City of Woodbury. 1994. *Surface Water Management Plan*. Bonestroo Rosene Andelik & Associates. St. Paul, MN.
- Cole, G.A. 1983. *Textbook of Limnology: 3rd ed*. C.V. Mosby Company. St. Louis, MI.
- Coon Creek Watershed District. 1985. *MSA-509 Plan*. Israelson and Associates, Inc Bloomington, MN.
- Environmental Research Group, Inc. 1986. *Management Alternatives Report on the Diagnostic-Feasibility Study for Golden Lake. Anoka County, MN*. Environmental Research Group, Inc.

- Forest Lake Watershed Management Organization. 1988. *Surface Water Management Plan*. Washington County Soil and Water Conservation District.
- Gersmehl, C., J. Drake and D. Brown. 1986. *Minnesota Water: A Geographical Perspective*. Water Resources Research Center. Public Report Series No. 4.
- Gun Club Lake WMO. 1989. *Management Plan*. James M. Montgomery Engineers (Montgomery Watson). Wayzata, MN.
- Hartsoe, J.A. and R.A Osgood. 1991. *A 1991 Study of the Water Quality of 17 Metropolitan Area Lakes*. Metropolitan Council Publ. 590-92-006.
- Hanson, K. 1995. *Personal Communication*. City of Inver Grove Heights, MN.
- Kernik, S. 1995-1999. *Personal Communication*. City of Woodbury, MN.
- Ludvig, M. 1994. *Lake Waconia Surface Water Report*. Carver County Environmental Services. Chaska, MN.
- Maloney, T.E. (ed.). 1979. *Lake and Reservoir Classification Systems*. U.S. EPA Environmental Research Lab. Corvallis, OR. EPA-600/3-79-074.
- Marine on St. Croix Watershed Management Organization. 1991. *Water Management Plan*. Washington County Soil and Water Conservation District.
- McComas, S. 2003. *Using Barley Straw to Improve Water Clarity in Valley Lake, in 2002, Lakeville Minnesota*. Blue Water Science. St. Paul, MN.
- McComas, S. and R. Anhorn. 2004. *The Use of Barley Straw as an Organic Carbon Amendment to Enhance Heterotrophic Processes and Increase Water Clarity on Valley Lake*. Unpublished study.
- McComas, S. 2005. *Personal Communication*. Blue Water Science. St. Paul, MN.
- Metropolitan Council. 1986. *Water Resources Management: Development Guide/Policy Plan*. St. Paul, MN.
- Metropolitan Council. 1997. *Lake McCarrons Wetland Treatment System—Phase III Study Report*. St. Paul, MN.
- Middle St. Croix Watershed Management Organization. 1991. *Watershed Management Plan*. Washington County Soil and Water Conservation District.
- Minnehaha Creek Watershed District. 1987. *Water Resources Management Plan*. James M. Montgomery Engineers (Montgomery Watson). Wayzata, MN.
- Minnehaha Creek Watershed District. 1996. *Water Resources Management Plan*. Wenck Associates. Maple Plain, MN.

- Minnesota Department of Conservation. 1967. *Metropolitan Lake Inventory*. Minnesota Department of Conservation. St. Paul, MN.
- Minnesota Department of Conservation. 1968. *Inventory of Minnesota Lakes*. Minnesota Department of Conservation. St. Paul, MN.
- Minnesota Department of Natural Resources. Various years. *Lake Fishery Surveys*. Minnesota Department of Natural Resources. St. Paul, MN.
- Minnesota Department of Natural Resources. 1996. *Report on the Status of the DNR Metro Region Trout Resources*. A Metro Region Trout Committee Report. Minnesota Department of Natural Resources. St. Paul, MN.
- Minnesota Department of Natural Resources. 2004. **Draft**. *Evaluation of the potential to selectively control Eurasian watermilfoil with fluridone herbicide in Minnesota 2001-2003*. Minnesota Department of Natural Resources. St. Paul, MN.
- Minnesota Pollution Control Agency. 1985. *A Citizens' Guide to Lake Protection*. Minnesota Pollution Control Agency. St. Paul, MN.
- Minnesota Pollution Control Agency. 1994. *Lake Assessment Program: Lake Waconia*. Minnesota Pollution Control Agency. St. Paul, MN.
- Minnesota Pollution Control Agency. 2003. *Fact Sheets on Lake Transparency Trends*. Minnesota Pollution Control Agency. St. Paul, MN.
- Montgomery Watson. 1990. *Lake Alimagnet Diagnostic - Feasibility Study*. Montgomery Watson. Wayzata, MN.
- Montgomery Watson. 1994-1998. *Personal Communication*. Montgomery Watson. Wayzata, MN.
- Newman, R.M., D.W. Ragsdale, and D.D. Biesboer. 1996. *Can Eurasian Watermilfoil be Managed in Minnesota by Biological Control with Native or Naturalized Insects?* Third progress report to the Minnesota Department of Natural Resources, Ecological Services, St. Paul, MN.
- Nichols, A.B. 1992. *Citizens Monitor Water Quality*. Water Environment and Technology. March, 1993. pp.55-59.
- Oberts, G.L. 1982. *Nonpoint Source Pollution in the Metropolitan Area: Technical Completion Report*. Metropolitan Council Publ. No. 10-82-016.
- Oberts, G.L and R.A Osgood. 1988. *Lake McCarrons Wetland Treatment System: Final Report the Function of the Wetland Treatment System and the Impacts on Lake McCarrons*. Metropolitan Council Publ. No. 590-88-095.
- Osgood, R. 1981. *A Study of the Water Quality of 60 Lakes in the Seven-County Metropolitan Area*. Metropolitan Council Publ. No. 01-81-047.

- Osgood, R. 1982a. *A 1981 Study of the Water Quality of 30 Lakes in the Seven-County Metropolitan Area*. Metropolitan Council Publ. No. 10-82-005.
- Osgood, R.A. 1982b. *Using Carlson's Trophic State Indices in Regional Water Quality Assessment*. Water Resources Bulletin 18:67-74.
- Osgood, R.A. 1983. *Diagnostic-Feasibility Study of Seven Metropolitan Area Lakes. Part One: General Overview*. Metropolitan Council Publ. No. 10-83-092. *Part Two: Bryant, Elmo, Fish, George, Riley, Spring and Square Lakes*. Metropolitan Council Publ. Nos. 10-83-093 a-g.
- Osgood, R.A. 1984a. *A 1983 Study of the Water Quality of 28 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 10-84-037.
- Osgood, R.A. 1984b. *A 1984 Study of the Water Quality of 43 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 10-84-172.
- Osgood, R.A. 1985. *A 1985 Study of the Water Quality of 32 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 10-85-156.
- Osgood, R.A. 1988a. *A 1986/1987 Study of the Water Quality of 10 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 590-88-037.
- Osgood, R.A. 1988b. *The Limnology, Ecology and Management of Twin Cities Metropolitan Area Lakes*. Metropolitan Council Publ. No. 590-88-123.
- Osgood, R.A. 1989a. *An Evaluation of the Effects of Watershed Treatment Systems on the Summertime Phosphorus Concentration in Metropolitan Area Lakes*. Metropolitan Council Publ. No. 590-89-062b.
- Osgood, R.A. 1989b. *A 1989 Study of the Water Quality of 20 Metropolitan Area Lakes*. Metropolitan Council Publ. No. 590-89-129.
- Osgood, R.A. 1989c. *An Evaluation of Lake and Stream Monitoring Programs in the Twin Cities Metropolitan Area*. Metropolitan Council Publ. 590-89-128.
- Osgood, R.A. 1989d. *Assessment of Lake Use-Impairment in the Twin Cities Metropolitan Area*. Metropolitan Council Publ. No. 590-89-130.
- Osgood, R.A. 1990. *A 1990 Study of the Water Quality of 21 Lakes in the Twin Cities Metropolitan Area*. Metropolitan Council Publ. No. 590-90-182.
- Osgood, R.A. and J.E. Stiegler. 1990. *The Effects of Artificial Circulation on a Hypereutrophic Lake*. Water Resource Bulletin 26:209-217.
- Osgood, R.A. 2000. *City of Victoria Local Water Resources Management Plan: Victoria Lakes – 2000*. Ecosystem Strategies. Shorewood, MN.
- Prior Lake - Spring Lake Watershed District. 1991. *Water Resources Management Plan*. James M. Montgomery Engineers (Montgomery Watson). Wayzata, MN.

- Prior Lake - Spring Lake Watershed District. 1998. *Draft-Water Resources Management Plan*. James M. Montgomery Engineers (Montgomery Watson). Wayzata, MN.
- Prior Lake - Spring Lake Watershed District. 2001. *Final Report on the Prior-Spring Lake Improvement Project*. James M. Montgomery Engineers (Montgomery Watson). Wayzata, MN.
- Prior Lake - Spring Lake Watershed District. 2004. *A Sustainable Water Quality Management Plan for Spring and Prior Lakes*. PLSLWD, Prior Lake, MN.
- Rice Creek Watershed District. 1987. *Water Resources Management Plan*. James M. Montgomery Engineers (Montgomery Watson). Wayzata, MN.
- Riley-Purgatory-Bluff Creek Watershed District. 1982. *Management Plan*. Barr Engineering. Minneapolis, MN.
- Short Elliot Hendrickson Inc. 1998. *How's the Water? Plymouth, Minnesota Water Resources Management Plan*. Short Elliot Hendrickson. St. Paul, MN.
- Shingle Creek Watershed Management Organization. 2003. *Watershed Management Plan*. Wenck Associates, Inc. Maple Plain, MN.
- Simpson, J.T. 1991. *Volunteer Lake Monitoring: A Methods Manual*. EPA 440/4-91-002.
- Six Cities Watershed Management Organization. 1988. *Water Management Plan*. Barr Engineering. Minneapolis, MN.
- South Washington Watershed District. 1997. *Watershed Management Plan*. Bonestrro, Rosene, Anderlik and Associates. St. Paul, MN.
- Square Lake Clean Water Partnership Report. 2001. *Square Lake, Washington County Phase I Resource Investigation..* Washington County Soil and Water Conservation District. Stillwater, MN.
- Square Lake Clean Water Partnership Report. 2001. *Square Lake, Washington County Phase I Resource Investigation..* Washington County Soil and Water Conservation District. Stillwater, MN.
- St. Croix Basin Water Resources Planning Team. 2004. *St. Croix Basin Phosphorus-Based Water Quality Goals*.
- U.S. Environmental Protection Agency. 1997. *Watershed Protection: Clean Lakes Case Study. Use of Aquatic Weevils to Control a Nuisance Weed in Lake Bomoseen, Vermont*. EPA-841-F-97-002.
- U.S. Geological Survey. 1976. *Hydrology of the Lakes in the Minneapolis-St. Paul Metropolitan Area: A Summary of Available Data*. Water-Resources Investigations 76-85.
- U.S. Geological Survey. 2001. *Response of the St. Croix River Pools, Wisconsin and Minnesota, to Various Phosphorus-Loading Scenarios*. Water-Resources Investigations Report 02-4181.

- Valley Branch Watershed District. 2005. *Watershed Management Plan*. Barr Engineering Company. Minneapolis, MN.
- Vermillion River Watershed Management Commission. 1989. *Watershed Management Plan*. James M. Montgomery Engineering. Wayzata, MN.
- Washington Conservation District. 2002. *August 2001-July 2002 Zooplanton Monitirng Summary Report for Square Lake in Washington County, MN*. Washington Conservation District. Stillwater, MN.
- Wenck Associates, Inc. 1987. *Lake Diagnostic/Feasibility Study: Prepared for Forest Lake Watershed Management Organization*. Wenck Associates Inc. Wayzata, MN.
- Wenck Associates, Inc. 1987. *Diagnostic/Feasibility Study for Crooked Lake, Anoka County*. Wenck Associates Inc. Wayzata, MN.
- Wetzel, R.G. 1983. *Limnology*. Saunders College Publishing. Chicago, IL.
- Wilson, B. 1990. *Lake Water Quality Summary of Shields, Bone, Half Breed, and Forest Lakes*. Forest Lake Watershed Management Organization.

APPENDIX A
2005 CAMP Lake/Watershed Characteristics

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
Acorn 82-102	44	296	7:1	3.0	0.7	440	100	0	N			
Alimagnet 19-21	109	1,094	10:1	3.0	1.5	545	100	12	N	C	3.2	
Armstrong 82-116-02	39			1.5	1.0	128	100		N	N		
Barker 82-76	45	823	19:1	9.0	4.4	648			Y	N		
Bass (Henn) 27-98	194	3,100	16:1	9.4	3.1	1,979	82		Y	N	2.3	
Bass (StLP) 27-15	95											
Bass (Wash)82-35	81			4.3			100		N	N		
Bavaria 10-19	200	711	3.5:1	18.3	5.6	3,674	40		Y	Y		Centrachid
Benton 10-69	115	322	3:1	2.0			100		N	N		
Benz 82-120	36						100		N	N		
Big Carnelian 82-49	455	1,900	4:1	20.0	9.8	14,560	28		Y	Y		
Big Comfort 13-53	219			14.3			41		Y	Y		
Big Marine 82-52	1,706	2,659	1.5:1	15.2	7.6	42,527	67		Y	Y		
Birch 13-42	65											
Bone 82-54	212	5,177	24:1	9.8	3.7	2,820	59	3	Y	Y		
Brickyard 10-225	17			13.1			35		Y	N		
Burandt 10-84	93			7.3			72		Y	N		

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
Campbell 10-127	72			2.0			100		N	N		
Carol 82-17	63	375	6:1	1.8	0.9	186	100		N	N		
Cates 70-18	27			4.0			100		N	N		
Cenaiko 2-654	29			9.1			40		Y	N	0.6	Stocked w/Trout - Fishing Pier
Clear 82-163	400			8.5	3.7	4,800	67		Y	Y	3.9	Walleye
Cloverdale 82-9	45	819	18:1	8.5	3.0	450	86		Y	N		
Cobblecrest 27-53	10									N		
Cobblestone19-456												
Colby 82-94	71	8,088	114:1	3.4			100		N	N		
Cornelia 27-28	71	8,088	114:1	3.4			100		N	N		
Courthouse 10-5	10			17.4			30		Y	N	0.6	Stocked w/Trout
Crystal(Bnsv)9-27	292	2,001	7:1	11.3	3.1	2,920	72		Y	Y		Panfish - Fishing Pier
Crystal(rob) 27-34	76	1,272	17:1	10.4	3.7	917	68		Y	Y	1.4	Centrarchid - Fishing Pier
Dean 70-74	128						100		N	N		
DeMontreville82-101	160	1,108	7:1	7.3	2.4	1,280	90		Y	Y		
Downs 82-110	35	2,400	69:1	2.1	1.5	175	100		N	N		
Eagle(Crv)10-121	233	1,050	4.5:1	4.0	1.2	920	100		N	Y		Natural Environment
Eagle(m.g.) 27-111	291	3,220	11:1	10.4	3.8	3,667	68		Y	Y	3.2	Centrarchid
Earley 19-33	29	1,629	56:1							N		
East 19-349	40											
East Boot 82-34	47	93	2:1	8.2	0.9	282	84		Y	Y		

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
Edina 27-29				1.0			100		N	N		
Edith 82-4	81	1,576	19:1	13.0					Y			
Elmo 82-106	284	1,191	4:1	41.7			22		Y			
Farquhar 19-23	63	353	6:1	3.0	1.4	290	100		N	N		
Fireman's 10-226	8			7.0			88		Y			
Fish (Grant) 82-137	21			10.4			67		Y			
Fish(Scott) 70-69	171	660	4:1	8.5	4.4	2,468	43		Y	Y		Centrachid
Fish (Wash) 82-64	72	683	9.5:1	3.0	1.5	360	100		N	N		
Forest 82-159	2,249	4,285	2:1	11.5	3.4	24,986	68	14	Y	Y		
French 27-127	352	870	4:1	1.0					N	Y		
Gaystock 10-31	105			5.0			100		N	N		
George Watch 2-5	528			2.0	1.5	2,587	100		N	Y		
German 82-56	109											
Goetschel 82-313	22	2,812	122:1	4.2	1.2	88	100		N	N		
Goggins 82-77	11						100		N	N		
Golden 2-45	57	7,680	135:1	7.3	2.5	463	90	1	Y	Y	1.5	
Goose (Sendia) 82-59	83			7.6	2.4	664	55			Y		
Goose(Wac)10-89	407	1,100	27:1	3.0	1.5	2,035	100		N	C		Natural Environment
Grace 10-218	22			6.7			79					
Hafften 27-199	43		13.4				60		Y	Y		
Half Breed 82-80	75	303	4:1	10.3	1.7	420	67		Y	N		

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
Hart 2-81	8						100		N	N		
Harvey 27-??				0.7			100		N	N		
Hay 82-65	33									N		
Hazeltine 10-14	236			2.0			100		N	N		
Henry 10-175	77			1.5			100		N	N		
Herbers Pnd 82-15-01				2.0			100		N	N		
Highland 2-79	22			1.0			100		N	N		
Hydes 10-88	215	430	2:1	5.5	3.0	2,150	88		Y	Y		
Island 2-22	67			6.7			87		Y	N		
Jane 82-104	155	1,402	9:1	12.0	3.7	1,860	72		Y	Y		
Jellum's 82-5202	72	333	4.6:1	4.9	2.4	569	100		N	N		
Keller (Brn)19-25	60			2.5	1.5	300	100		N	N		
Kingsley 19-30	44	193	4:1	4.0			100		N	N	1.7	
Kismet 82-333										N		
Klawitter 82-368	4.5	168	37:1				100					
La 82-97	35			3.5			100		N	N	1.3	
Lac Lavon19-446	69	306	4:1	9.8			26		Y	N	2.3	Stocked w/Trout - Fishing Pier
Langton 62-49	30	257	9:1	1.5	1.2	120	100		N			
Lee 19-29	25	324	13:1	5.2			100		N	N	1.0	
Legion Pond 82-462	16	224	14:1									
Libbs 27-85	23			2.1			100		N	N		

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
Lily 82-23	52			17.4			73		Y	Y		Centrachid - Fishing Pier
Little Carnelian 82-14	162	565	3.5:1	21.3	10.7	5,686			Y	N	1.7	
Little Johanna 62-58	35			12.0			67		N	N		
Long(ap val)19-22	36			3.5			100		N	N		
Long(Maht) 82-130	48			7.7			92		Y	N		
Long (May)82-30	88			3.7			100		N	Y		
Long (P.S.) 82-118	62	2,060	33:1	10.4	3.6	744	55		Y	N		
Long(Still) 82-21	71			6.7			96		N	N		
Long (Wash) 82-68	35	381	11:1	2.1	1.1	126	100		N	N		
Loon 82-15	64	407	6.4:1	4.9	2.4	206	100		N	N		
Lotus 10-6	246	1,033	4:1	8.8	4.3	3,500	74		Y	Y		
Louise 82-25	48	616	13:1	3.7	1.8	283	100		N	N		
MacDonald Pnd 82-62	12			2.7			100		N	N		
Maple Marsh 82-38	38	148	4:1	3.4	1.7	212	100		N	N		
Maria 10-58	169			1.0			100		Y	N		
Marion 19-26	560			6.4			81		Y	Y		
Markgrafs 82-89	46	413	10:1	2.4			100		N	N	2.6	Rearing
Markley 70-21	27			3.7			100		N	N		
McDonald 82-10	54	1,051	19:1	3.7	1.8	324	100		N	N		
McKusick 82-20	46			4.7			100		N	N	1.6	
Meadow 27-57	11	121	11:1	1.2			100		N	N	0.7	

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
Mergen's 82-482	12	1,383	115:1	1.3			100		N	N		
Miller 10-29	145	16,701	115:1	4.3	3.1	1,479	100		N	N		
Mitchell 27-70	112			5.8			97		N	Y		
Moody 13-23	35			14.6			63		Y	N		
Mud 82-26-02	62	899	15:1	2.1	1.1	224	100		N	N		
North Twin 82-18	69	187	3:1	1.8	0.9	207	100		N	N		
Northwood 27-627	15	1,341	89:1	1.5	0.8	41	100		N	N		
Oak 10-93	339			3.4			100		N	N		
O'Connor 82-2	38									N		
Olson 82-103	89	200	2:1	4.5	2.1	623	100		N	Y		
Oneka 82-140	381			2.1	1.2	1,524	100		N	N		Wildlife
Orchard 19-31	250	2,012	8:1	10.0	3.0	2,500	75		Y	Y		Centrachid
Pamela 27-675	18			1.5			100		N	N		
Parkers 27-107	97	950	10:1	11.3	3.7	1,164	70		Y	Y		
Peltier 2-4	174	68,082	391:1	4.9	2.1	3,255	100		N	Y		Gamefish
Pike(m.g.) 27-111	59	919	16:1	11.9	2.0	395	95		Y	Y	1.5	Centrachid
Pike(ramsy)62-69	35			4.9	2.1	252	100		N	N		Gamefish
Pike (scott) 70-76	57	1,991	35:1	2.7			100		N	N		
Pine Tree 82-122	174			7.9	3.0	1,740	91		Y	N		Centrachid
Powers 82-92	57	1,238	22:1	12.5			57	2	Y	N	1.8	Centrachid
Prior(lower)70-26	827	19,560	24:1	18.3	4.1	11,120	46	1	Y	Y		Centrachid

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
Prior(upper)70-72	340	16,460	48:1	15.2	3.1	3,460	93	2	Y	Y		Centrachid
Region Prk 82-87	16	600	38:1	5.8			100		N	N		
Reitz 10-52	79	3,711	47:1	11.0	4.0	1,027	58		Y	Y		
Riley 10-2	297	4,796	16:1	15.0	6.6	6,429	34		Y	Y	2.9	
Ryan 27-58	20	5,510	157:1	10.7	64.8	312	56		Y	N	0.6	
Sand 82-67	46			5.5	2.4	368	46	2		N	1.8	
Schmidt 27-102	37	190	4:1	9.1	1.5	207	92		Y	N	1.6	
School 13-57	48											
Schutz 10-18	105	943	9:1	15.0	6.0	2,100	27		Y	N		
Schroeder Pnd 82-301				3.0			100		N	N		
Seidl's 19-95	14	415	30:1	5.0			100	5	N	N		Rearing
Shaver 27-86	11									N		
Shields 82-162	27			8.2			85		Y	N	0.8	
Silver 82-16	98	455	4.6:1	3.4	1.7	549	100		N	N		
South Oak 27-661										N		
South Rice 27-645	3.2	63	20:1	2.5	0.5	5.4	100		N	N		
S. School Section 82-151	125			8.0			41					
South Twin 82-19	54	63	1.2:1	4.0	2.0	356	100		N	N		
Spring (Scott)70-54	630	13,500	21:1	11.3	5.6	11,500	50	2	Y	Y	5.0	
Square 82-46	193	782	4:1	20.7	9.0	5,694	65	5	Y	Y	2.2	Stocked w/Trout
Staples 82-28	24	127	5.3:1	4.3	2.1	165	100		N	N		

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
St. Croix 82-1	8,600	4,918,790		23.8					Y	Y		
St. Joe 10-11	14			15.9			46		Y	Y		
Sunfish 82-107	50	526	11:1							N		
Sunnybrook 82-133	16	630	39:1	6.1	2.0	104			Y	N		
Sunset 82-153	124			5.2			100		N	N	2.3	Gamefish
Sunset Pnd19-451	60			3.7			100		N	N	1.9	
Swede 10-95	376			4.0			100		N	Y		
Sweeney 27-35	66	2,400	36:1	8.0	3.6	790	52		Y	N		Panfish
Tamarack 10-10	24			20.0			41		Y	N		
Terrapin 82-31	86			4.6			100		N	N		
Turtle 82-36	44	699	16:1	2.4	1.2	172	100		N	N		
Twin(Bnsv)19-28	11						100					
Twin(U)(b.p.)27-42	137	3,657	31:1	2.4	0.9	397	100		Y	N	2.8	Centrachid
Twin(M)(cry)27-42	69	4,053	72:1	14.0	4.9	918	57		Y	Y	1.4	Centrachid
Twin(L)(rob)27-42	46	5,322	176:1	6.7	2.3	340	83		Y	Y	1.2	Centrachid
Twin(StLP) 27-656										N		
Valentine 62-71	60	2,237	37:1	4.0	1.5	300	100		N			
Valley 19-348	15	117	8:1	3.2			100	1	N	N		
Virginia 10-18	110	772	7:1	10.4	3.3	1,210	88		Y	Y		
Waconia 10-59	3,000	7,880	4:1	11.3	4.0	38,632	53		Y	Y	6.8	Centrachid
West Boot 82-44	110	209	2:1	11.9	5.9	2,090	56		Y	Y		

Lake DNR #	Surface Area(ac)	Watershed Area(ac)	Ratio	Max Depth(m)	Mean Depth(m)	Volume (ac-ft)	% Littoral	# Inlets	Termo- cline?	Public Access	Shr Length (miles)	DNR Classification
West Lakeland 82-488	27	1,139	347:1						N	N		
Westwood 27-711	41			2.0			100		N	N		
Wilmes 82-90	41	2,247	55:1	5.5						Y	1.3	
Windsor 27-82	14									N		
Winkler 10-66	129	2,758	21:1									
Wood(Brns)19-24	9	157	17:1	4.5			100	1	N	N		Panfish

APPENDIX B

2005 Volunteer Lake Monitors

<u>WMO/WD/City</u>	<u>Lake</u>	<u>DNR #</u>	<u>Volunteer</u>
Anoka Co. Parks	Cenaiko	02-0654	Anoka Co. Parks
	Highland	02-0079	Anoka Co. Parks
	Island	02-0022	Anoka Co. Parks
Apple Valley	Cobblestone	19-0456	City of Apple Valley
	Farquhar	19-0023	Rick Bruneau
	Long (Apple Valley)	19-0022	Cherie Serie/Al Kettlekamp
Bassett Creek WMO	Northwood	27-0627	Steve Bur
	Parkers	27-0107	Bob Videen
	South Rice	27-0645	Steve Streff
	Sweeney (Site-1)	27-0035	Dave Hanson
	Westwood	27-0711	Carlson Family
Black Dog WMO	Crystal	19-0027	Arnett Family
	Keller	19-0025	Glen Gramse
	Kingsley	19-0030	Green Family
	Lac Lavon	19-0446	Wally Shaver
	Orchard	19-0031	Tom/Dorothy Goodwin
Browns Creek WMO	Benz	82-0120	Washington Co. SWCD
	Goggins	82-0077	Washington Co. SWCD
	Kismet	82-0333	Washington Co. SWCD
	Long (Stillwater)	82-0021	Washington Co. SWCD
	South School Section	82-0151	Washington Co. SWCD
Burnsville	Alimagnet	19-0021	John Ritter
	Earley	19-0033	Mary Oaster
	Twin (Burnsville)	19-0028	Bernie DeMaster
	Wood	19-0024	David Bess
Carnelian-Marine WD	Barker	82-0076	Washington Co. SWCD
	Bass	82-0035	Washington Co. SWCD
	Big Carnelian	82-0049	Washington Co. SWCD
	Big Marine	82-0052	Washington Co. SWCD
	Carol	82-0017	Washington Co. SWCD
	East Boot	82-0034	Washington Co. SWCD
	Fish	82-0064	Washington Co. SWCD
	German	82-0056	Washington Co. SWCD
	Herber's Pond	82-0015-01	Washington Co. SWCD
	Jellum's (Site-1)	82-0052-01	Washington Co. SWCD
	Little Carnelian	82-0014	Washington Co. SWCD
	Long	82-0068	Washington Co. SWCD
	Loon	82-0015-02	Washington Co. SWCD
	Louise	82-0025	Washington Co. SWCD
	MacDonald's Pond	82-0062	Washington Co. SWCD
	Maple Marsh	82-0038	Washington Co. SWCD
	Mud	82-0026-02	Washington Co. SWCD
	North Twin	82-0018	Washington Co. SWCD
	Schroeder's Pond	82-0301	Washington Co. SWCD
	Silver	82-0016	Washington Co. SWCD
	South Twin	82-0019	Washington Co. SWCD
	Staples	82-0028	Washington Co. SWCD
	Turtle	82-0036	Washington Co. SWCD
	West Boot	82-0044	Washington Co. SWCD

<u>WMO/WD/City</u>	<u>Lake</u>	<u>DNR #</u>	<u>Volunteer</u>
Carver Co.	Bavaria	10-0019	John Ryski
	Benton	10-0069	Carver Co. Env. Services
	Brickyard	10-0025	Carver Co. Env. Services
	Burandt	10-0084	Don Westermann
	Campbell	10-0127	Carver Co. Env. Services
	Courthouse	10-0005	Carver Co. Env. Services
	Eagle	10-0121	Carver Co. Env. Services
	Fireman's	10-0226	Carver Co. Env. Services
	Gaystock	10-0031	Carver Co. Env. Services
	Goose (Waconia)	10-0089	Carver Co. Env. Services
	Hazeltine	10-0014	Carver Co. Env. Services
	Hydes	10-0088	Carver Co. Env. Services
	Maria	10-0058	Carver Co. Env. Services
	Miller	10-0029	Carver Co. Env. Services
	Oak	10-0093	Ed Foley
	Swede	10-0095	Wayne Hubin
	Waconia	10-0059	Carver Co. Env. Services
	Winkler	10-0066	Carver Co. Env. Services
Chanhassen	Lotus	10-0006	Shelly Strohmaier
	Riley	10-0002	David Florenzano
	St. Joe	10-0011	Sue Morgan/Linda Scott
Comfort Lk-Forest Lk WD	Big Comfort	13-0053	Washington Co. SWCD
	Birch	13-0042	Washington Co. SWCD
	Bone	82-0054	Washington Co. SWCD
	Forest-West	82-0159	Washington Co. SWCD
	Halfbreed (Sylvan)	82-0080	Washington Co. SWCD
	Moody	13-0023	Washington Co. SWCD
	School	13-0057	Washington Co. SWCD
	Shields	82-0162	Washington Co. SWCD
Conservation League/Edina	Cornelia	27-0028-01	Conservation League of Edina
	Edina	27-0029	Conservation League of Edina
	Pamela	27-0675	Conservation League of Edina
Eden Prairie	Mitchell	27-0070	Gordon Warner
Elm Creek	Henry	27-0175	Tom Hoverson
IGH/SSP	Seidl's	19-0025	Harv Bartz
Lakeville	East	19-0349	City of Lakeville
	Lee	19-0029	David Zook
	Marion	19-0026	Wally and Ardyce Potter
	Valley	19-0348	City of Lakeville
Lower St. Croix WMO	O'Connor	82-0002	Jeff Keene
Marine/St.Croix WD	Goose (New Scandia)	82-0059	Washington Co. SWCD
	Hay	82-0065	Washington Co. SWCD
	Long (May)	82-0030	Washington Co. SWCD
	Sand	82-0067	Washington Co. SWCD
	Square	82-0046	Washington Co. SWCD
	Terrapin	82-0031	Washington Co. SWCD
Middle St. Croix WMO	McKusick	82-0020	Washington Co. SWCD

<u>WMO/WD/City</u>	<u>Lake</u>	<u>DNR #</u>	<u>Volunteer</u>
Minnehaha Creek WD	Schutz	10-0018	Mike Shouldice
	Tamarack	10-0010	Mike Shouldice
	Virginia	10-0015	Renay Leone
Minnetonka	Shaver	27-0086	Peter Davis and Family
	Windsor	27-0082	Sue Proudfoot/Karen Mueller
Pioneer-Sarah WD	Hafften	27-0199	Todd Fellman/Jim Van Someren
Prior Lake-Spring Lake WD	Cates	70-0018	Tom Sletta
	Fish	70-0069	Steve Pierson
	Pike (Site-1)	70-0076	David/Mona Hanson
	Prior (Lower) (Site-1)	70-0026-01	Walt Burris
	Prior (Upper) (Site-1)	70-0072-01	Madison Groves
	Spring	70-0054	Bill Tisdell
Rice Creek WD	George Watch	02-0005	Wargo Nature Center
	Golden	02-0045	City of Circle Pines
	Hart	02-0081	Ray Muno
	Langton (Site-1)	62-0049-01	Yul Yost
	Langton (Site-2)	62-0049-02	Yul Yost
	Langton (North Basin)	62-0204	Yul Yost
	Little Johanna	62-0058	Jason Johnson
	Long (Mahtomedi)	82-0130	Kitty Francy-Payton
	Peltier	02-0004	Wayne LeBlanc
	Pike	62-0069	Philip Goodrich
	Pine Tree	82-0122	Gene Berwald
	Sunset	82-00153	Diane and Bob Coderre
	Valentine	62-0071	Bob Kistler
St. Croix Basin Planning	Lake St. Croix(Upper Pool S-2)	82-0001	Jim and Roberta Harper
	Lake St. CroixMid Pool S-3)	82-0001	Cecilia and Harry Martin
	Lake St. Croix(Mid Pool S-5)	82-0001	Richard and Sheryl Lindhom
	Lake St. Croix(Lower Pool S-6)	82-0001	Rick Meierotto
	Lake St. Croix(Lower Pool S-7)	82-0001	Carpenter Nature Center
St. Louis Park	Bass	27-0015	Jason Westrum
	Cobblecrest	27-0053	Jim and Gramham Kellogg
	Twin (St. Louis Pk)	27-0656	Ed Voyles
Schmidt Lake Assoc.	Schmidt	27-0102	Dale Wahlstrom
Shakopee	Dean	70-0074	Gerlach Family
Shingle Creek WMC	Bass	27-0098	Marvin Groth
	Crystal (Robbinsdale)	27-0034	Wayne Sicora
	Eagle	27-0111-01	Sharon Collins
	Meadow	27-0057	Diane Stauner
	Twin (Middle)	27-0042-02	Roni Brunner/Bob Hill
	Twin (Lower)	27-0042-03	Roni Brunner/Bob Hill
South Washington WD	Armstrong	82-0116-02	Washington Co. SWCD
	Powers	82-0092	Washington Co. SWCD
	Regional Park	82-0087	Washington Co. SWCD
Stillwater	Lily	82-0023	Washington Co. SWCD

<u>WMO/WD/City</u>	<u>Lake</u>	<u>DNR #</u>	<u>Volunteer</u>
Valley Branch WD	Cloverdale	82-0009	Kevin Bjork
	DeMontreville	82-0101	Washington Co. SWCD
	Downs	82-0110	Wesley Sly Family
	Edith	82-0004	Court Storey
	Elmo	82-0106	Wendy Griffin
	Goetschel	82-0313	John Longtin
	Jane	82-0104	Chuck Taylor
	Klawitter	82-0368	Bonnie Jurand
	Legion Pond	82-0462	Winkels Family
	Long (Pine Springs)	82-0118	Bill Feely/Jim Malkowski
	McDonald	82-0010	Steve Groves
	Mergen's	82-0482	Chris Moosbrugger
	Olson	82-0103	Washington Co. SWCD
	Sunfish	82-0107	Tom Hilpish
	Sunnybrook	82-0133	Arnie Johnson
Woodbury	Colby	82-0094	Hvass Family
	La	82-0097	City of Woodbury
	Markgrafs	82-0089	Terry Riley
	Wilmes	82-0090	Bill Aamodt

APPENDIX C
Lakes Sampled either by Metropolitan Council or Volunteer Programs, 1980 - 2005
(Numbers indicate sampling visits per year, while ^v denotes volunteer monitoring)

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Alimagnet	19-21																^v 12	^v 10	^v 10	^v 10	^v 10	^v 10	^v 8	^v 9	^v 12	^v 10	^v 10
Ann	10-12						5				13													13			
Armstrong	82-116 -02																			^v 15	^v 10	^v 13	^v 14	^v 15	^v 14	^v 14	^v 14
Assumption	10-63																				^v 1						
Auburn-East	10-44				10																						
Auburn-West	10-44				10			17	18				12			13											
Aue	10-28																				^v 1						
Bald Eagle (Site-1)	62-2	4	5		5																					13	13
Bald Eagle (Site-2)	62-2																									13	13
Barnes	10-109																				^v 1						
Barker	82-96																					^v 5	^v 5	^v 7	^v 7	^v 7	^v 7
Bass	27-98	4														^v 16			^v 15		^v 15		^v 13		^v 9		^v 15
Bass (St. Louis Park)	27-15																							^v 12			^v 12
Bass (Washington Co.)	82-35																					^v 14	^v 5	^v 7	^v 7	^v 7	^v 7
Battle Creek	82-91														^v 14	^v 13	^v 11	^v 13									
Bavaria	10-19				5			17	18							13		^v 11	^v 12	^v 15	^v 12	^v 14	^v 14	^v 14	^v 19	^v 16	^v 18
Benton	10-69																				^v 13	^v 14	^v 14		^v 15		^v 14
Benz	82-120																			^v 8							^v 14
Berliner	10-103																				^v 1						
Big Carnelian	82-49					5					13					13			13			^v 14	^v 7	^v 14	^v 14	^v 14	^v 14

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Big Comfort	13-53																			^3		^14	^14	^14	^14	^14	^13
Big Marine	82-52	4	5			5					13					13			13			^14	^7	^14	^14	^14	^14
Birch	13-42																										^10
Birch	62-24	2																									^14
Bluebill Bay	19-449																		^8								
Bone	82-54					5					13				^7		^14		^14	^14	^14		^14	^14	^14	^14	^14
Brand	10-110																				^1						
Braunworth	10-107																				^1						
Brickyard	10-225																							^14	^13	^14	^14
Bryant	27-67	2	5	16		5					13	13	12														
Burandt	10-84																				^7	^13	^9			^18	^22
Bush	27-47					5									13	13					13		13			13	
Byllesby	19-6														^14	^14	^13										
Calhoun	27-31		5			5																					
Campbell	10-127																				^2	^14		^10			^14
Carol	82-17																					^5	^5	^7	^7	^7	^7
Carver	82-166									20					^15	^15	^16	^9									
Cates	70-18																							^14	^13	^15	^13
Cedar (Minneapolis)	27-39					5																					
Cedar (Scott Co.)	70-91	4	5			5						13			14					13			13				13
Cedar Island	27-119																^13					^13		^11			
Cenaiko	2-654																		^12	^11	^13	^11	^13	^12	^12	^14	^14

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Centerville	2-6	4	5		5																	13	13/v 4	v1	13	13	
Charley	62-62						5																				
Christmas	27-137	4	5				5												13	13	13			13	13		
Chub	19-20	2													v14	v14	v11										
Clear (Forest Lake)	82-163	4				5						13			v11	v12	v12	v11	v10	v11	v10	v9	v12	v12	v12	v6	
Cleary	70-22					5																					
Cloverdale	82-9																						v10	v10	v11	v13	v12
Cobblecrest	27-53																							v4		v14	v16
Cobblestone	19-456																										v14
Colby	82-94															v13	v14	v13	v13	v12	v12	v9	v10	v10	v10	v10	v6
Comfort	13-53																		v3								
Coon	2-42	4				5										13			13								
Cornelia	27-28																								v7		v11
Courthouse (Chaska)	10-5																	v2	v14	v13	v13	v14	v14	v14	v14	v14	v14
Cowley	27-169																	v12									
Crane	27-734														v9												
Crooked	2-84				5						13				v15	v15	v14	v14	v12	v14	v14						
Crystal (Burnsville)	19-27	2			5						13					13	13	13	13	13	v12	v10	v14	v15	v15	v15	v16
Crystal (Robbinsdale)	27-34							17	19	19						v15			v11				v8				v7
Crystal (Spring Lake)	70-61																		v12		v11						
Cynthia	70-52	2																									
Dan Patch	70-16																		v15								

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05	
Dean	70-74																							'7	'7	'6	'7	
Deeg	19-117																						'12					
Deep	62-18						5																					
DeMontreville	82-101	4				5							12		'15		14						13			13	'14	'7
Diamond (Dayton)	27-125	2														'13										13		
Downs	82-110																				'14		'9	'9	'6	'7	'9	
Dutch	27-181					5																						
Eagle (Maple Grove)	27-111-01	4			5			17	18				11		'15			'14	'14	'14		'6		'4			'6	
Eagle (Young America)	10-121	4	5				5											12		'15	'14	'14	'12	'14	'14	13	'14	
Eagle Point	82-109			2											'14													
Earley	19-33															'10	'11	'9	'10	'10	'9	'8	'6	'10	'9	'6	'7	
East	19-349																										'13	
East Boot	82-34																					'14	'14	'14	'14	'14	'14	
East Twin	2-133	2	5		5						13						13			13								
Edina	27-29																									'10	'10	
Edith	82-4																										'6	
Egg	82-147																						'3					
Elmo	82-106	4	5	16		5				19			12			'11											'9	
Farquhar	19-23	4														'15	'16	'14	'15		'15	'13	'11	'13	'14	'14	'15	
Fireman's	10-226																						'12	'14	'14	'14	'14	
Fish (Eagan)	19-57										13																	
Fish (Grant Twmsp)	82-137																							'5	'5	'4		

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Fish (Maple Grove)	27-118	4	5	16			5					13															
Fish (Scott County)	70-69	4				5						13					13		✓2	✓13	✓8	✓12	✓9	✓14	✓13	✓11	✓13
Fish (Washington Co.)	82-64																					✓5	✓14	✓7	✓7	✓7	✓7
Forest - East (3)	82-159	4				5						13			✓7			✓12						13			13
Forest - Middle (2)	82-159					5						13			✓7			✓12						13			13
Forest - West (1)	82-159					5						13			✓7			✓12	✓14	✓15	✓14	✓14	✓14	✓14	✓14	✓14	✓14
French	27-127																						✓11	✓10	✓7	✓7	
Gables	82-82																			✓8	✓5						
Gaystock	10-31																				✓2	✓14	✓14				✓14
George	2-91	4	5	16		5					13					13				13							
George Watch	2-5																	✓14	✓12	✓11	✓11	✓6	✓7	✓8	✓9	✓10	✓12
German	82-56																							✓7	✓7	✓7	✓7
Gervais	62-7						5																				
Goetschel	82-313																							✓11	✓9	✓4	✓15
Goggins	82-77																				✓13	✓14	✓14	✓14	✓14	✓14	✓14
Golden	2-45	2											12		14			✓13	✓11	✓15	✓13	✓13	✓12	✓11	✓11	✓10	✓11
Goose (Lakeville)	19-360																✓13	✓13									
Goose (New Scandia)	82-59															✓15	✓15	✓13	✓13	✓15						✓7	✓7
Goose (Waconia)	10-89																✓9	✓7	✓15	✓15	✓14	✓11	✓14	✓14	✓14	✓14	✓14
Grace	10-218																							✓11	✓14	✓14	
Grass	27-681																		✓12								
Hafften	27-199																					13	13			13	✓15

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Half Breed (Sylvan)	82-80														✓7			✓14		✓15	✓14	✓14	✓14	✓14	✓14	✓14	✓14
Ham	2-53					5									✓15	✓13		✓13	✓9	✓14							
Harriet	27-16					5																					
Hart	2-81																									✓6	✓4
Harvey	27-???																									✓10	
Haughey	27-187																							✓4			
Hay	82-65																			✓14	✓13	✓14	✓14	✓4	✓7	✓7	✓7
Hazeltine	10-14																				✓1	✓14	✓14				✓14
Henry	27-175																✓10										✓11
Herber's Pond	82-15-01																									✓14	✓14
Highland	2-79																				✓13	✓11	✓13	✓12	✓12	✓14	✓14
Holland	19-65				10	16	15			20					13							13					
Horseshoe (Wash. Co.)	82-74																				✓1						
Horseshoe (Dakota Co.)	19-32																✓11	✓10									
Hydes	10-88						5						12		13			12			✓11	✓4	✓9	✓14	✓15	✓14	✓14
Independence	27-176	4	5		5							13			✓14	✓15											
Isabelle	19-4															✓14											
Island (Linwood)	2-22				7																				✓12	✓14	✓14
Jane	82-104					5		17	18				12			✓12						13				✓15	✓13
Jellum's (Site-1)	82-52-01																					✓14	✓14	✓12	✓14	✓14	✓14
Jellum's (Site-2)	82-52-02																							✓11	✓11		
Johanna	62-78		5				5				13																

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Jonathan	10-217																							'13			
Josephine	62-57						5				13																
Jubert	27-165																					'11					
Keller (Burnsville)	19-25																	13	13	'13	'15	'14	'12	'13	'15	'15	'14
Keller (Maplewood)	62-10						5																				
Kingsley	19-30														5		'11	'10	'9			'14	'14	'15	'14	'15	'16
Kismet	82-333																			'14	'13	'14	'14	'14	'14	'14	'13
Klawitter	82-368																							'13	'13	'14	'13
Kohlman	62-6						5																				
La	82-97															'13	'11	'13	'11	'10	'10	'8	'6	'5	'6	'3	'13
Lac Lavon	19-446																		'11	'10	'10	'9	'2	'7	'12	'12	'12
Laddie	2-72	4													'13	'14	'12					'13	'13	'14	'10		
Langdon	27-182					5																					
Langton (Site-1)	62-49-01																										'14
Langton (Site-2)	62-49-02																										'14
Langton (North Basin)	62-204																										'14
Lee	19-29															'14	'15	'14	'13			'12	'13	'11	'9	'15	'9
Legion Pond	82-462																										'14
Libbs	27-85																									'10	
Lily	82-23																'15	'14	'14	'15	'13	'14	'14	'14	'7	'7	'7
Linwood	2-26	4	5		7						13					13			13								
Lippert	10-104																				'1						

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Little Carnelian	82-14																					'14	'7	'14	'14	'14	'14
Little Johanna	62-58																						'12	'16	'15	'8	'6
Little Long	27-179-02	4				5						13								13			13		13		
Long (Apple Valley)	19-22																		'16					'11	'13	'12	'15
Long (Carver Co.)	10-16																				'2		'13		'5		
Long (Mahtomedi)	82-130																								'11	'9	'12
Long (May Twnsp)	82-30														'14	'14	'14	'13	'14		'14	'14	'14	'14	'14	'7	'7
Long N (New Brighton)	62-67						5																				
Long S (New Brighton)	62-67						5																				
Long (Orono)	27-160				5																						
Long (Pine Springs)	82-118														'14										13	'15	'14
Long (Stillwater)	82-21																'14	'7		'14	'13	'14	'14	'14	'14	'14	'14
Long (Washington Co.)	82-68																					'5	'14	'7	'7	'7	'7
Loon	82-15																					'14	'14	'7	'7	'7	'7
Lost	27-103														'13												
Lotus	10-6						5					13									13	13			'5	'10	'8
Louise	82-25																					'5	'5	'7	'7	'7	'7
Lucy	10-7						5																				
MacDonald's Pond	82-1501																									'14	'14
Magda	27-65																				'14	'13			'11		
Maple Marsh	82-38																					'5	'5	'7	'7	'7	'7
Marcott (site 1)	19-263																'15										

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Marcott (site 2)	19-41																✓15	✓13	✓10	✓10	✓12	✓10	✓6	✓5			
Maria	10-58																				✓2	✓14	✓14				✓13
Marion (Lakeville)	19-26	2	5		5						13					✓15					✓15	✓14	✓13	✓14	✓14	✓15	✓16
Markgrafs	82-89															✓15	✓11	✓12	✓10	✓15	✓10	✓10	✓9	✓13	✓14	✓14	✓14
Markley	70-21																		✓11	✓13	✓12	✓14	✓13	✓9	✓6	✓4	
Marsh	10-54																				✓1						
Marshan	2-7																	✓10	✓13	✓10	✓9	✓8	✓7				
Martin	2-34				7															13							
McCarrons	62-54					12	20	17	18	19	13	13	12		14	13	16	13			18	13	13	13		13	13
McDonald	82-10																				✓11		✓14	✓9	✓12	✓12	✓14
McDonough	19-76						5														13						
McKusick	82-20															✓14	✓14	✓14	✓14	✓14	✓13	✓14	✓14	✓14	✓14	✓14	✓14
McMahon (Carls)	70-50	2				5											13			13			13				13
Meadow	27-57																	✓12			✓12			✓9			✓10
Medicine	27-104	4	5		10							13	12														
Mergen's	82-482																					✓10			✓3	✓2	✓6
Meuwissen	10-70																				✓1						
Miller	10-29																	✓6	✓13		✓12	✓14	✓13	✓13	✓14	✓14	✓14
Minnetonka (Lower)	27-133	4	5																								
Minnetonka (Upper)	27-133	2	5																								
Minnewashta	10-9					5						13			13				13	13	13			13	13		
Mitchell	27-70																13				13	13			13	✓14	✓14

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Moody	13-23																										'14
Mooney	27-134														'14	'10											
Moore	2-75																				'14						
Mud	82-26-02																					'5	'5	'7	'7	'7	'7
Myers	10-68																				'1						
Nokomis	27-19	4				5																					
North Twin	82-18																					'5	'5	'7	'7	'7	'7
Northwood	27-627																					'12	'10	'13	'12	'12	'10
Oak	10-93																				'2		'14	'13	'12	'14	'14
O'Connor	82-2																										'8
O'Dowd	70-95					5										13			13			13		13			13
Olson	82-103												12		'15		14					13			13	'14	'7
Oneka	82-140																				'13	'11	'11	'9	'6	'5	
Orchard	19-31	4	5		5						13				13					13	'15	'13	'13		'14	'14	'14
Otter	2-3	2			5																						
Owasso	62-56	4			5																						
Pamela	27-675																										'10
Parkers	27-107	4										13					13				13	'12		'14	'15	'15	'15
Parley	10-42					5		17	18				12					12			13		13		13		
Patterson	10-86																				'2						
Peltier	2-4				5										'14	'16	'15	'14	'14	'13	'13	'14	'13	'17	'15	'15	'16
Phalen	62-13	4	5				5																				

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Pickereel	2-103	2															13										
Pierson	10-53	2	5		5						13						13						13	13	13		
Pike (Maple Grove)	27-111-02																	√14	√15	√13		√13					
Pike (Ramsey Co.)	62-69																				√14	√10	√14	√14	√14	√15	√15
Pike (Scott Co.) [Site-1]	70-76-1																		√9		√10	√9	√9	√11	√15	√15	√13
Pike {Scott Co.} [Site-2]	70-76-2																							√11			
Pine Tree	82-122						5								√14	√14	√16	√14	√15	√15	√13	√14	√9	√12	√7	√8	√12
Pleasant (New Prague)	70-98														13												
Pleasant (North Oaks)	62-46						5																				
Pomerleau	27-100																	√9			√10		√6		√3		
Powers	82-92															√12	√13	√13	√12	√9	√10	√8	√5	√7	√14	√14	√14
Prior (Lower) [Site-1]	70-26-1					5						13						13	√15	√14	√13	√9	√14	√16	√13	√12	√12
Prior (Lower) [Site-2]	70-26-2																			√14	√13	√9	√14	√15			
Prior (Upper) [Site-1]	70-72-1	4	5			5						13						13	√15	√14	√13	√9	√14	√12	√13	√10	√9
Prior {Upper} [Site-2]	70-72-2																							√12			
Raven	19-369																√13	√6	√8								
Rebecca	27-192				10	12	12																				
Red Rock	27-76																				12	13			13	13	
Regional Park	82-87																			√12	√14	√12	√13	√14	√15	√15	√14
Reitz	10-52						5						12		13						√15	√13	√7	√13	√14	√14	15
Reshanau	2-9	2																			√7	√1	√6				
Rice	10-78	2																			√1						

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Riley	10-2	2	5	16			5	17	18			13	12		13				13			13		13	14	15	14
Rutz	10-89																				1	14	14	14			
Ryan	27-58																	14		5		9		4	6		
Sand (New Scandia)	82-67														7	14	14	13						14	7	7	7
Sarah	27-191	4			5																						
Scheuble	10-85																					1					
Schmidt (Smith)	27-102																14			12		12	9			14	9
School	13-57																										14
Schroeder's Pond	82-301																									14	14
Schultz	19-75					5	5															13					
Schutz	10-18					5																6	10	6	8	9	11
Seidl's	19-95																15	14	14	15	16	14	14	15	8	14	14
Shaver	27-86																										14
Shields	82-162														6	14	14	13	13	14	14	14	14	14	14	14	14
Silver	82-16																					14	5	7	7	7	7
Simley	19-37																10	16	14	15	16	14	12	14			
Snail	62-73	4					5																				
South Oak	27-661																							12	15		
South Rice	27-645																					9	14	15	14	14	15
South School Section	82-151																14	7		14							14
South Twin	82-19																					5	5	7	7	7	7
Spring (Anoka Co.)	2-71																						11				

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Spring (Prior Lake)	70-54	4	5	16		5						13						13	✓12			✓6	✓11	✓13	✓14	✓14	✓13
Square	82-46	4	5	16	6	7	7				13				✓11	✓14	✓14	✓13	✓14	19	✓14	✓14	✓15	✓14	✓14	✓14	✓14
Staring	27-78	4					5										13				13		13			13	
Staples	82-28																					✓14	✓5	✓7	✓7	✓7	✓7
Steiger	10-45					12					13						13										
St. Croix (Upper Basin S-2)	82-1																										✓10
St. Croix (Mid Basin S-3)	82-1																										✓11
St. Croix (Mid Basin S-5)	82-1																										✓8
St. Croix (Lower Basin S-6)	82-1																										✓11
St. Croix (Lower Basin S-7)	82-1																										✓8
St. Joe	10-11																									✓17	✓8
Success	27-634																	✓10							✓11		
Sucker	62-28						5																				
Sullivan	2-80														✓14	✓14	✓15		✓15	✓14	✓13	✓11	✓11	✓12	✓12		
Sunfish	82-107																					✓10					✓13
Sunnybrook	82-133																				✓14		✓13	✓10	✓12	✓10	✓16
Sunset	82-153					5									✓14	✓14	✓12	✓13	✓16	✓12	✓10	✓13	✓13	✓18	✓20	✓15	✓17
Sunset Pond	19-451															✓14	✓14	✓14	✓12	✓10		✓13	✓11	✓10	✓12	✓11	
Swan	10-82																				✓1						
Swede	10-95	2																13					13	✓14	✓16	✓13	✓14
Sweeney (South) [Site-1]	27-35																					✓11	✓9	✓14	✓13	✓14	✓11
Sweeney (North) [Site-2]	27-35																					✓11	✓9				

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
Tamarack	10-10																						'10	'11	'12	'11	'11
Tanners	82-115	2								20					'14	'13	'12	'14									
Terrapin	82-31																									'7	'7
Thole	70-120-01					5										13			13			13		13			13
Thomas	19-67	2																									
Tiger	10-108																				'1						
Turtle	62-61	4	5		5																						
Turtle (Washington Co.)	82-36																					'5	'5	'7	'7	'7	'7
Twin (Burnsville)	19-28																				'6		'13	'11	'6	'2	'11
Twin-Lower (Robbinsd.)	27-42-03											12			'14			13		'5		13			'13		'8
Twin-Middle (Crystal)	27-42-02						5					12						13	'11		'13	13			'13		'8
Twin-Upper (Br. Center)	27-42-01										12				'14			11		'15		'11		'13		'14	
Twin-South (May Twnsp)	82-48																		'13	'13							
Twin (St. Louis Park)	27-656																							'12	'14	'14	'11
Vadnais	62-38						5																				
Valentine	62-71																						'14	'13	'12	'12	'9
Valley	19-348																'15	'14	'11		'8	'14	'14	'14	'14	'14	'13
Virginia	10-15																					'11	'12	'14	'12	'15	'13
Wabasso	62-82	4	5		5						12																
Waconia	10-59	4	5				5					13				'16	'13	'15	'17	'15	'14	'14	'14	'15	'14	12	'14
Wasserman	10-48				5			17	18							13			13	13	13			13	13		
Weaver	27-117				5			17	18																		

LAKE	ID #	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05
West Boot	82-44																					'14	'14	'14	'14	'14	'14
West Lakeland	82-488																					'2					
Westwood	27-711														'13							'15	'14	'10	'9	'7	'7
Whaletail (Site-1)	27-184-01																									13	13
Whaletail (Site-2)	27-184-02	4				5														13			13			13	13
White Bear	82-167	4	5			5																					
Wilmes	82-90															'14	'15	'14	'15	'15	'14	'13	'13	'10	'12	'12	'10
Windsor	27-82																									'12	'14
Winkler	10-66																				'8	'6	'6		'13		'14
Wolsfeld	27-157	4																									
Wood (Burnsville)	19-24																	'10	'14	'15	'15	'14	'13	'14	'14	'14	'14
Young America	10-105																				'1						
Zumbra	10-41					5						13												13			

Metropolitan Council Lake Report Questionnaire

1. The Council publishes a lake water quality report annually.
 - Are you familiar with the report? yes no
 - Have you used any of the data? yes no
 - Describe how you have used report findings.

2. These questions pertain to the content of the Lake Report.
 - What types of information from the report do you find most useful?

 - What portions of the report should be reduced, removed or are not useful?

 - What types of new data should be included in the report? Why?

 - Is the information in the report explained fully? Is it easy to understand?

 - Are the charts and graphs easy to understand?

 - Does the format add to understanding the information?

 - Other suggestions pertaining to writing, format design and graphics:

3. How often should the report be published? less frequently more frequently annually

4. Demographics of report users
 - Please circle occupation/employer

 - City County State

 - Government Official Government Staff Other (please specify)

Thank you for your help. Please return to:

Metropolitan Council
c/o Randy Anhorn
2400 Childs Road
St. Paul, MN 55106