This document is made available electronically by the Minnesota Legislative Reference Library as part of an ongoing digital archiving project. http://www.leg.state.mn.us/lrl/lrl.asp

Minnesota Statewide Quality Reporting and Measurement System:

Appendices to Minnesota Administrative Rules, Chapter 4654

Minnesota Department of Health

November 2010



Minnesota Statewide Quality Reporting and Measurement System:

Appendices to Minnesota Administrative Rules, Chapter 4654

November 2010

For more information, contact: Division of Health Policy Health Economics Program Minnesota Department of Health PO Box 64882 St. Paul, MN 55164-0882 www.health.state.mn.us

Phone: (651) 201-3550 Fax: (651) 201-5179 TDD: (651) 201-5797

Upon request, this material will be made available in an alternative format such as large print, Braille or cassette tape. Printed on recycled paper.

TABLE OF CONTENTS

INTRO	ODUCTION	5
APPEN	NDIX A – REQUIRED PHYSICIAN CLINIC QUALITY MEASURE DATA.	7
APPEN	NDIX B – REQUIRED HOSPITAL QUALITY MEASURE DATA	13
	NDIX C – REQUIRED AMBULATORY SURGICAL CENTER QUALITY SURES	34
APPEN	NDIX D – OTHER STANDARDIZED QUALITY MEASURES	37
APPEN	NDIX E – SUBMISSION SPECIFICATIONS	39
I.	SUBMISSION REQUIREMENTS FOR PHYSICIAN CLINICS	39
II.	SUBMISSION REQUIREMENTS FOR HOSPITALS	43
III.	SUBMISSION REQUIREMENTS FOR AMBULATORY SURGICAL CENTERS	47

INTRODUCTION

Minnesota Statutes 62U.02 requires the Commissioner of Health to establish standards for measuring health outcomes and develop a standardized set of measures to assess the quality of health care services offered by health care providers. In addition, Minnesota Statutes 62U.02 requires the Commissioner of Health to issue annual public reports on provider quality using a subset of measures from the standardized set of measures. The Department of Health has contracted with Minnesota Community Measurement (MNCM) to lead a consortium of organizations, including Stratis Health, the Minnesota Medical Association (MMA), the Minnesota Hospital Association (MHA), and the University of Minnesota School of Public Health, to assist in the completion of these tasks.

Measures that will be used for public reporting are identified in Appendices A, B and C. The standardized set of measures are defined in the body of the rule and include the measures identified in Appendices A, B, C, and D. The hospital measures in Appendix B and the ambulatory surgical center measures in Appendix C are defined by the referenced national quality organizations and will likely change over time as modified by the national quality organizations.

APPENDIX A REQUIRED PHYSICIAN CLINIC QUALITY MEASURE DATA

Data Elements Specification Information	Data Required for Reporting Beginning in Calendar Year 2011 an	nd Every Year Thereafter	
Diabetes Optimal diabetes care (ODC) composite These measures are used to assess the percent of adult patients who have type I or type II diabetes with optimally managed modifiable risk factors: • HbA1c (less than 8 percent) • Low-density lipoprotein (LDL) cholesterol (less than 100 mg/dL) • Blood pressure control (less than 140/90 mm Hg) • Daily aspirin use if patient has diagnosis of IVD or valid contraindication to aspirin • Documented tobacco free (Urgent Care Centers are not required to submit data on this measure.) Physician clinics must submit the following data for the optimal diabetes care measure and for each of the five component measures: Patient identification methodology Submit the following two data elements by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay:): • Denominator: Number of patients meeting the criteria for inclusion in the measure if submitting on the full population OR Number of patients in data submission if submitting a sample	Measure Name and Purpose	Data Elements	Specification Information
Optimal diabetes care (ODC) composite These measures are used to assess the percent of adult patients who have type I or type II diabetes with optimally managed modifiable risk factors: HbA1c (less than 8 percent) Low-density lipoprotein (LDL) cholesterol (less than 100 mg/dL) Blood pressure control (less than 140/90 mm Hg) Daily aspirin use if patient has diagnosis of IVD or valid contraindication to aspirin Documented tobacco free (Urgent Care Centers are not required to submit data on this measure.) Physician clinics must submit the following data for the optimal diabetes care measure and for each of the five component measures: Physician clinics must submit the following data for the optimal diabetes care: Dates of service: 01/01/2010 – 12/31/2010. Minnesota Community Measurement; Revised 08/13/2010. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/ healthreform OR Number of patients in data submitting a sample	Data Required for Reporting Beginning in January 2011 (2010 Da	ates of Service) and Every Year Thereafter	
These measures are used to assess the percent of adult patients who have type I or type II diabetes with optimally managed modifiable risk factors: I HbA1c (less than 8 percent) Low-density lipoprotein (LDL) cholesterol (less than 100 mg/dL) Blood pressure control (less than 140/90 mm Hg) Daily aspirin use if patient has diagnosis of IVD or valid contraindication to aspirin Documented tobacco free (Urgent Care Centers are not required to submit data on this measure.) the optimal diabetes care measure and for each of the five component measures: Patient identification methodology Submit the following two data elements by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay:): Number of patients meeting the criteria for inclusion in the measure if submitting on the full population OR Number of patients in data submission if submitting a sample	Diabetes		
 Numerator: Number of patients meeting the targets in the measure Denominator: Number of patients meeting the criteria for inclusion in the measure Number of patients meeting the exclusion criteria Calculated rate 	These measures are used to assess the percent of adult patients who have type I or type II diabetes with optimally managed modifiable risk factors: HbA1c (less than 8 percent) Low-density lipoprotein (LDL) cholesterol (less than 100 mg/dL) Blood pressure control (less than 140/90 mm Hg) Daily aspirin use if patient has diagnosis of IVD or valid contraindication to aspirin Documented tobacco free (Urgent Care Centers are not required to submit data on this	the optimal diabetes care measure and for each of the five component measures: Patient identification methodology Submit the following two data elements by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay:): Denominator: Number of patients meeting the criteria for inclusion in the measure if submitting on the full population OR Number of patients in data submission if submitting a sample Numerator: Number of patients meeting the targets in the measure Denominator: Number of patients meeting the criteria for inclusion in the measure Number of patients meeting the exclusion criteria	diabetes care: Dates of service: 01/01/2010 – 12/31/2010. Minnesota Community Measurement; Revised 08/13/2010. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/

Measure Name and Purpose	Data Elements	Specification Information
Data Required for Reporting Beginning in January 2011 (2010 Da	ates of Service) and Every Year Thereafter	
Cardiovascular Conditions		
Optimal vascular care (OVC) composite These measures are used to assess the percent of adult patients who have ischemic vascular disease (IVD) with optimally managed modifiable risk factors: Low-density lipoprotein (LDL) cholesterol (less than 100 mg/dL) Blood pressure (BP) control: For patients with IVD and diabetes, BP less than 140/90 mm Hg For all other IVD patients, BP less than 130/80 mm Hg Daily aspirin use or contraindication to aspirin Documented tobacco free (Urgent Care Centers are not required to submit data on this measure.)	Physician clinics must submit the following data for the optimal vascular care measure and for each of the four component measures: Patient identification methodology Submit the following two data elements by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay): Denominator: Number of patients meeting the criteria for inclusion in the measure if sumitting on the full population OR Denominator: Number of patients in data submission if submitting a sample Numerator: Number of patients meeting the targets in the measure	Specifications for optimal vascular care: Dates of service: 01/01/2010 – 12/31/2010. Minnesota Community Measurement; Revised 8/13/2010. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/healthreform
	 Denominator: Number of patients meeting the criteria for inclusion in the measure 	
	 Number of patients meeting the exclusion criteria 	
	■ Calculated rate	

Measure Name and Purpose	Data Elements	Specification Information
Data Required for Reporting Beginning in February 2011 (2010 Dates of Service) and Every Year Thereafter		
Behavioral Health Conditions		
Depression remission at six months This measure is used to assess the percent of adult patients who have major depression or dysthymia who have reached remission at six months (+/- 30 days) after being identified as having an initial PHQ-9 score greater than 9. Remission is identified as a PHQ-9 score less than 5. (Urgent Care Centers are not required to submit data on this measure.)	Physician clinics must submit the following data for the depression remission at six months measure: Patient identification methodology Submit the following two data elements by three bands of initial PHQ9 scores (10-14; 15-19; 20 and above): Denominator: Number of patients meeting the criteria for inclusion in the measure Numerator: Number of patients meeting the targets in the measure Number of patients meeting the exclusion criteria Number of patients for whom a follow up six month (+/- 30 days) PHQ9 assessment was not completed. Calculated rate	Specifications for depression: Minnesota Community Measurement; Revised 7/13/2010. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/ healthreform
Measure Name and Purpose	Data Elements	Specification Information

Measure Name and Purpose	Data Elements	Specification Information
Data Required for Reporting Beginning in July 2011 (July 1, 2010	- June 30, 2011 Dates of Service) and Every Year Th	ereafter
Respiratory Conditions		
Optimal asthma care These measures are used to assess the percent of pediatric and adult asthma patients who are receiving optimal care. Optimal care is	Physician clinics must submit the following data for the optimal asthma care measure and for each of the three component measures:	Specifications for optimal asthma care: Dates of service: 07/01/2010- 06/30/2011.

Measure Name and Purpose	Data Elements	Specification Information
Data Required for Reporting Beginning in July 2011 (July 1, 2010	- June 30, 2011 Dates of Service) and Every Year Th	ereafter
defined as: Asthma is well controlled Patient is not at increased risk of exacerbations Patient has a current written asthma action/management plan (Urgent Care Centers are not required to submit data on this measure.)	 Patient identification methodology Within two separate age bands, ages 5-17 and 18-50, submit the following two data elements by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay): Denominator: Number of patients meeting the criteria for inclusion in the measure if submitting on the full population OR Number of patients in data submission if submitting a sample (NOTE: One sample per age band is required for this measure.) Numerator: Number of patients meeting the targets in the measure Denominator: Number of patients meeting the criteria for inclusion in the measure Number of patients meeting the exclusion criteria Calculated rate 	Minnesota Community Measurement; April 2010. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/ healthreform
Preventive Care		
Colorectal cancer screening This measure is used to assess the percent of adult patients who are up to date with appropriate colorectal cancer screening. The	Physician clinics must submit the following data for the colorectal cancer screening measure: Patient identification methodology	Specifications for colorectal cancer screening: Dates of service: 07/01/2010 – 06/30/2011. Minnesota

Measure Name and Purpose	Data Elements	Specification Information
Data Required for Reporting Beginning in July 2011 (July 1, 2010	- June 30, 2011 Dates of Service) and Every Year Th	ereafter
 Colonoscopy within ten years Sigmoidoscopy within five year Stool Blood Tests (FOBT or FIT) within the reporting period (Urgent Care Centers are not required to submit data on this measure.) 	 Submit the following two data elements by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay): Denominator: Number of patients meeting the criteria for inclusion in the measure if submitting on the full population OR Number of patients in data submission if submitting a sample Numerator: Number of patients meeting the targets in the measure Denominator: Number of patients meeting the criteria for inclusion in the measure Number of patients meeting the exclusion criteria Calculated rate 	Community Measurement; Revised 5/27/2010. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/ healthreform

Measure Name and Purpose	Data Elements	Specification Information
Data Required for Reporting Beginning in February 2011 (2010 D	ates of Service) and Every Year Thereafter	
Health Information Technology (HIT)		
Health information technology (HIT) This survey is used to assess a medical group's adoption and use of Health Information Technology (HIT) in their clinical practice.	Internet-based survey as updated in 2011	Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/

Measure Name and Purpose	Data Elements	Specification Information
Data Required for Reporting Beginning in February 2011 (2010 Dates of Service) and Every Year Thereafter		
		healthreform

Data Required for Reporting Beginning in Calendar Year 2013 and Every Year Thereafter		
Measure Name and Purpose	Data Elements	Specification Information
Data Required for Reporting Beginning in February 2013 (2012 D	ates of Service) and Every Year Thereafter	
Patient Experience of Care		
Patient experience of care This survey will be used to assess adult patient experience of care. MDH will require use of the Clinician and Group Consumer Assessment of Healthcare Providers and Systems (CG-CAHPS) visit specific tool.		Specifications for patient experience: Survey period 09/01/2012-11/30/2012. Minnesota Community Measurement; Revised 10/4/10. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/healthreform

APPENDIX B REQUIRED HOSPITAL QUALITY MEASURE DATA

Data Required for Reporting Beginning in Calendar Year 2011 and Every Year Thereafter		
Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and	Every Year Thereafter	
Hospital Compare Quality Measures, Centers for Medicare & Me	dicaid Services (CMS) and The Joint Commission	
Acute myocardial infarction (AMI) – Acute myocardial infarction (AMI) / heart attack process of care measures for applicable hospital discharge dates The hospital process of care measures include the following measures related to heart attack care: Aspirin at arrival (AMI-1) – This measure is used to assess the percent of acute myocardial infarction (AMI) patients who received aspirin within 24 hours before or after hospital arrival. Aspirin prescribed at discharge (AMI-2) – This measure is used to assess the percent of acute myocardial infarction (AMI) patients who are prescribed aspirin at hospital discharge. ACEI or ARB for LVSD (AMI-3) – This measure is used to assess the percent of acute myocardial infarction (AMI) patients with left ventricular systolic dysfunction (LVSD) who are prescribed an ACEI or ARB at hospital discharge. For purposes of this measure, LVSD is defined as chart documentation of a left ventricular ejection fraction (LVEF) less than 40% or a narrative description of left ventricular systolic (LVS) function consistent with moderate or severe systolic dysfunction. Adult smoking cessation advice/counseling (AMI-4) – This measure is used to assess the percent of acute myocardial infarction (AMI) patients with a history of smoking cigarettes, who are given smoking cessation advice or	Hospitals must submit data for each of the hospital compare acute myocardial infarction (AMI) / heart attack process of care quality measures. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate	Specifications manual for national hospital inpatient quality measures, version 3.2b, discharges 10-01-10 (4Q10) through 03-31-11 (1Q11). Centers for Medicare & Medicaid Services (CMS), The Joint Commission; April 2010 or as updated. Measure specifications can be found on the Centers for Medicare & Medicaid Services (CMS), QualityNet website http://qualitynet.org

Measure Name and Purpose	Data Elements	Specification Information	
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter			
counseling during hospital stay. For purposes of this measure, a smoker is defined as someone who has smoked cigarettes anytime during the year prior to hospital arrival.			
 Beta-blocker prescribed at discharge (AMI-5) – This measure is used to assess the percent of acute myocardial infarction (AMI) patients who are prescribed a beta-blocker at hospital discharge. 			
• Fibrinolytic therapy received within 30 minutes of hospital arrival (AMI-7a) – This measure is used to assess the percent of acute myocardial infarction (AMI) patients receiving fibrinolytic therapy during the hospital stay and having a time from hospital arrival to fibrinolysis of 30 minutes or less.			
Primary PCI received within 90 minutes of hospital arrival (AMI-8a) – This measure is used to assess the percent of acute myocardial infarction (AMI) patients receiving primary percutaneous coronary intervention (PCI) during the hospital stay with a time from hospital arrival to PCI of 90 minutes or less.			
Heart failure (HF) – All heart failure (HF) process of care measures for applicable hospital discharge dates The hospital process of care measures include the following measures related to heart failure care: Discharge instructions (HF-1) – This measure is used to assess the percent of heart failure patients discharged home	Hospitals must submit data for each of the hospital compare heart failure process of care quality measures. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality	Specifications manual for national hospital inpatient quality measures, version 3.2b, discharges 10-01-10 (4Q10) through 03-31-11 (1Q11). Centers for Medicare & Medicaid Services (CMS),	
with written discharge instructions or educational material given to patient or caregiver at discharge or during hospital stay addressing all of the following: activity level, diet, discharge medications, follow-up appointment, weight monitoring, and what to do if symptoms worsen. • Evaluation of LVS function (HF-2) – This measure is used to	 Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate 	 Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate The Joint Commit 2010 or as update Measure specification found on the Cen Medicare & Medicare	The Joint Commission; April 2010 or as updated. Measure specifications can be found on the Centers for Medicare & Medicaid Services (CMS), QualityNet
assess the percent of heart failure patients with		Services (Civis), QuantyNet	

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
documentation in the hospital record that left ventricular systolic (LVS) function was evaluated before arrival, during hospitalization, or is planned for after discharge. ACEI or ARB for LVSD (HF-3) – This measure is used to assess the percent of heart failure patients with left ventricular systolic dysfunction (LVSD) who are prescribed an ACEI or ARB at hospital discharge. For purposes of this measure, LVSD is defined as chart documentation of a left ventricular ejection fraction (LVEF) less than 40% or a narrative description of left ventricular systolic (LVS) function consistent with moderate or severe systolic dysfunction. Adult smoking cessation advice/counseling (HF-4) – This measure is used to assess the percent of heart failure patients with a history of smoking cigarettes, who are given smoking cessation advice or counseling during hospital stay. For purposes of this measure, a smoker is defined as someone who has smoked cigarettes anytime during the year prior to hospital arrival.		website http://qualitynet.org
 Pneumonia (PN) – Pneumonia (PN) process of care measures for applicable hospital discharge dates The hospital process of care measures include the following measures related to pneumonia care: Pneumococcal vaccination (PN-2) – This measure is used to assess the percent of pneumonia patients, age 65 and older, who were screened for pneumococcal vaccine status and were administered the vaccine prior to discharge, if indicated. Blood cultures performed in the emergency department prior to initial antibiotic received in hospital (PN-3b) – This measure is used to assess the percent of pneumonia patients whose initial emergency room blood culture specimen was 	Hospitals must submit data for each of the hospital compare pneumonia process of care quality measures. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate	Specifications manual for national hospital inpatient quality measures, version 3.2b, discharges 10-01-10 (4Q10) through 03-31-11 (1Q11). Centers for Medicare & Medicaid Services (CMS), The Joint Commission; April 2010 or as updated. Measure specifications can be found on the Centers for Medicare & Medicaid Services (CMS), QualityNet

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
collected prior to first hospital dose of antibiotics. This measure focuses on the treatment provided to Emergency Department patients prior to admission orders.		website http://qualitynet.org
• Adult smoking cessation advice/counseling (PN-4) – This measure is used to assess the percent of pneumonia patients with a history of smoking cigarettes who are given smoking cessation advice or counseling during hospital stay. For purposes of this measure, a smoker is defined as someone who has smoked cigarettes anytime during the year prior to hospital arrival.		
 Initial antibiotic received within 6 hours of hospital arrival (PN-5c) – This measure is used to assess the percent of pneumonia patients who received their first dose of antibiotics within 6 hours after arrival at the hospital. 		
■ Initial antibiotic selection for community-acquired pneumonia (CAP) in immunocompetent patients (PN-6) — This measure is used to assess the percent of immunocompetent patients with Community-Acquired Pneumonia who receive an initial antibiotic regimen during the first 24 hours that is consistent with current guidelines.		
 Influenza vaccination (PN-7) – This measure is used to assess the percent of pneumonia patients age 50 years and older, hospitalized during October, November, December, January, February, or March who were screened for influenza vaccine status and were vaccinated prior to discharge, if indicated. 		
Surgical care improvement project (SCIP) – All surgical care improvement project (SCIP) process of care measures for applicable hospital discharge dates The hospital process of care measures include the following measures related to surgical care improvement project: • Prophylactic antibiotic received within one hour prior to	Hospitals must submit data for each of the hospital compare surgical care improvement project (SCIP) process of care quality measures. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality	Specifications manual for national hospital inpatient quality measures, version 3.2b, discharges 10-01-10 (4Q10) through 03-31-11 (1Q11). Centers for Medicare & Medicaid Services (CMS),

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
surgical incision * (SCIP-Inf-1) – This measure set is used to assess the percent of surgical patients with prophylactic antibiotics initiated within one hour prior to surgical incision. *Patients who received vancomycin or a fluoroquinolone for prophylactic antibiotics should have the antibiotics initiated within two hours prior to surgical incision. Due to the longer infusion time required for vancomycin or a fluoroquinolone, it is acceptable to start these antibiotics within two hours prior to incision time.	 Mumerator: Number of patients meeting the targets in each of the quality measures Calculated rate 	The Joint Commission; April 2010 or as updated. Measure specifications can be found on the Centers for Medicare & Medicaid Services (CMS), QualityNet website http://qualitynet.org
 Prophylactic antibiotic selection for surgical patients (SCIP-Inf-2) – This measure set is used to assess the percent of surgical patients who received prophylactic antibiotics consistent with current guidelines (specific to each type of surgical procedure). 		
Prophylactic antibiotics discontinued within 24 hours after surgery end time * (SCIP-Inf-3) – This measure set is used to assess the percent of surgical patients whose prophylactic antibiotics were discontinued within 24 hours after surgery end time (within 48 hours for coronary artery bypass graft (CABG) or other cardiac surgery) *The Society of Thoracic Surgeons (STS) Practice Guideline for Antibiotic Prophylaxis in Cardiac Surgery (2006) indicates that there is no reason to extend antibiotics beyond 48 hours for cardiac surgery and very explicitly states that antibiotics should not be extended beyond 48 hours even with tubes and drains in place for cardiac surgery.		
■ Cardiac surgery patients with controlled 6 a.m. postoperative blood glucose (SCIP-Inf-4) — This measure is used to assess the percent of cardiac surgery patients with controlled 6 A.M. blood glucose (≤ 200 mg/dL) on postoperative day one (POD 1) and postoperative day two (POD 2) with Surgery End Date being postoperative day zero (POD 0).		
 Surgery patients with appropriate hair removal (SCIP-Inf-6) – This measure is used to assess the percent of surgery patients 		

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
with appropriate surgical site hair removal. No hair removal, or hair removal with clippers or depilatory is considered appropriate. Shaving is considered inappropriate.		
Surgery patients on beta-blocker therapy prior to arrival who received a beta-blocker during the perioperative period (SCIP-Card-2) – This measure is used to assess the percent of surgery patients on beta-blocker therapy prior to arrival who received a beta-blocker during the perioperative period. The perioperative period for the SCIP Cardiac measures is defined as 24 hours prior to surgical incision through discharge from post-anesthesia care/recovery area.		
 Surgery patients with recommended venous thromboembolism prophylaxis ordered (SCIP-VTE-1) – This measure is used to assess the percent of surgery patients with recommended venous thromboembolism (VTE) prophylaxis ordered anytime from hospital arrival to 24 hours after Surgery End Time. 		
 Surgery patients who received appropriate venous thromboembolism prophylaxis within 24 hours prior to surgery to 24 hours after surgery (SCIP-VTE-2) – This measure is used to assess the percent of surgery patients who received appropriate venous thromboembolism (VTE) prophylaxis within 24 hours prior to Surgical Incision Time to 24 hours after Surgery End Time. 		
Outpatient acute myocardial infarction (AMI) and chest pain measures The hospital outpatient process of care measures include the following measures related to acute myocardial infarctions (AMI) and chest pain emergency department care: Median time to fibrinolysis (OP-1) – This measure is used to assess the percent of patients with extended median time	Hospitals must submit data for each of the outpatient acute myocardial infarction (AMI) and chest pain quality measures. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures	Specifications manual for hospital outpatient department quality measures, version 4.0, encounter dates 01-01-11 (1Q11) through 06-30-11 (2Q11). Centers for Medicare & Medicaid Services (CMS); May 2010 or as updated.

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
from emergency department (ED) arrival to administration of fibrinolytic therapy in ED acute myocardial infarction (AMI) patients with ST-segment elevation or left bundle branch block (LBBB) on the electrocardiogram (ECG) performed closest to ED arrival and prior to transfer.	 Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate 	Measure specifications can be found on the Centers for Medicare & Medicaid Services (CMS), QualityNet website http://qualitynet.org
■ Fibrinolytic therapy received within 30 minutes of emergency department (ED) arrival (OP-2) – This measure is used to assess the percent of emergency department (ED) acute myocardial infarction (AMI) patients receiving fibrinolytic therapy during the ED stay and having a time from ED arrival to fibrinolysis of 30 minutes or less.		
• Median time to transfer to another facility for acute coronary intervention (OP-3) – This measure is used to assess the percent of patients with extended median time from emergency department (ED) arrival to time of transfer to another facility for acute coronary intervention for ED acute myocardial infarction (AMI) patients.		
Aspirin at arrival (OP-4) – This measure is used to assess the percent of emergency department (ED) acute myocardial infarction (AMI) patients or chest pain patients (with Probable Cardiac Chest Pain) without aspirin contraindications who received aspirin within 24 hours before ED arrival or prior to transfer.		
• Median time to ECG (OP-5) – This measure is used to assess the percent of patients with extended median time from emergency department (ED) arrival to electrocardiogram (ECG) (performed in the ED prior to transfer) for acute myocardial infarction (AMI) or chest pain patients (with Probable Cardiac Chest Pain).		
Outpatient surgery department measures The hospital outpatient process of care measures include the	Hospitals must submit data for each of the outpatient surgery department quality measures. This data	Specifications manual for hospital outpatient department quality measures, version 4.0,

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
 Timing of antibiotic prophylaxis (prophylactic antibiotic initiated within one hour prior to surgical incision*) (OP-6) – This measure is used to assess the percent of surgical patients with prophylactic antibiotics initiated within one hour * prior to surgical incision. *Patients who received vancomycin or a fluoroquinolone for prophylaxis should have the antibiotic initiated within two hours prior to surgical incision. Due to the longer infusion time required for vancomycin or a fluoroquinolone, it is acceptable to start these antibiotics within two hours prior to incision time. Prophylactic antibiotic selection for surgical patients (OP-7) – This measure is used to assess the percent of surgical patients who received prophylactic antibiotics consistent with current guidelines (specific to each type of surgical procedure). 	 Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate 	encounter dates 01-01-11 (1Q11) through 06-30-11 (2Q11). Centers for Medicare & Medicaid Services (CMS); May 2010 or as updated. Measure specifications can be found on the Centers for Medicare & Medicaid Services (CMS), QualityNet website http://qualitynet.org
Home Management Plan of Care Given to Patient/Caregiver for Pediatric Asthma (CAC-3) – This measure is used to assess the number of pediatric asthma inpatients with documentation that they or their caregivers were given a written Home Management Plan of Care (HMPC) document. The HMPC document addresses all of the following: Arrangements for follow-up care Environmental control and control of other triggers Method and timing of rescue actions Use of controllers Use of relievers 	Hospitals must submit data for the home management plan of care given to patient/caregiver for pediatric asthma (CAC-3) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure. Numerator: Number of patients meeting the targets in the quality measure Calculated rate	Specifications manual for national hospital inpatient quality measures, version 3.2b, discharges 10-01-10 (4Q10) through 03-31-11 (1Q11). Centers for Medicare & Medicaid Services (CMS), The Joint Commission; April 2010 or as updated. Measure specifications can be found on the Centers for Medicare & Medicaid Services (CMS), QualityNet website http://qualitynet.org

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
Appropriate Care Measures (ACM)		
Acute myocardial infarction appropriate care measure (AMI-ACM) The ACM is a pass/fail measure at the individual patient level that asks whether eligible patients have received ALL of the appropriate care for acute myocardial infarction (AMI). The following individual measures are included in the AMI-ACM: AMI-1: Aspirin at arrival AMI-2: Aspirin prescribed at discharge AMI-3 ACEI or ARB for LVSD AMI-4: Adult smoking cessation advice/counseling AMI-5: Beta blocker prescribed at discharge AMI-7a: Thrombolytic within 30 minutes of hospital arrival AMI-8a: PCI within 90 minutes of hospital arrival	Hospitals must submit data for each of the hospital compare acute myocardial infaction (AMI)/heart attack process measures. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate	Specifications for acute myocardial infaction appropriate care: Stratis Health; 2009. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/healthreform
Heart failure appropriate care measure (HF-ACM) The ACM is a pass/fail measure at the individual patient level that asks whether eligible patients have received ALL of the appropriate care for heart failure (HF). The following individual measures are included in the HF-ACM: HF-1: Discharge instructions HF-2: LVF assessment HF-3: ACEI or ARB for LVSD HF-4: Adult smoking cessation advice/counseling	Hospitals must submit data for each of the hospital compare heart failure (HF) process measures. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate	Specifications for heart failure appropriate care: Stratis Health; 2009. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/healthreform

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
Pneumonia appropriate care measure (PN-ACM) The ACM is a pass/fail measure at the individual patient level that asks whether eligible patients have received ALL of the appropriate care for pneumonia (PN). The following individual measures are included in the PN-ACM: PN-2: Pneumococcal vaccination PN-3a: Blood cultures within 24 hours before or after ICU admission PN-3b: Blood cultures before antibiotic PN-4: Adult smoking cessation advice/counseling PN-5c: Initial antibiotic within 6 hours of hospital arrival PN-6: Initial antibiotic selection for CAP in immunocompetent patient	Hospitals must submit data for each of the hospital compare pneumonia (PN) process measures. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate	Specifications for pneumonia appropriate care: Stratis Health; 2009. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/healthreform

Measure Name and Purpose	Data Elements	Specification Information	
Measures Required for Reporting Beginning in January 2011 and	Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
Agency for Healthcare Research and Quality (AHRQ) Inpatient Quality Indicators (IQI)			
Abdominal aortic aneurysm (AAA) repair volume (IQI 4) – This measure is used to assess the raw volume of provider-level abdominal aortic aneurysm (AAA) repair (surgical procedure).	Hospitals must submit data for the abdominal aortic aneurysm (AAA) repair volume (IQI 4) quality measure. This data includes the following information: • Volume	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for	

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
		Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Abdominal aortic aneurysm (AAA) repair mortality rate (IQI 11) – This measure is used to assess the number of deaths per 100 discharges with procedure code of abdominal aortic aneurysm (AAA) repair.	Hospitals must submit data for the abdominal aortic aneurysm (AAA) repair mortality rate (IQI 11) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Coronary artery bypass graft (CABG) volume (IQI 5) – This measure is used to assess the raw volume of provider-level coronary artery bypass graft (CABG) (surgical procedure).	Hospitals must submit data for the coronary artery bypass graft (CABG) volume (IQI 5) quality measure. This data includes the following information: • Volume	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for
		Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov

Measure Name and Purpose	Data Elements	Specification Information	
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter			
Coronary artery bypass graft (CABG) mortality rate (IQI 12) – This measure is used to assess the number of deaths per 100 discharges with a procedure code of coronary artery bypass graft (CABG).	Hospitals must submit data for the coronary artery bypass graft (CABG) mortality rate (IQI 12) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov	
Percutaneous transluminal coronary angioplasty (PTCA) volume (IQI 6) – This measure is used to assess the raw volume of provider-level percutaneous transluminal coronary angioplasty (PTCA) (surgical procedure).	Hospitals must submit data for the percutaneous transluminal coronary angioplasty (PTCA) volume (IQI 6) quality measure. This data includes the following information: • Volume	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov	
Percutaneous transluminal coronary angioplasty (PTCA) mortality rate (IQI 30) – This measure is used to assess the number of deaths per 100 percutaneous transluminal coronary angioplasties (PTCAs).	Hospitals must submit data for the percutaneous transluminal coronary angioplasty (PTCA) mortality rate (IQI 30) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated.	

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
	 measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate 	Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Hip fracture mortality rate (IQI 19) – This measure is used to assess the number of deaths per 100 discharges with principal diagnosis code of hip fracture.	Hospitals must submit data for the hip fracture mortality rate (IQI 19) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Mortality for selected conditions composite measure This measure is used to assess the number of deaths for acute myocardial infarction (AMI), congestive heart failure (CHF), acute stroke, GI hemorrhage, hip fracture, and pneumonia. This composite measure includes the Agency for Healthcare Research and Quality (AHRQ) Inpatient Quality Indicators (IQI) related to hospital inpatient mortality for specific conditions: Acute myocardial infarction (AMI) mortality rate (IQI 15) Congestive heart failure (CHF) mortality rate (IQI 16) Acute stroke mortality rate (IQI 17)	Hospitals must submit data for the mortality for selected conditions composite measure and for each of the mortality for selected conditions composite measure component indicators. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate	See specific mortality for selected conditions composite measure component indicators for more information. Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
 GI Hemorrhage mortality rate (IQI 18) Hip fracture mortality rate (IQI 19) Pneumonia mortality rate (IQI 20) 		Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Agency for Healthcare Research and Quality (AHRQ) Patient Saf	<u> </u>	Driver Control (DOD)
Pressure ulcer (PSI 3) – This measure is used to assess the number of cases of decubitus ulcer per 1,000 discharges with a length of stay greater than 4 days. (Behavorial health only hospitals are not required to submit data on this measure.)	Hospitals must submit data for the pressure ulcer (PSI 3) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate	Patient Safety Indicators (PSI) Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Death among surgical inpatients with serious treatable complications (PSI 4) – This measure is used to assess the number of deaths per 1,000 patients having developed specified complications of care during hospitalization.	Hospitals must submit data for the death among surgical inpatients with serious treatable complications (PSI 4) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in each of the quality measure Calculated rate	Patient Safety Indicators (PSI) Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
		hrq.gov
Postoperative pulmonary embolism (PE) or deep vein thrombosis (DVT) (PSI 12) – This measure is used to assess the number of cases of deep vein thrombosis (DVT) or pulmonary embolism (PE) per 1,000 surgical discharges with an operating room procedure.	Hospitals must submit data for the postoperative pulmonary embolism (PE) or deep vein thrombosis (DVT) (PSI 12) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate	Patient Safety Indicators (PSI) Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Obstetric trauma – vaginal delivery with instrument (PSI 18) – This measure is used to assess the number of cases of obstetric trauma (3 rd or 4 th degree lacerations) per 1,000 instrument-assisted vaginal deliveries.	Hospitals must submit data for the obstetric trauma – vaginal delivery with instrument (PSI 18) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate	Patient Safety Indicators (PSI) Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Obstetric trauma – vaginal delivery without instrument (PSI 19) – This measure is used to assess the number of cases of obstetric trauma (3 rd or 4 th degree lacerations) per 1,000 without instrument assistance.	Hospitals must submit data for the obstetric trauma – vaginal delivery without instrument (PSI 19) quality measure. This data includes the following information:	Patient Safety Indicators (PSI) Technical Specifications, version 4.1. Agency for Healthcare Research and

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
	 Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate 	Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Patient safety for selected indicators composite measure This measure is used to assess the number of potentially preventable adverse events for pressure ulcer, iatrogenic pneumothorax, selected infections due to medical care, postoperative hip fracture, postoperative deep vein thrombosis (DVT) or pulmonary embolism (PE), postoperative sepsis, postoperative wound dehiscence, and accidental puncture or laceration (separately). This composite measure includes all of the Agency for Healthcare Research and Quality (AHRQ) Patient Safety Indicators related to hospital inpatient mortality for specific conditions: Pressure ulcer (PSI 3) Iatrogenic pneumothorax (PSI 6) Selected infections due to medical care (PSI 7) Postoperative hip fracture (PSI 8) Postoperative pulmonary embolism (PE) or deep vein thrombosis (DVT) (PSI 12) Postoperative sepsis (PSI 13) Postoperative wound dehiscence (PSI 14) Accidental puncture or laceration (PSI 15)	Hospitals must submit data for the patient safety for selected indicators composite measure and for each of the patient safety for selected indicators composite measure component indicators. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate	See specific patient safety for selected indicators composite measure component indicators for more information. Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and Every Year Thereafter		
Agency for Healthcare Research and Quality (AHRQ) Pediatric P	ratient Safety Indicators (PDI)	
Pediatric heart surgery volume measure (PDI 7) – This measure is used to assess the number of inhospital congenital heart surgeries for pediatric patients.	Hospitals must submit data for the pediatric patients undergoing surgery for congenital heart disease volume (PDI 7) quality measure. This data includes the following information: • Volume	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be
		found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Pediatric heart surgery mortality rate measure (PDI 6) – This measure is used to assess the number of inhospital deaths in pediatric patients undergoing surgery for congenital heart disease.	Hospitals must submit data for the pediatric patients undergoing surgery for congenital heart disease repair mortality rate (PDI 6) quality measure. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in the quality measure Numerator: Number of patients meeting the targets in the quality measure Calculated rate	Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov
Pediatric patient safety for selected indicators composite measure This measure is used to assess the number of potentially preventable adverse events for accidental puncture or laceration, pressure ulcer,	Hospitals must submit data for the pediatric patient safety for selected indicators composite measure and for each of the pediatric patient safety for selected	See specific pediatric patient safety for selected indicators composite measure component

Measure Name and Purpose	Data Elements	Specification Information
Measures Required for Reporting Beginning in January 2011 and	Every Year Thereafter	
iatrogenic pneumothorax, postoperative wound dehiscence, postoperative sepsis, and selected infections due to medical care (separately). This composite measure includes all of the Agency for Healthcare Research and Quality (AHRQ) Patient Safety Indicators related to hospital inpatient mortality for specific conditions: - Accidental puncture or laceration (PDI 1) - Pressure ulcer (PDI 2) - Iatrogenic pneumothorax (PDI 5) - Postoperative sepsis (PDI 10) - Postoperative wound dehiscence (PDI 11) - Selected infections due to medical care (PDI 12)	 indicators composite measure component indicators. This data includes the following information: Denominator: Number of patients meeting the criteria for inclusion in each of the quality measures Numerator: Number of patients meeting the targets in each of the quality measures Calculated rate 	indicators for more information. Inpatient Quality Indicators Technical Specifications, version 4.1. Agency for Healthcare Research and Quality (AHRQ); December 2009 or as updated. Measure specifications can be found on the Agency for Healthcare Research and Quality (AHRQ), Quality Indicators website http://www.qualityindicators.a hrq.gov

Measure Name and Description	Data Elements	Specification Information
Measure Required for Reporting in Calendar Year 2011 and Every Year Thereafter (2010 Dates of Service)		
Vermont Oxford Network (VON)		
Late sepsis or meningitis in very low birth weight neonates This measure is used to assess the infection rate for inborn and outborn infants meeting certain age and weight requirements.	Hospitals with a level 3 neontal intensive care unit (NICU) must submit data for the late sepsis or meningitis in very low birth weight neonates. This data includes the following information:	Specifications for late sepsis or meningitis in very low birth weight neonates: Vermont Oxford Network.
	 Denominator: Number of patients meeting the criteria for inclusion in the quality 	Measure specifications can be found on the Vermont Oxford

Measure Name and Description	Data Elements	Specification Information
Measure Required for Reporting in Calendar Year 2011 and Ever	ry Year Thereafter (2010 Dates of Service)	
	 measure. Numerator: Number of patients meeting the targets in the quality measure Calculated rate 	Network website http://www.vtoxford.org
Measures Required for Reporting Beginning in August 2011 and l	Every Year Thereafter (2011 Dates of Service)	
National Healthcare Safety Network (NHSN)		
Central line-associated bloodstream infection (CLABSI) event This measure is used to assess the infection rate of patients with a central line-associated bloodstream infection (CLABSI) event by inpatient hospital unit.	Hospitals must submit data for the central line- associated bloodstream infection (CLABSI) event by neonatal and pediatric intensive care units. This data includes the following information for each intensive care unit:	Specifications for the central line-associated bloodstream infection (CLABSI) event: Center for Disease Control and Prevention; June 2010.
	 Denominator: Number of patients meeting the criteria for inclusion in the quality measure. Numerator: Number of patients meeting the targets in the quality measure Calculated rate 	Measure specifications can be found on the Centers for Disease Control and Prevention website http://www.cdc.gov/nhsn/psc_da.html

Measure Name and Description	Data Elements	Specification Information	
Measure Required for Reporting in Calendar Year 2011 and Every Year Thereafter			
Health Information Technology (HIT)			
Health information technology (HIT) This survey is used to assess a hospital's adoption and use of Health Information Technology (HIT) in its clinical practice.	The information technology supplement of the American Hospital Association (AHA) annual survey and any additional Minnesota specific questions.		
Measures Required for Reporting Beginning in January 2011 (20)	10 Discharge Dates) and Every Year Thereafter		
Patient Experience of Care	Patient Experience of Care		
Patient experience of care This measure is used to assess adult patients' perception of their hospital care using a national survey called the Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS). (This measure is not required for hospitals with less than 500 admissions in the previous calendar year.)	Consumer assessment of healthcare providers and systems hospital (HCAHPS) survey	Consumer assessment of healthcare providers and systems hospital survey (HCAHPS), Version 5.0. Centers for Medicare & Medicaid Services (CMS); March 2010 or as updated. Measure specifications can be found on the HCAHPS website http://www.hcahpsonline.org/	

Measure Name and Description	Data Elements	Specification Information	
Retired Measures			
Hospital Compare Quality Measures, Center for Medicare & Medicaid Services (CMS) and The Joint Commission			
Surgical care improvement project (SCIP) Colorectal surgery patients with immediate postoperative normothermia (SCIP-Inf-7) – This measure is used to assess the percent of colorectal surgery patients with immediate normothermia (greater than or equal to 96.8° F) within the first fifteen minutes after leaving the operating room.	Hospitals are no longer required to submit data for this measure.	Dates of service ending 06/30/10 (2Q10).	

APPENDIX C REQUIRED AMBULATORY SURGICAL CENTER MEASURES

Data Required for Reporting Beginning in Calendar Year 2011 and Every Year Thereafter			
Measure Name and Description	Data Elements	Specification Information	
Measures Required for Reporting Beginning in July 2011 (Dates of Service July 1, 2010- June 30, 2011) and Every Year Thereafter			
Prophylactic intravenous (IV) antibiotic timing – This measure is used to assess the percent of ambulatory surgery center (ASC) patients who were administered antibiotics for prevention of surgical site infection on time	Ambulatory surgical centers must submit data for the prophylactic intravenous (IV) antibiotic timing quality measure. This data includes the following information: Patient identification methodology Submit the following three data elements: Denominator: Number of patients meeting the criteria for inclusion in the measure Denominator: Number of patients in data submission if submitting a sample Numerator: Number of patients meeting the targets in the measure Number of patients meeting the exclusion criteria Calculated rate Distribution of the ambulatory surgical center's total patient population by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay).	ASC quality measures: implementation guide – Version 1.3. ASC Quality Collaboration; April 2008. This measure was developed by the Ambulatory Surgery Centers (ASC) Quality Collaboration. Measure specifications can be found on the ASC Quality Collaboration website http://www.ascquality.org/doc uments/ASCQualityCollaborat ionImplementationGuide.pdf	
Hospital transfer/admission – This measure is used to assess the percent of ambulatory surgery center (ASC) patients who are transferred or admitted to a hospital upon discharge from the ASC.	Ambulatory surgical centers must submit data for the hospital transfer/admission quality measure. This data includes the following information: Patient identification methodology	ASC quality measures: implementation guide – Version 1.3. ASC Quality Collaboration; April 2008. This measure was developed	

Measure Name and Description	Data Elements	Specification Information	
Measures Required for Reporting Beginning in July 2011 (Dates	Measures Required for Reporting Beginning in July 2011 (Dates of Service July 1, 2010- June 30, 2011) and Every Year Thereafter		
	 Submit the following three data elements: Denominator: Number of patients meeting the criteria for inclusion in the measure Denominator: Number of patients in data submission if submitting a sample Numerator: Number of patients meeting the targets in the measure Number of patients meeting the exclusion criteria Calculated rate Distribution of the ambulatory surgical center's total patient population by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay). 	by the Ambulatory Surgery Centers (ASC) Quality Collaboration. Measure specifications can be found on the ASC Quality Collaboration website http://www.ascquality.org/doc uments/ASCQualityCollaborat ionImplementationGuide.pdf	
Appropriate surgical site hair removal – This measure is used to assess the percent of ambulatory surgery center (ASC) patients who have appropriate surgical site hair removal.	Ambulatory surgical centers must submit data for the appropriate surgical site hair removal quality measure. This data includes the following information: Patient identification methodology Submit the following three data elements: Denominator: Number of patients meeting the criteria for inclusion in the measure Denominator: Number of patients in data submission if submitting a sample Numerator: Number of patients meeting the targets in the measure Number of patients meeting the exclusion criteria	ASC quality measures: implementation guide — Version 1.3. ASC Quality Collaboration; April 2008. This measure was developed by the Ambulatory Surgery Centers (ASC) Quality Collaboration. Measure specifications can be found on the ASC Quality Collaboration website http://www.ascquality.org/doc uments/ASCQualityCollaborat ionImplementationGuide.pdf	

Measure Name and Description	Data Elements	Specification Information
Measures Required for Reporting Beginning in July 2011 (Dates of Service July 1, 2010- June 30, 2011) and Every Year Thereafter		
	 Calculated rate Distribution of the ambulatory surgical center's total patient population by primary payer type (i.e., private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay). 	

APPENDIX D OTHER STANDARDIZED QUALITY MEASURES

Measure Name	Measure Elements	Specification Information
Unlimited Availability		
Healthcare Effectiveness Data and Information Set (HEDIS)	All Healthcare Effectiveness Data and Information Set (HEDIS) measures as of HEDIS 2010, or as updated, that are applicable to physician clinics are included in the standardized set of quality measures.	Healthcare Effectiveness Data and Information Set (HEDIS) 2010 Volume 2: Technical Specifications. National Committee for Quality Assurance (NCQA); 2009 or as updated.
National Quality Forum (NQF) endorsed measures	All NQF-endorsed measures as of August 1, 2010, or as updated, that are applicable to physician clinics and hospitals are included in the standardized set of quality measures, excluding those requiring use of proprietary databases or registries.	More information about these measures can be found on the National Quality Forum (NQF), website http://qualityforum.org

Measure Name	Measure Elements	Specification Information	
Time-Limited Availability: These Measures are Available for Use for Three Years (2010 Dates of Service Through 2012 Dates of Service)			
Pediatric asthma	 For patients, ages 5 to 19 years: At the last asthma visit, is the asthma severity level (intermittent or persistent) documented? If it is documented that the patient has persistent asthma, is the patient on an anti-inflammatory medication? If the patient is a smoker or exposed to second hand smoke, is advice regarding smoking cessation documented in the last year? 	Clinical Review: Measure Specifications for Pediatric Asthma. Medica; Pediatric Asthma; July 2009. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/ healthreform	

• If the patient is a smoker, or exposed to second hand smoke, is advice regarding smoking cessation documented at the last visit?
Is the child's environment tobacco smoke free?
Is there evidence in the medical records that the patient received a written Asthma Action Plan?
• Unless contraindicated, is there evidence in the medical record that the patient has received an influenza vaccine within the past year?

Measure Name	Measure Elements	Specification Information
Time-Limited Availability: These Measures are Available for Use for Two Years (2010 Dates of Service Through 2011 Dates of Service)		
Adult depression	 At the time of initial diagnosis in the past year, was there documentation of at least 5 of the symptoms of depression, or that the patient met the criteria for depression using a standardized depression diagnosing tool? At the time of the initial diagnosis in the past year, is there documentation that the patient was assessed for alcohol consumption? If the patient was at the clinic for a primary care visit more than 6 weeks after the initial diagnosis, is there documentation of ongoing assessment of their depression status? Was the patient's depression status reassessed using the PHQ-9 tool at a follow up visit within 6 months after diagnosis? Was the PHQ-9 depression diagnostic tool used for the initial depression diagnosis? 	Clinical Review: Measure Specifications for Adult Depression. Medica; Pediatric Asthma; July 2009. Measure specifications can be found on the Minnesota Department of Health website http://www.health.state.mn.us/ healthreform

APPENDIX E SUBMISSION SPECIFICATIONS

I. Submission Requirements for Physician Clinics

- 1. **Registration.** Each physician clinic must register electronically and obtain a login user ID and password from the commissioner or commissioner's designee beginning January 1, 2010 and no later than February 10, 2010 and no later than February 10 of each subsequent year, and must supply data elements, including the following:
 - **a. Physician clinic information:** Name, mailing address, clinic national provider identifier (NPI);
 - **b.** Contact information for individual(s) responsible for submitting data: Company, name, title, mailing address, telephone number, fax number, e-mail address;
 - **c.** Contact information for physician clinic general contact: Name, title, mailing address, telephone number, fax number, e-mail address;
 - **d.** Clinical staff information for the previous calendar year: Name, national provider identifier (NPI), board certifications for all clinical staff that have provided health care services at the physician clinic during the previous calendar year;
 - e. Description of health care services provided by the physician clinic.
 - f. Medical group affiliation.

2. Data Submission.

a. Measures for which physician clinics may submit on their full patient population or a random sample in 2011. (NOTE: Beginning in 2012, physician clinics with electronic medical records in place for the prior full measurement period are required to submit data on their full patient population.)

Optimal diabetes care (ODC) and optimal vascular care (OVC). Each physician clinic, except ambulatory surgical centers, must submit the data required to calculate the applicable quality measures, as described in Appendix A and including the number of patients receiving the applicable health care services allocated according to primary payer type (private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay) to the commissioner or the commissioner's designee. Specifically, this includes patient identification methodology, numerator and denominator by primary payer type, number of patients meeting the exclusion criteria, and calculated rate. If submitting a sample, the denominator for the entire patient population does not need to be allocated by primary payer type. A physician clinic may work with a single subcontractor to submit the required data on their behalf. Data may be submitted beginning January 1, 2011 and no later than February 15, 2011 and beginning January 1 and no later than February 15 of each subsequent year.

Optimal asthma care. Each physician clinic, except ambulatory surgical centers, must submit the data required to calculate the applicable quality measures, as described in Appendix A to the commissioner or the commissioner's designee. For this measure, this includes identifying the patients in two separate age bands, ages 5-17 and ages 18-

50. If the physician clinic chooses to submit a sample, there must be one sample per age band. Within these two age bands, data on the number of patients receiving the applicable health care services must be allocated according to primary payer type (private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay). Specifically, this includes patient identification methodology, separation of the data by age bands, numerator and denominator by primary payer type, number of patients meeting the exclusion criteria, and calculated rate. If submitting a sample, the denominator for the entire patient population does not need to be allocated by primary payer type. A physician clinic may work with a single subcontractor to submit the required data on their behalf. Data may be submitted beginning July 1, 2011 and no later than August 15, 2011 and beginning July 1 and no later than August 15 of each subsequent year.

Colorectal cancer screening. Each physician clinic, except ambulatory surgical centers, must submit the data required to calculate the applicable quality measures, as described in Appendix A and including the number of patients receiving the applicable health care services allocated according to primary payer type (private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay) to the commissioner or the commissioner's designee. Specifically, this includes patient identification methodology, numerator and denominator by primary payer type, number of patients meeting the exclusion criteria, and calculated rate. If submitting a sample, the denominator for the entire patient population does not need to be allocated by primary payer type. A physician clinic may work with a single subcontractor to submit the required data on their behalf. Data may be submitted beginning July 1, 2011 and no later than August 15, 2011 and beginning July 1 and no later than August 15 of each subsequent year.

- i. **Data Submission Requirements.** A physician clinic may satisfy the data submission requirement for these quality measures by completing the following steps:
 - 1. **Patient identification methodology.** Identify patients meeting the criteria for inclusion in the measure. Use the measurement specifications referenced in Appendix A to determine eligibility for each patient, only including patients that meet denominator criteria for each measure in the list. Develop a list of the eligible patients for each measure using a practice management, billing system, or electronic medical record.
 - 2. Data Collection: Total Population versus Sample. Identification of the population of patients eligible for the denominator for each measure is accomplished via a query of a practice management system or an electronic medical record. Use the measurement specifications referenced in Appendix A to determine eligibility for each patient, only including patients that meet denominator criteria for each measure in the list. Physician clinics may choose one of the following options:
 - a. Full patient population. Physician clinics are encouraged to submit data using their full patient population for each

- measure, but may use a random sampling methodology, as described below.
- b. Random sampling methodology. Physician clinics may submit data on a random sample of relevant patients in 2011. At a minimum, physician clinics must select 60 patients for the random sample population and must oversample by at least 20 patients. If a physician clinic's total population for a particular measure is less than 60, the physician clinic must submit data using their full patient population for that measure. In subsequent years, physician clinics with an electronic medical record in place for a full measurement period will be expected to submit data on a full population basis. (NOTE: For the optimal asthma measure, there must be one sample per age band, one for ages 5-17 and one for ages 18-50.)
- 3. **Data submission template**. Use the data submission template supplied annually by the commissioner or the commissioner's designee as a data collection tool. Data elements may be either extracted from an electronic medical record system or abstracted through medical record review.
- 4. **Data file upload.** Submit data electronically to the commissioner or the commissioner's designee.
- 5. **Data validation.** Physician clinics must maintain documentation for the data described in Appendix A, including the methodology used to determine patients meeting the criteria for inclusion in each measure and the data submission template, for purposes of data validation.
- b. Measures for which physician clinics may only submit data on their full patient population in 2011.

Depression remission at six months. Each physician clinic, except ambulatory surgical centers, must submit the data required to calculate the applicable quality measures, as described in Appendix A to the commissioner or the commissioner's designee. For this measure the data elements must be submitted by three bands of initial PHQ9 scores (10-14; 15-19; 20 and above). The number of patients for whom a follow up six month (+/- 30 days) PHQ9 assessment was not completed is a required data element as well. Specifically the required data includes, patient identification methodology, numerator and denominator separated by three bands of initial PHQ9 scores, number of patients who did not get a second PHQ9 assessment, number of patients meeting the exclusion criteria, and calculated rate. A physician clinic may work with a single subcontractor to submit the required data on their behalf. Data may be submitted beginning February 7, 2011 and no later than February 25, 2011.

i. **Data Submission Requirements.** A physician clinic may satisfy the data submission requirement for these quality measures by completing the following steps:

- 1. **Patient identification methodology.** Identify patients meeting the criteria for inclusion in the measure. Use the measurement specifications referenced in Appendix A to determine eligibility for each patient, only including patients that meet denominator criteria for each measure in the list. Develop a list of the eligible patients for each measure using a practice management, billing system, or electronic medical record
- 2. **Data Collection: Total Population.** Identification of the population of patients eligible for the denominator for each measure is accomplished via a query of a practice management system or an electronic medical record. Use the measurement specifications referenced in Appendix A to determine eligibility for each patient, only including patients that meet denominator criteria for each measure in the list. For this measure physician clinics must submit data using their full patient population.
- 3. **Data submission template**. Use the data submission template supplied annually by the commissioner or the commissioner's designee as a data collection tool. Data elements may be either extracted from an electronic medical record system or abstracted through medical record review.
- 4. **Data file upload.** Submit data electronically to the commissioner or the commissioner's designee.
- 5. **Data validation.** Physician clinics must maintain documentation for the data described in Appendix A, including the methodology used to determine patients meeting the criteria for inclusion in each measure and the data submission template, for purposes of data validation.
- **3. Health information technology (HIT) survey.** Each physician clinic must complete the internet-based survey available annually from the commissioner or commissioner's designee beginning February 15, 2010 and no later than March 15, 2010 and beginning February 15 and no later than March 15 of each subsequent year.
- **4. Patient experience of care survey.** Each physician clinic must use a vendor certified by CMS. Each physician clinic must either select a CMS-certified vendor of its choice or use the services of a centralized vendor coordinated by the commissioner or the commissioner's designee. Survey period will include patients seen September 1, 2012 November 30, 2012.

Minnesota Statewide Quality Reporting and Measurement System

¹ CMS only certifies vendors to administer the CMS-required HCAHPS survey. CMS does not certify vendors to administer CG-CAHPS for physician clinics. For purposes of fulfilling state requirements under Chapter 4654, physician clinics must use a vendor certified by CMS to administer HCAHPS.

II. Submission Requirements for Hospitals

- 1. Data Submission for Hospital Compare Measures, Centers for Medicare & Medicaid Services (CMS) and The Joint Commission. Each hospital must submit the data described in Appendix B required to calculate the applicable quality measures. There are two ways hospitals may satisfy this requirement:
 - a. Submission to the Centers for Medicare & Medicaid Services (CMS). If a hospital normally submits data for all cases for these quality measures to the Centers for Medicare & Medicaid Services (CMS), using the Centers for Medicare & Medicaid Services' (CMS) existing schedule, specifications, and processes, and continues to do so, the hospital will have satisfied their data submission requirements for these quality measures provided that the hospital also signs an authorization form allowing the data to be published on the U.S. Department of Health & Human Services Hospital Compare website for *all* cases for each applicable quality measure; or
 - **b.** Submission directly to commissioner or commissioner's designee. If a hospital does not submit data for these quality measures to the Centers for Medicare & Medicaid Services (CMS), the hospital must submit data to the commissioner or the commissioner's designee according to the following schedule:

Inpatient Quality Measures

Discharge Dates	Data Submission Deadline
Third Quarter, 2010: July 1 – September 30	February 15, 2011
Fourth Quarter, 2010: October 1 – December 31	May 15, 2011
First Quarter, 2011: January 1 – March 31	August 15, 2011
Second Quarter, 2011: April 1 – June 30	November 15, 2011

Outpatient Quality Measures

Discharge Dates	Data Submission Deadline
Third Quarter, 2010: July 1 – September 30	February 1, 2011
Fourth Quarter, 2010: October 1 – December 31	May 1, 2011
First Quarter, 2011: January 1 – March 31	August 1, 2011
Second Quarter, 2011: April 1 – June 30	November 1, 2011

- i. Data collection and analysis.
 - 1. Hospitals must use the CMS Abstraction & Reporting Tool (CART), available from the Centers for Medicare & Medicaid Services (CMS), for the collection and analysis of the data required to calculate each measure.

- 2. Use the measurement specifications referenced in Appendix B to determine whether each patient is eligibile for inclusion in the measurement calculation.
- ii. Data validation. At their own expense, hospitals must have their data validated by a third-party vendor using protocols and standards consistent with those of the Centers for Medicare & Medicaid Services (CMS) to verify that the data is consistent and reproducible.
- iii. Data submission. Submit data electronically to the commissioner or the commissioner's designee on a form provided by the commissioner or the commissioner's designee.
- **2. Data Submission for Appropriate Care Measures (ACM).** Each hospital must submit the data described in Appendix B required to calculate the applicable quality measures according to the following schedule:

Discharge Dates	Data Submission Deadline
Third Quarter, 2010: July 1 – September 30	February 15, 2011
Fourth Quarter, 2010: October 1 – December 31	May 15, 2011
First Quarter, 2011: January 1 – March 31	August 15, 2011
Second Quarter, 2011: April 1 – June 30	November 15, 2011

There are two ways hospitals may satisfy this requirement.

- a. Each hospital may authorize a single organization to complete the following steps and submit the data on their behalf:
 - Signs an authorization form allowing Stratis Health to release summary level ACM data, calculated from data in the CMS national data repository, to MHA for publication on the MHA website Minnesota Hospital Quality Report.
- b. Each hospital may perform the following steps itself:
 - i. Data collection and analysis. Identify the patients meeting the criteria for inclusion in the measure. Use the measurement specifications referenced in Appendix B to determine eligibility for each patient, only including patients that meet denominator criteria. The ACM is a pass/fail measure at the individual patient level that asks whether eligible patients have received ALL of the appropriate care for the condition they are being treated for. A patient is included if the patient meets denominator criteria for at least one of the measures in a topic. These topics include heart failure, acute myocardial infarction, and pneumonia. Within each topic, a patient must meet numerator criteria for each measure in which the patient meets denominator criteria to be considered as having appropriate care (Pass).
 - ii. Data submission. Submit data electronically to the commissioner or the commissioner's designee.

3. Data Submission for Inpatient Quality Indicators (IQI), Patient Safety Indicators (PSI), and Pediatric Patient Safety Indicators (PDI), Agency for Healthcare Research and Quality. Each hospital must submit the data described in Appendix B required to calculate the applicable quality measures according to the following schedule:

Discharge Dates	Data Submission Deadline
Third Quarter, 2010: July 1 – September 30	January 28, 2011
Fourth Quarter, 2010: October 1 – December 31	April 29, 2011
First Quarter, 2011: January 1 – March 31	July 23, 2011
Second Quarter, 2011: April 1 – June 30	October 22, 2011

There are two ways hospitals may satisfy this requirement.

- a. Each hospital may authorize a single organization to complete the following steps and submit the data on their behalf:
 - i. Apply Version 4.2, or the most recent version of the Quality Indicator software, available from the Agency for Healthcare Research and Quality's (AHRQ), to the hospital's discharge data. A hospital must participate in verifying the results of the analysis as needed.
 - ii. Validate the data.
 - 1. In the event data validation procedures show that data is inaccurate, hospitals must correct the inaccurate information and resubmit corrected data. Resubmitted data must be verified for accuracy.
 - 2. The results of the analysis using the Quality Indicator software for each hospital must be verified for accuracy by each hospital prior to submission.
 - iii. Submit the data to the commissioner or the commissioner's designee.
- b. Each hospital may perform the following steps itself:
 - i. Apply Version 4.2, or the most recent version of the Quality Indicator software, available from the Agency for Healthcare Research and Quality's (AHRQ), to its discharge data.
 - ii. Validate the data submission through a third-party vendor.
 - iii. Submit data electronically to the commissioner or the commissioner's designee on a form provided by the commissioner or the commissioner's designee.
- **4. Data Submission for Vermont Oxford Network (VON).** Each hospital with a level 3 neonatal intensive care unit (NICU) must submit the data required to calculate the applicable quality measure, as described in Appendix B, to VON.
 - a. Each hospital with a level 3 NICU must submit applicable data on the specified patients to VON, for measure calculation and inclusion in VON's annual report to the hospital, according to the following VON data submission schedule:

Discharge Dates	Data Submission Deadline
All 2010 Dates of Service	June 30, 2011

- b. Each hospital with a level 3 NICU must submit summary level results electronically for the previous calendar year to the commissioner or the commissioner's designee by October 31, 2011 and every year thereafter.
- **5. Data Submission for the National Healthcare Safety Network (NHSN).** Each hospital with a neonatal and/or pediatric intensive care unit must submit the data required to calculate the applicable quality measure, as descibed in Appendix B, on the following schedule:

Discharge Dates	Data Submission Deadline
First Quarter, 2011: January 1 – March 31	August 15, 2011
Second Quarter, 2011: April 1 – June 30	November 15, 2011
Third Quarter, 2011: July 1 – September 30	January 15, 2012
Fourth Quarter, 2011: October 1 – December 31	April 15, 2012

- a. Each hospital with a neonatal and/or pediatric intensive care unit must electonically submit summary level data according to NHSN definitions, for each intensive care unit, to the commissioner or the commissioner's designee for measure calculation.
- **6. Health information technology (HIT) survey.** Each hospital must complete the survey available annually from the commissioner or commissioner's designee in calendar year 2010 and each subsequent year.
- **7. Patient expereince of care survey.** Each hospital must complete the HCAHPS survey using a CMS-cerified vendor.

III. Submission Requirements for Ambulatory Surgical Centers

- 1. **Registration.** Each ambulatory surgical center must register electronically and obtain a login user ID and password from the commissioner or commissioner's designee beginning March 1, 2011 and no later than April 1, 2011 and no later than April 1 of each subsequent year; and must supply data elements, including the following:
 - **a. Ambulatory Surgical Center information:** Name, mailing address, ambulatory surgical center national provider identifier (NPI);
 - **b.** Contact information for individual(s) responsible for submitting data: Company, name, title, mailing address, telephone number, fax number, e-mail address;
 - **c. Contact information for ambulatory surgical center general contact:** Name, title, mailing address, telephone number, fax number, e-mail address;
 - **d.** Clinical staff information for the previous calendar year: Name, national provider identifier (NPI), board certifications for all clinical staff that have provided health care services at the ambulatory surgical center during the previous calendar year;
 - f. Medical group affiliation if applicable.
- 2. Data Submission. Each ambulatory surgical center must submit the data required to calculate the applicable quality measures, as described in Appendix C, including the number of patients receiving the applicable health care services, as well as the distribution of the ambulatory surgical center's total patient population by primary payer type (private insurance, Medicare, Minnesota Health Care Programs, uninsured/self-pay) to the commissioner or the commissioner's designee. An ambulatory surgical center may work with a single subcontractor to submit the required data on their behalf. Data may be submitted beginning July 1, 2011 and no later than August 15, 2011 and beginning July 1 and no later than August 15 of each subsequent year. (NOTE: After the initial 2011 reporting year, each ASC must allocate the data required to calculate the applicable quality measures by primary payer type when the commissioner or the commissioner's designee determines the results must be risk adjusted. In 2012, based on current measures, this would apply to the hospital transfer/admission measure.)
 - a. Prophylactic intravenous (IV) antibiotic timing; Appropriate surgical site hair removal; and Hospital transfer/admission.
 - An ambulatory surgical center may satisfy the data submission requirements for these quality measures by completing steps formally outlined in the Summary Data Submission Guide.
 - i. Patient identification methodology. Use the measurement specifications referenced in Appendix C to determine eligibility for each patient, only including patients that meet denominator criteria for each measure in the list. Ambulatory surgical centers may submit data on a random sample of relevant patients in 2011. Ambulatory surgical centers with fewer than 60 relevant patients for each measure must submit data on all relevant patients. In subsequent years, ambulatory surgical centers with an electronic medical record in place for a full measurement period will be expected to submit data on a full population basis.

- **ii. Data submission template.** Use the data submission template supplied annually by the commissioner or the commissioner's designee as a data collection tool. Data elements may be either extracted from an electronic medical record system or abstracted through medical record review.
- **iii. Data file upload.** Submit data electronically to the commissioner or the commissioner's designee.
- iv. **Data validation.** Ambulatory surgical centers must maintain documentation for the data described in Appendix C including the methodology used to determine patients meeting the criteria for inclusion in each measure and the data submission template for purposes of data validation.

