This document is made available electronically by the Minnesota Legislative Reference Library as part of an ongoing digital archiving project. http://www.leg.state.mn.us/lrl/lrl.asp (Funding for document digitization was provided, in part, by a grant from the Minnesota Historical & Cultural Heritage Program.)

INVENTORY OF PEAT RESOURCES

KOOCHICHING COUNTY, MINNESOTA

APPENDIX E SITE DESCRIPTIONS

TEGISLATIVE REFERENCE LIBRARY STATE OF MINNESOTA

MINNESOTA DEPARTMENT OF NATURAL RESOURCES Neither the State of Minnesota nor the Department of Natural Resources, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Reference to a company or product name does not impring privately owned rights.

Reference to a company or product name does not imply approval or recommendation of the product by the State of Minnesota or the Department of Natural Resources to the exclusion of others that may meet specifications.

This report was prepared with the support of the U.S. Department of Energy, Grant

This report was prepared with the support of the U.S. Department of Energy, Grant No. DE-FG01-79ET14692. However, any opinions, findings, conclusions, or recommendations expressed herein are those of the authors and do not necessarily reflect the views of DOE.

INVENTORY OF PEAT RESOURCES KOOCHICHING COUNTY, MINNESOTA

APPENDIX E
SITE DESCRIPTIONS

MINNESOTA DEPARTMENT OF NATURAL RESOURCES DIVISION OF MINERALS

PEAT INVENTORY PROJECT

Funding Provided By U.S. Department of Energy, Grant No. DE-FG01-79ET14692

> Hibbing, Minnesota 1980

CEGISLATIVE REFERENCE LIBRARY
STATE OF MINNESOTA

This report is on deposit at
Minnesota Department of Natural Resources
Division of Minerals
345 Centennial Office Building
St. Paul, Minnesota 55155

INVENTORY OF PEAT RESOURCES KOOCHICHING COUNTY, MINNESOTA

APPENDIX E
SITE DESCRIPTIONS

MINNESOTA DEPARTMENT OF NATURAL RESOURCES DIVISION OF MINERALS

PEAT INVENTORY PROJECT

Funding Provided By U.S. Department of Energy, Grant No. DE-FG01-79ET14692

> Hibbing, Minnesota 1980

This report is on deposit at

Minnesota Department of Natural Resources
Division of Minerals

345 Centennial Office Building
St. Paul, Minnesota 55155

INVENTORY OF PEAT RESOURCES KOOCHICHING COUNTY, MINNESOTA

APPENDIX E SITE DESCRIPTIONS

This appendix contains supplementary data on the peat resources of Koochiching County, Minnesota. Included are site locations; peat profile descriptions, including underlying soil texture; vegetation descriptions; microrelief; and depth to water table.

Location: 800 feet S and 2280 feet E of the NW corner of Sec. 27, T.63N., R.22W. (NW1/4 of SE1/4 of NE1/4 of NW1/4, Sec. 27, T.63N., R.22W.)

Vegetation: Black spruce crown cover of about 90 percent; sparse understory consists of some Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry and snowberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer Depth (cm)
Hemic 0-168
Clay 168+

Location: 825 feet N and 1500 feet E of the SW corner of Sec. 35, T.63N., R.22W. (Gov. Lot 2, Sec. 35, T.63N., R.22W.)

Vegetation: Black spruce crown cover of about 65 percent; sparse understory consists of some willow. Labrador tea, sedges, and grasses; ground cover consists mostly of feather mosses with some Sphannum mosses.

Microrelief: 25 cm.

Depth To Water Table: 7 cm.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm)
Hemic	0-122
Fibric	122-229
Sapric	229-518
Silty	518+
clay	
loam	

Location: 2000 feet S and 200 feet E of the NW corner of Sec. 27, T.64N., R.22W. (Gov. Lot 5, Sec. 27, T.64N., R.22W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; understory consists of Labrador tea with some leatherleaf, sedges, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 25 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm
Hemic	0-213
Sapric	213-305
Sandy	305+
loam	

Location: 1900 feet S and 950 feet E of the NW corner of Sec. 15, T.66N., R.22W. (SW1/4 of NE1/4 of SW1/4 of NW1/4, Sec. 15, T.66N., R.22W.)

Vegetation: Black spruce crown cover of about 95 percent with scattered paper birch and aspen; lush understory consists mostly of speckled alder with Labrador tea, leatherleaf, bog rosemary, grasses, and some willow; ground cover consists of Sphagnum mosses with some feather mosses.

Microrelief: 40 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 28, 1979.

Layer	Depth (cm
Hemic	0-366
Sapric	366-442
Silty	442+
clay	

Location: 450 feet S and 1980 feet W of the NE corner of Sec. 18, T.66N., R.22W. (SE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 18, T.66N., R.22W.)

Vegetation: Scattered paper birch and black spruce; understory consists of Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: D. Mellem and P. Guertin on November 28, 1979.

Layer	Depth (cm)	
Hemic	0-213	
Sandy	213+	
clay		

Location: 460 feet N and 180 feet E of the SW corner of Sec. 4, T.67N., R.22W.) (NW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 4, T.67N., R.22W.)

Vegetation: Scattered paper birch and ash; lush understory consists mostly of grasses with some sedges and bog birch.

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 19, 1980.

.ayer	Depth (cm
lemic	0-274
Silty	274+
lay	
oam	

Location: 440 feet N and 2525 feet E of the SW corner of Sec. 4, T.67N., R.22W. (NE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 4, T.67N., R.22W.)

Vegetation: Scattered black spruce; understory consists of Labrador tea with some bog birch, willow, grasses, raspberry, and fern; ground cover consists of Sphagnum mosses with some other mosses.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 19, 1980.

Layer Depth (cm)

Layer	Depth (cm)
Hemic	0-411
Sapric	411-427
Silt	427+
loam	

Location: 1180 feet N and 500 feet W of the SE corner of Sec. 5, T.67N., R.22W. (NW1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 5, T.67N., R.22W.)

Vegetation: Scattered black spruce and nothern white cedar; understory consists of sedges with some grasses; ground cover consists mostly of mosses. Microrelief: 30 cm.

Depth To Water Table: 10 cm.

Described By: H. Mooers and D. Haverkost on May 19, 1980.

Layer	Depth (cm)	
Hemic	0-244	
Sapric	244-305	
Limnic	305-335	
Loamy sand	335+	

Location: 1000 feet N and 200 feet W of the SE corner of Sec. 5, T.67N., R.22W. (SE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 5, T.67N., R.22W.)

Vegetation: Scattered black spruce, northern white cedar, and paper birch; lush understory consists of grasses and bog birch with some sedges.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 19, 1980.

.ayer	Depth (cm
łemic	0-213
Sapric	213-244
/ery	244+
ine	
oamy	
and	

Vegetation: Black spruce crown cover of about 65 percent with scattered tamarack and aspen; sparse understory consists of some grasses; ground cover consists of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 14, 1980.

Layer Depth (cm)
Hemic 0-305
Clay 305+
loam

Location: 325 feet S and 1600 feet W of the NE corner of Sec. 9, T.67N., R.22W. (SE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 9, T.67N., R.22W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; understory consists of Labrador tea, grasses, and willow; ground cover consists mostly of Sphagnum mosses with some cranberry, snowberry, and other mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 14, 1980.

Layer	Depth (cm)
Hemic Silty	0-457 457+
clay	10. 1

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 15, 1980.

Layer	Depth (cm
Hemic	0-427
Silty clay	427+
loam	

Location: 320 feet S and 2110 feet W of the NE corner of Sec. 9. T.67N., R.22W.

Vegetation: Black spruce crown cover of about 40 percent with scattered ash and

tamarack; sparse understory consists of some Labrador tea, leatherleaf, and

grasses; ground cover consists of Sphagnum and other mosses with some

(NE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 9, T.67N., R.22W.)

Location: 310 feet S and 2610 feet W of the NE corner of Sec. 9, T.67N., R.22W. (NW1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 9, T.67N., R.22W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered ash and tamarack; understory consists of Labrador tea with some grasses and leatherleaf; ground cover consists mostly of mosses with some Sphagnum mosses and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 15, 1980.

Layer	Depth (cm	
Hemic	0-396	
Silt	396+	
loam		

Location: 350 feet S and 2050 feet E of the NW corner of Sec. 9, T.67N., R.22W. (SW1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 9, T.67N., R.22W.)

Vegetation: Black spruce crown cover of about 60 percent with an ash crown cover of about 35 percent with scattered tamarack; ground cover consists mostly of mosses with some Sphagnum mosses.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 15, 1980.

Layer	Depth (cm)
Hemic Silt	0-517 517+
loam	 '

Location: 375 feet S and 1640 feet E of the NW corner of Sec. 9, T.67N., R.22W. (NE1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 9, T.67N., R.22W.)

Vegetation: Black spruce crown cover of about 65 percent with scattered northern white cedar and paper birch; understory consists of Labrador tea and bog birch with some cotton grass; ground cover consists of Sphagnum and other mosses with some snowberry and cranberry.

Microrelief: 35 cm.

cranberry.

Microrellef: 35 cm.

Depth To Water Table: 4 cm.

Described By: H. Mooers and D. Haverkost on May 15, 1980.

ayer	Depth (cm)
emic	0-315
ilt	315+

Location: 480 feet S and 1080 feet E of the NW corner of Sec. 9, T.67N., R.22W. (SE1/4 of NE1/4 of NW1/4 of NW1/4, Sec. 9, T.67N., R.22W.)

Vegetation: Scattered black spruce, aspen, tamarack, and paper birch; understory consists of alder with some grasses and bog birch; ground cover consists mostly of Sphagnum mosses.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 15, 1980.

Layer	Depth (cm
Hemic Silty clay loam	0-183 183+

Location: 100 feet S and 100 feet E of the NW corner of Sec. 9, T.67N., R.22W. (NW1/4 of NW1/4 of NW1/4, Sec. 9, T.67N., R.22W.)

Vegetation: Consists mostly of sedges and grasses.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 19, 1980.

Layer	Depth (cm)
Hemic	0-122
Limnic	122-335
Silt	335+
ioam	

Location: 500 feet S and 150 feet E of the NW corner of Sec. 9, T.67N., R.22W. (SW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 9, T.67N., R.22W.)

Vegetation: Consists mostly of grasses.

Microrelief: 5 cm.

Depth To Water Table: 45 cm.

Described By: H. Mooers and D. Haverkost on May 15, 1980.

Layer	Depth (cm)
Hemic Silty	0-61 61+
clay loam	

Location: 500 feet S and 400 feet E of the NW corner of Sec. 9, T.67N., R.22W.

Vegetation: Scattered black spruce; lush understory consists mostly of grasses and boo birch.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 15, 1980.

Layer	Depth (cm)
Muck Silty clay	0-183 183+

Location: 1780 feet S and 2340 feet E of the NW corner of Sec. 9, T.67N., R.22W.

Vegetation: Black spruce and tamarack crown cover of about 80 percent; lush understory consists of speckled alder, paper birch, willow, aspen, black ash, Labrador tea, grasses, and sedges; ground cover consists of feather mosses with some Sohagnum mosses.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and R. Graber on November 28, 1979.

Layer	Depth (cm)
Hemic	0-229
Sapric	229-259
Silty	259+
clay	

Location: 2200 feet N and 1250 feet W of the SE corner of Sec. 7, T.68N., R.22W.

Vegetation: Scattered black spruce and aspen; lush understory consists mostly of Labrador tea with some willow, leatherleaf, grasses, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 15 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 27, 1979.

(SW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 7, T.68N., R.22W.)

Layer	Depth (cm)
Fibric	0-152
Hemic	152-427
Sapric	427-457
Clay	457+

Location: 1150 feet S and 1880 feet W of the NE corner of Sec. 5, T.68N., R.22W.

Vegetation: Scattered black spruce and tamarack; sparse understory consists of

leatherleaf, swamp laurel, sedges, cotton grass, and grasses; ground cover

(SW1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 5, T.68N., R.22W.)

Described By: R. Graber and D. Haverkost on November 27, 1979.

consists mostly of Sphagnum mosses.

Depth To Water Table: At surface.

Microrellef: 15 cm.

Layer	Depth (cm)
Hemic Silty clay	0-46 46+

Location: 1780 feet N and 90 feet E of the SW corner of Sec. 11, T.68N., R.22W. (NW1/4 of SW1/4 of SW1/4, Sec. 11, T.68N., R.22W.)
Vecetation: Scattered black spruce: sparse understory consists of leatherleaf and

Location: 1250 feet N and 650 feet W of the SE corner of Sec. 11, T.67N., R.22W.

Vegetation: Scattered paper birch: understory consists of some speckled alder

Depth (cm)

(SW1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 11, T.67N, R.22W.)

Described By: H. Mooers and D. Haverkost on May 14, 1980.

Laver

Muck

Vegetation: Scattered black spruce; sparse understory consists of leatherleaf and grasses with some swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 15 cm.

and doowood

Depth To Water Table: Not visible.

Microrelief: 5 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer	Depth (cm
Fibric	0- 91
Hemic	91-427
Sapric	427-457+
Bottom	
unknown	

Location: 2420 feet N and 15 feet E of the SW corner of Sec. 4, T.69N., R.22W. (NW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 4, T.69N., R.22W.)

Vegetation: Black spruce crown cover of about 65 percent; sparse understory consists of some Labrador tea, swamp laurel, bog rosemary, and grasses; ground cover consists mostly of Sphagnum mosses with some snowberry and pitcher plant.

Microrelief: 15 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and B. Sether on March 20, 1980.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-213
Fibric	213-244
Hemic	244-305
Fibric	305-335
Hemic	335-487
Silt	487+
loam	

Location: 1930 feet N and 1760 feet E of the SW corner of Sec. 5, T.69N., R.22W. (NE1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 5, T.69N., R.22W.)

Vegetation: Tamarack crown cover of about 90 percent with scattered black spruce; understory consists of bog rosemary, grasses, speckled alder, Labrador tea, and leatherleaf; ground cover consists of Sphagnum mosses.
Microrellef: Negligible.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Hemic Bottom	0-548+
unknown	

Location: 200 feet S and 1240 feet W of the NE corner of Sec. 7, T.69N., R.22W. (NE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 7, T.69N., R.22W.)

Vegetation: Black spruce crown cover of about 30 percent with scattered black ash and tamarack; understory consists of Labrador tea and grasses with some speckled alder, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer	Depth (cm)
Fibric	0-183
Hemic	183-442+
Bottom	
unknown	

Location: 20 feet S and 2350 feet W of the NE corner of Sec. 7, T.69N., R.22W. (NW1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 7, T.69N., R.22W.)

Vegetation: Black spruce crown cover of about 65 percent; sparse understory consists of some leatherleaf, swamp laurel, grasses, and speckled aider; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrelief: 30 cm.

Depth To Water Table: 30 cm.

Described By: H. Mooers, B. Sether, and S. Nelson on March 20, 1980.

Layer	Depth (cm)	
Hemic	0-244	
Fibric	244-274	
Hemic	274-304	
Fibric	304-396	
Hemic	396-487	
Silty	487+	
clay		

Location: 20 feet S and 2600 feet E of the NW corner of Sec. 8, T.69N., R.22W. (NE1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 8, T.69N., R.22W.)

Vegetation: Scattered tamarack and black spruce; sparse understory consists of some bog rosemary, leatherleaf, speckled alder, sedges, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: B. Sether and J. Dahl on March 14, 1980.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-213
Fibric	213-274
Hemic	274-517
Loamy	517+
sand	
with	
pebbles	

Location: 20 feet S and 2350 feet W of the NE corner of Sec. 9, T.69N., R.22W. (NW1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 9, T.69N., R.22W.)

Vegetation: Scattered black spruce; sparse understory consists of some speckled alder, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some snowberry.

Microrelief: 10 cm.

Depth To Water Table: Not recorded.

Described By: H. Mooers and D. Haverkost on March 14, 1980.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-487
Sapric	487-608
Silty	608+
clay	

Location: 1140 feet S and 750 feet E of the NW corner of Sec. 9. T.69N., R.22W. (SW1/4 of SE1/4 of NW1/4 of NW1/4, Sec. 9, T.69N., R.22W.)

Vegetation: Black spruce crown cover of about 95 percent with scattered tamarack; understory consists of grasses and some leatherleaf, swamp laurel, and bog rosemary; ground cover consists of Sphagnum mosses with some snowberry.

Microrellef: 15 cm.

Microrelief: 15 cm.

Depth To Water Table: 15 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-548+
Bottom	
unknown	

Location: 2550 feet W and 20 feet S of the NE corner of Sec. 10, T.69N., R.22W.

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack

sists of some Sphagnum and feather mosses, snowberry, and cranberry.

Described By: H. Mooers, B. Sether, J. Dahl, and D. Haverkost on March 14, 1980.

and northern white cedar; sparse understory consists of some willow, speckled

alder, Labrador tea, leatherleaf, bog rosemary, and sedges; ground cover con-

(NW1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 10, T.69N., R.22W.)

Location: 2350 feet S and 20 feet E of the NW corner of Sec. 9, T.69N., R.22W. (SW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 9, T.69N., R.22W.)

Vegetation: Black spruce crown cover of about 40 percent; sparse understory consists of some leatherleaf and swamp laurel; ground cover consists mostly of

Microrelief: 15 cm.

Depth To Water Table: At surface.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-335
Fibric	335-396
Hemic	396-457
Fibric	457-487
Hemic	487-669
Fibric	669-700
Sapric	700-775
Silty	775+
clay	

Sphagnum mosses with some snowberry and pitcher plant.

Described By: H. Mooers, B. Sether, S. Nelson, and D. Haverkost on March 17,

Location: 330 feet S and 2300 feet E of the NW corner of Sec. 17, T.69N., R.22W. (SW1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 17, T.69N., R.22W.)

Vegetation: Scattered black spruce; sparse understory consists of Labrador tea, leatherleaf, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some snowberry.

Microrellef: 15 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Fibric	0-183
Hemic	183-366
Fine	366+
sand	

Layer	Depth (cm)	
Fibric	0-152	
Hemic	152-548+	
Bottom		
unknown		

Location: 1050 feet N and 20 feet E of the SW corner of Sec. 9. T.69N., R.22W. (NW1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 9, T.69N., R.22W.)

Vegetation: Black spruce crown cover of about 65 percent; understory consists of Labrador tea with some grasses, leatherleaf, bog rosemary, and swamp laurel;

ground cover consists mostly of Sphagnum mosses. Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: L. Severson, B. Sether, and S. Nelson on April 2, 1980.

Hemic 0-122 Fibric 122-213 Hemic 213-427 Fibric 427-457 Hemic 457-487 Fibric 487-517 Hemic 517-700 Silty 700+	Layer	Depth (cm)
IOam	Fibric Hemic Fibric Hemic Fibric Hemic Silty	122-213 213-427 427-457 457-487 487-517 517-700

Location: 1250 feet S and 2300 feet W of the NE corner of Sec. 19, T.70N., R.22W. (SW1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 19, T.70N., R.22W.)

Vegetation: Black spruce crown cover of about 75 percent; understory consists of Labrador tea and leatherleaf with some swamp laurel and grasses; ground cover consists mostly of Sphagnum mosses with some snowberry.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 27, 1979.

ayer	Depth (cm)
ibric	0- 91
emic	91-183
lay	183+

Location: 750 feet N and 560 feet E of the SW corner of Sec. 20, T.70N., B.22W. (SE1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 20, T.70N., R.22W.)

Vegetation: Scattered black spruce and paper birch; understory consists of speckled alder, willow, Labrador tea, grasses, and cattails; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 46 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer	Depth (cm)
Hemic	0-30
Clay	30+

Location: 1700 feet S and 640 feet E of the NW corner of Sec. 33, T.70N., R.22W.

Vegetation: Scattered black spruce; understory consists of leatherleaf and grasses

Location: 2350 feet S and 1625 feet W of the NE corner of Sec. 17, T.63N., R.23W.

Vegetation: Black spruce crown cover of about 60 percent with scattered

tamarack; sparse understory consists of some Labrador tea; ground cover

consists mostly of Sphagnum mosses with some club mosses, cranberry, and

(SW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 17, T.63N., R.23W.)

with some swamp laurel and bog rosemary; ground cover consists mostly of

(SE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 33. T.70N., R.22W).

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer

Fibric Hemic Sandy clay

Sphagnum mosses with some cranberry.

Depth To Water Table: At surface.

Microrellef: 22 cm

blueberry.

Microrellef: 25 cm.

Depth To Water Table: Not Visible.

Location: 1480 feet S and 1850 feet F of the NW corner of Sec. 32, T.70N, R.22W. (NE1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 32, T.70N., R.22W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; understory consists of leatherleaf, Labrador tea, bog rosemary, and grasses; ground cover consists of Sphagnum mosses with some feather mosses and snowberry.

Microrelief: 20 cm

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Fibric	0-183
Hemic	183-457
Sapric	457-518
Sandy	518+
clay	

Location: 1900 feet N and 180 feet W of the SE corner of Sec. 4, T.63N., R.23W. (SE1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 4, T.63N., R.23W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered northern white cedar, paper birch, and aspen; sparse understory consists of some Labrador tea, sedges, and grasses; ground cover consists of Sphagnum

Microrelief: 20 cm.

Depth To Water Table: 30 cm.

Described By: H. Mooers and B. Sether on April 11, 1980.

Depth (cm)	
0-183	Layer
183-427	Hemio
427+	Sand

mosses with some other mosses, cranberry, snowberry, and bunchberry.

Layer	Depth (cm)
Hemic	0-122
Sand	122+

Location: 1070 feet S and 550 feet W of the NE corner of Sec. 18, T.63N., R.23W. (SW1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 18, T.63N., R.23W.) Vegetation: Cut over area; consists mostly of grasses with some willow.

Microrellef: 5 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 20, 1980.

Layer	Depth (cm)
Hemic Clay Ioam	0-183 183+

Location: 2500 feet N and 20 feet W of the SE corner of Sec. 32, T.70N., R.22W. (NE1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 32, T.70N., R.22W.)

Vegetation: Scattered black spruce and tamarack: sparse understory consists of some leatherleaf, swamp laurel, bog rosemary, and grasses; ground cover consists mostly of Sphagnum mosses with some snowberry.

Microrelief: 7 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers, B. Sether, and S. Nelson on March 20, 1980.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-152
Fibric	152-183
Hemic	183-213
Fibric	213-244
Hemic	244-274
Fibric	274-335
Hemic	335-578
Clay	578+

Location: 350 feet N and 1080 feet W of the SE corner of Sec. 14, T.63N., R.23W. (NW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 14, T.63N., R.23W.)

Vegetation: Black spruce crown cover of about 85 percent; sparse understory consists of some Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm

Depth To Water Table: Not visible.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm)
Hemic	0-183
Silty	183+
clav	

Location: 1180 feet S and 2400 feet W of the NE corner of Sec. 21, T.63N., R.23W. (SW1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 21, T.63N., R.23W.)

Vegetation: Northern white cedar crown cover of about 35 percent with scattered paper birch, ash, and balsam fir; lush understory consists mostly of grasses with some bog birch, willow, alder, sedges, reeds, and cattails.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 20, 1980.

Layer	Depth (cm)
Hemic	0- 90
Sapric	90-165
Sandy	165+
loam	

Depth (cm) Layer Hemic 0-213 Sapric 213-274 Hemic 274-518 518+ Silty clay

Described By: H. Mooers and B. Sether on April 11, 1980.

Location: 1920 feet N and 1460 feet W of the SE corner of Sec. 29, T.64N., R.23W. (NE1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 29, T.64N., R.23W.)

Vegetation: Scattered black spruce, tamarack, paper birch, and balsam; lush understory consists mostly of alder with some sedges, grasses, Labrador tea, and dogwood; ground cover consists of some Sphagnum mosses, other mosses, cranberry, and snowberry.

Microrellef: 20 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm
Hemic	0-396
Sand	396+

Location: 1000 feet S and 2240 feet W of the NE corner of Sec. 34, T.63N., R.23W. (SE1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 34, T.63N., R.23W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered balsam, northern white cedar, paper birch, and aspen; ground cover consists mostly of Sphagnum mosses.

Microrellef: 10 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and D. Haverkost on May 20, 1980.

Layer	Depth (cm)
Muck	0-12
Loam	12+

Location: 1450 feet S and 1180 feet E of the NW corner of Sec. 34, T.64N., R.23W. (NE1/4 of NE1/4 of SW1/4 of NW1/4, Sec. 34, T.64N., R.23W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; sparse understory consists of Labrador tea with some alder; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm)
Silty	0+

Location: 750 feet N and 1190 feet W of the SE corner of Sec. 20, T.66N., R.23W. (SW 1/4 of NW1/4 of SE1/4 of SE1/4, Sec. 20, T.66N., R.23W.)

Vegetation: Northern white cedar crown cover of about 70 percent with scattered paper birch and black spruce: understory consists of Labrador tea with some speckled alder and grasses; ground cover consists of feather mosses.

Microrellef: 25 cm.

Depth To Water Table: Near surface.

Described By: R. Graber and D. Haverkost on November 28, 1979.

Layer	Depth (cm)
Hemic	0-335
Limnic	335-366
Medium sand	366+

Location: 775 feet S and 2050 feet W of the NE corner of Sec. 25, T.66N., R.23W. (Gov Lot 2, Sec. 25, T.66N., R.23W.)

Vegetation: Black spruce crown cover of about 95 percent with scattered tamarack, paper birch, and aspen; understory consists of Labrador tea, leatherleaf, and some cotton grass; ground cover consists of Sphagnum mosses with some snowberry.

Microrellef: 30 cm.

Depth To Water Table: Not Visible.

Described By: R. Graber and D. Haverkost on November 28, 1979.

Layer	Depth (cm)
Hemic Clay Ioam	0-91 91+

Location: 280 feet N and 2100 feet W of the SE corner of Sec. 28, T.67N., R.23W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 28, T.67N., R.23W.) Vegetation: Black spruce crown cover of about 85 percent; understory consists of

Vegetation: Black spruce crown cover of about 85 percent; understory consists of Labrador tea; ground cover consists of Sphagnum mosses, feather mosses, and cranberry.

Microrellef: 15 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 28, 1979.

Layer	Depth (cm)
Hemic	0-122
Limnic	122-152
Silty	152+
clay	
loam	

Location: 2150 feet S and 180 feet E of the NW corner of Sec. 19, T.69N., R.23W. (Gov. Lot 2, Sec. 19, T.69N., R.23W.)

Vegetation: Black spruce crown cover of about 95 percent with scattered tamarack; lush understory consists of leatherleaf, Labrador tea, swamp laurel, bog rosemary, cotton grass and some grasses; ground cover consists of Sphagnum mosses.

Microrellef: 20 cm.

Depth To Water Table: Near surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer Depth (cm)
Hemic 0-427
Clay 427+

Location: 740 feet S and 50 feet W of the NE corner of Sec. 28. T.69N., R.23W. (NE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 28, T.69N., R.23W.)

Vegetation: Black spruce crown cover of about 95 percent with scattered tamarack; lush understory consists mostly of leatherleaf with swamp laurel, grasses, and some bog rosemary and cotton grass; ground cover consists of Sphagnum mosses.

Microrelief: 25 cm.

Depth To Water Table: Near surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Hemic	0-548+
Bottom	
unknown	

Location: 580 feet S and 1800 feet E of the NW corner of Sec. 29, T.69N., R.23W. (SE1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 29, T.69N., R.23W.)

Vegetation: Black spruce and tamarack crown cover of about 85 percent; lush understory consists of leatherleaf, grasses, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-366
Clay	366+

Location: 1140 Feet S and 490 feet W of the NE corner of Sec. 3, T.70N., R.23W. (Gov. Lot 1, Sec. 3, T.70N., R.23W.)

Vegetation: Scattered black ash, black spruce, and balsam fir; sparse understory consists of willow, forbs, and grasses; ground cover consists of some cranberry.

Microrelief: Not recorded.

Sphagnum mosses.

Depth To Water Table: Not visible.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: D. Mellem and R. Graber on November 27, 1979.

Layer	Depth (cm)
Hemic	0-15
Silty	15+
clay	

Location: 490 feet N and 2500 feet E of the SW corner of Sec. 10, T.70N., R.23W.

Vegetation: Scattered tamarack and black spruce; lush understory consists mostly

of grasses with leatherleaf and swamp laurel; ground cover consists mostly of

Depth (cm)

0- 61

61-259

259+

(NE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 10, T.70N., R.23W.)

Described By: D. Mellem and P. Guertin on November 27, 1979.

Laver

Fibric

Hemic

Clay

Location: 950 feet S and 2450 feet E of the NW corner of Sec. 5, T.70N., R.23W. (Gov. Lot 3, Sec. 5, T.70N., R.23W.)

Vegetation: Northern white cedar crown cover of about 75 percent with scattered tamarack, paper birch, and aspen; lush understory consists of bog birch, sedges, grasses, and some speckled alder, willow, leatherleaf, and cattails; ground cover consists of Sphagnum mosses with some other mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)	
Hemic	0-366	
Sapric	366-442	
Clay	442+	

Location: 1020 feet N and 640 feet W of the SE corner of Sec. 14, T.70N., R.23W. (NW1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 14, T.70N., R.23W.)

Vegetation: Black spruce crown cover of about 30 percent; sparse understory consists of leatherleaf, Labrador tea, swamp laurel, bog rosemary, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 46 cm.

Depth To Water Table: 30 cm.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer	Depth (cm
Fibric	0-122
Hemic	122-274
Clay	274+
	Fibric Hemic

Location: 450 feet S and 150 feet W of the NE corner of Sec. 15, T.70N., R.23W.

Location: 1600 feet N and 2450 feet W of the SE corner of Sec. 9, T.70N., R.23W.

Vegetation: Scattered black spruce, poplar, and tamarack; sparse understory consists of some leatherleaf, swamp laurel, bog rosemary, sedges, and grasses;

ground cover consists mostly of Sphagnum mosses with some cranberry.

Depth (cm)

0-320

320-350

350-366 366+

(SW1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 9, T.70N., R.23W.)

Described By: R. Graber and D. Haverkost on November 27, 1979.

snowberry, false Solomon's seal, and pitcher plant.

Laver

Fibric

Hemic

Sapric

Sandy clay

Vegetation: Scattered poplar, black spruce, and tamarack; sparse understory consists of some sedges, leatherleaf, swamp laurel, and bog rosemary; ground cover consists of Sphagnum mosses, feather mosses, and cranberry.

Microrelief: 45 cm.

Microrellef: 15 cm.

Depth To Water Table: Not Visible.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 27, 1979.

(SE1/4 of NE1/4 of NE1/4, Sec. 15, T.70N., R.23W.)

Layer	Depth (cm)
Hemic Clay loam	0-290 290+

Location: 690 feet N and 2210 feet E of the SW corner of Sec. 15, T.70N., R.23W. (SW1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 15, T.70N., R.23W.)

Vegetation: Scattered black spruce and tamarack; understory consists mostly of grasses with some leatherleaf, swamp laurel, bog rosemary, and fern; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not Visible.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer

Fibric Sandy clay with pebbles Location: 1790 feet N and 140 feet E of the SW corner of Sec. 20, T.70N., R.23W. (NW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 20, T.70N., R.23W.)

Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf, Labrador tea, bog rosemary and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Depth (cm)	Layer	Depth (cm)
0-107	Fibric	0-305
107+	Hemic	305-442+
	Bottom	
	unknown	

Location: 1340 feet S and 170 feet W of the NE corner of Sec. 21, T.70N., R.23W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 21, T.70N., R.23W).

Vegetation: Black spruce crown cover of about 45 percent; understory consists of Labrador tea, leatherleaf, grasses, and some swamp laurel and bog rosemary; ground cover consists of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 30 cm.

Described By: D. Mellem and P. Guertin on November 27, 1979.

ayer	Depth (cm)
bric emic	0- 91 91-335
lay	335+

Vegetation: Scattered black spruce, tamarack, and poplar; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, bog rosemary, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 60 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

 Layer
 Depth (cm)

 Hemic
 0-183

 Sapric
 183-213

 Clay
 213+

Location: 1840 feet N and 1315 feet W of the SE corner of Sec. 25, T.70N., R.23W. (Gov. Lot 3, Sec. 25, T.70N., R.23W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; understory consists of Labrador tea and leatherleaf with some swamp laurel and grasses; ground cover consists of Sphagnum mosses with some feather mosses and snowberry.

Microrelief: 20 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Hemic Sampler would not penetrate	0-183 183+

Location: 150 feet S and 480 feet W of the NE corner of Sec. 29, T.70N., R.23W. (NW1/4 of NE1/4 of NE1/4, Sec. 29, T.70N., R.23W.)
Vecetation: Scattered black spruce, poplar, and tamarack: understory consists of

Vegetation: Scattered black spruce, poplar, and tamarack; understory consists of grasses with some speckled alder and fern; ground cover consists of Sphagnum mosses with some other mosses.

Microrelief: Negligible.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)	
Hemic	0-411	
Clay	411+	

Location: 2420 feet N and 400 feet E of the SW corner of Sec. 33, T.70N., R.23W. (NE1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 33, T.70N., R.23W.)

Vegetation: Scattered black spruce and aspen; understory consists of speckled alder, bog birch, willow, Labrador tea, leatherleaf, and some grasses; ground cover consists of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and R. Graber on November 27, 1979.

Layer	Depth (cm)
Hemic	0-244
Clay	244+

Location: 790 feet S and 1780 feet W of the NE corner of Sec. 13, T.63N., R.24W. (NW1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 13, T.63N., R.24W.)

Vegetation: Scattered black spruce, northern white cedar, and paper birch; sparse understory consists of some Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 15 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and D. Haverkost on May 20, 1980.

Layer	Depth (cm)
Hemic	0-335
Loam	335+

Location: 600 feet S and 1380 feet W of the NE corner of Sec. 14, T.63N., R.24W. (SE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 14, T.63N., R.24W.)

Vegetation: Northern white cedar crown cover of about 35 percent with scattered ash; lush understory consists mostly of grasses with some bog birch, cow slips, and sedges; ground cover consists of some Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on May 20, 1980.

.ayer	Depth (cm
lemic	0-213
.oam	213+

Location: 1715 feet N and 1740 feet W of the SE corner of Sec. 26, T.63N., R.24W. (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 26, T.63N., R.24W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, swamp laurel, bog rosemary, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 11, 1980.

unknown

Location: 1050 feet S and 950 feet W of the NE corner of Sec. 27, T.63N., R.24W. (SW1/4 of SW1/4 of NE1/4, Sec. 27, T.63N., R.24W.)

Venetation: Scattered black source and tamerack: sparse understory consists of

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, swamp laurel, bog rosemary, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 25 cm.

Depth To Water Table: 8 cm.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm)
Hemic Bottom unknown	0-609+

Location: 350 feet N and 2400 feet E of the SW corner of Sec. 28, T.63N, R.24W. (NE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 28, T.63N., R.24W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, swamp laurel, bog rosemary, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm)
Hemic	0-213
Sapric	213-244
Clay	244+

Layer Depth (cm)
Fibric 0- 61
Hemic 61-479+
Bottom

Location: 240 feet N and 1825 feet E of the SW corner of Sec. 31, T.65N., R.24W. (SE1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 31, T.65N., R.24W.)

Vegetation: Scattered tamarack and black spruce; sparse understory consists of some leatherleaf, swamp laurel, and bog rosemary; ground cover consists of Sphagnum mosses with some other mosses and cranberry.

Microrellet: 20 cm.

Depth To Water Table: Near surface.

mosses with some snowberry.

Depth To Water Table: Not visible.

Microrellef: 30 cm.

Described By: H. Mooers and B. Sether on April 11, 1980.

Laver Depth (cm) Hemic 0-427 +Bottom unknown

Location: 1480 feet S and 150 feet E of the NW corner of Sec. 5, T.66N., R.24W.

Vegetation: Scattered black spruce and poplar; sparse understory consists of

some Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum

Depth (cm) 0-229

229-244

244+

(NW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 5, T.66N., R.24W.)

Described By: R. Graber and D. Haverkost on November 28, 1979.

Laver

Hemic Sapric

Fine

sand

Location: 1675 feet N and 1800 feet W of the SE corner of Sec. 31, T.65N., R.24W. (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 31, T.65N., R.24W.)

Vegetation: Scattered black spruce; sparse understory consists of some grasses, leatherleaf, swamp laurel, bog rosemary, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm)
Hemic	0-366
Medium	366+
to	
coarse	
sand	

Location: 1475 feet S and 560 feet W of the NE corner of Sec. 11, T.66N., R.24W. (NW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 11, T.66N., R.24W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of Labrador tea and leatherleaf, with some swamp laurel and grasses; ground cover consists mostly of Sphagnum mosses and some cranberry.

Microrellef: 22 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and R. Graber on November 28, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-183
Silt	183+
loam	

Location: 175 feet N and 2500 feet W of the SE corner of Sec. 13, T.66N., R.24W.

(SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 13, T.66N., R.24W.)

Location: 1550 feet S and 75 feet W of the NE corner of Sec. 3, T.66N., R.24W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 3, T.66N., R.24W.)

Vegetation: Scattered black spruce; sparse understory consists of some Labrador

Sphagnum mosses with some cranberry.

Described By: D. Mellem and P. Guertin on November 28, 1979.

Laver

Fibric

Hemic

Sandv

clay loam

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

tea, leatherleaf, swamp laurel, and grasses; ground cover consists mostly of

Depth (cm)

0- 61

61-305

305+

Vegetation: Black spruce crown cover of about 95 percent; sparse understory consists of some Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: D. Mellem and P. Guertin on November 28, 1979.

Layer	Depth (cm)	
Hemic	0-152	
Loam	152+	

Location: 1000 feet S and 1050 feet W of the NE corner of Sec. 16, T.66N., R.24W. (SW1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 16, T.66N., R.24W.)

Vegetation: Scattered black spruce and paper birch; sparse understory consists of some Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses and snowberry.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 28, 1979.

Location: 50 feet S and 2025 feet W of the NE corner of Sec. 20, T.66N., R.24W. (NE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 20, T.66N., R.24W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of mosses with some Sphagnum mosses and snowberry.

Microrellef: 15 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 28, 1979.

Layer	Depth (cm)	Layer	Depth (cm)
Hemic Medium sand	0-76 76+	Hemic Fine sand	0-91 91+

Location: 2425 feet S and 2040 feet W of the NE corner of Sec. 28, T.66N., R.24W. (SE1/4 of SW1/4 of SW1/4 of NE1/4, Sec. 28, T.66N., R.24W.)

Vegetation: Scattered black spruce, tamarack, and paper birch; sparse understory consists of bog birch, Labrador tea, leatherleaf, and grasses; ground cover consists of Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: 122 cm.

Described By: D. Mellem and P. Guertin on November 28, 1979.

Layer	Depth (cm)
Hemic	0-107
Limnic	107-152
Gravel	152+
and	
sand	

Location: 125 feet S and 1800 feet W of the NW corner of Sec. 12, T.67N., R.24W. (NF1/4 of NW1/4 of NF1/4 of NW1/4, Sec. 12, T.67N., R.24W.)

Vegetation: Scattered black spruce; understory consists of Labrador tea. leatherleaf, bog rosemary, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 152 cm.

Described By: D. Mellem and P. Guertin on November 28, 1979.

Layer	Depth (cm
Hemic	0-427
Limnic	427-518
Silty	518+
clay	
loam	

Location: 1450 feet N and 75 feet W of the SE corner of Sec. 19, T.67N., R.24W. (SE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 19, T.67N., R.24W.)

Vegetation: Black spruce crown cover of about 40 percent, understory consists of Labrador tea with some leatherlear, ground cover consists mostly of Sphagnum mosses.

Microrellef: 22 cm.

Depth To Water Table: Not recorded.

Described By: D. Mellem and P. Guertin on November 28, 1979.

Depth (cm)
0-183
183+

Location: 845 feet N and 170 feet W of the SE corner of Sec. 4, T.68N., R.24W. (SE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 4, T.68N., R.24W.)

Vegetation: Black spruce crown cover of about 80 percent; lush understory con-

Vegetation: Black spruce crown cover of about 80 percent; lush understory consists mostly of Labrador tea with some leatherleaf; ground cover consists mostly of Sphagnum mosses with some feather mosses, lingen berry, cranberry, and snowberry.

Microrelief: 35 cm.

Depth To Water Table: 20 cm.

Described By: R. Graber and D. Haverkost on November 16, 1979.

Layer	Depth (cm)	
Fibric	0-122	
Hemic	122-305	
Sapric	305-335	
Hemic	335-427	
Clav	427+	

Location: 1750 feet N and 2350 feet E of the SW corner of Sec. 6, T.68N., R.24W. (NE1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 6, T.68N., R.24W.)

Vegetation: Scattered black spruce; sparse understory consists of leatherleaf and swamp laurei; ground cover consists mostly of Sphagnum mosses with cranberry and some other mosses.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on November 16, 1979.

Layer	Depth (cm)
Hemic	0-183
Silty	183+
clay	
with	
pebbles	

Location: 2125 feet S and 1185 feet W of the NE corner of Sec. 35, T.68N., R.24W. (NW1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 35, T.68N., R.24W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; lush understory consists mostly of leatherleaf with some Labrador tea and bog rosemary; ground cover consists of some Sphagnum mosses. Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 28, 1979.

Layer	Depth (cm)
Hemic	0-274
Sapric	274-290
Sandy	290+
clay	

Location: 1125 feet S and 150 feet E of the NW corner of Sec. 8, T.69N., R.24W. (SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 8, T.69N., R.24W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly

egetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some Labrador tea, leatherleaf, bog rosemary, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 35 cm.

Depth To Water Table: At surface.

Described By: D. Olson and R. Graber on November 16, 1979.

.ayer	Depth (cm)
apric	0-138
lemic	138-457
apric	457-518
Clay	518+

Location: 800 feet N and 1750 feet E of the SW corner of Sec. 12, T.69N., R.24W. (SE1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 12, T.69N., R.24W.)

Vegetation: Scattered black spruce; understory consists of Labrador tea, leatherleaf, swamp laurel, bog rosemary, and some grasses; ground cover consists of Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-259 259+

Location: 210 feet N and 950 feet E of the SW corner of Sec. 13, T.69N., R.24W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 13, T.69N., R.24W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of leatherleaf, swamp laurel, and bog rosemary; ground cover consists of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: 30 cm.

Described By: D. Mellem and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-183
Sandy	183+
clav	

Location: 195 feet N and 260 feet E of the SW corner of Sec. 14, T.69N., R.24W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 14, T.69N., R.24W.)

Vegetation: Black spruce crown cover of about 75 percent; sparse understory consists of some dogwood, Labrador tea, and grasses; ground cover consists of some snowberry.

Microrellef: 46 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer	Depth (cm)
Hemic Silty	0-107 107+
clav	

Location: 550 feet S and 2000 feet W of the NE corner of Sec. 19, T.69N., B.24W. (SE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 19, T.69N., R.24W.)

Vegetation: Scattered black spruce; understory consists mostly of grasses with some swamp laurel and leatherleaf; ground cover consists mostly of Sphagnum mosses

Microrellef: 30 cm.

Microrelief: 30 cm.

Depth To Water Table: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on November 16, 1979.

Layer Depth (cm) 0- 61 Fibric 61-229 Hemic Silty 229+ clay

Location: 1825 feet N and 110 feet W of the SE corner of Sec. 25, T.69N., R.24W.

Vegetation: Black spruce crown cover of about 75 percent with scattered

tamarack; understory consists of grasses, leatherleaf, and some swamp laurel;

(NE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 25, T.69N., R.24W.)

ground cover consists mostly of Sphagnum mosses.

Layer

Hemic

Sandy clay

Described By: D. Mellem and P. Guertin on November 27, 1979.

Location: 2500 feet S and 525 feet W of the NE corner of Sec. 21, T.69N., B.24W. (SW1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 21, T.69N., R.24W.)

Vegetation: Scattered black spruce; understory consists of Labrador tea with some leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's coal

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on November 16, 1979.

Layer	Depth (cm)
Hemic	0-183
Clay	183+

Location: 1740 feet N and 2475 feet E of the SW corner of Sec. 27, T.69N., R.24W. (NE1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 27, T.69N., B.24W.)

Vegetation: Scattered aspen in a clearing surrounded by black spruce; understory consists mostly of cotton grass with some Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses with some feather mosses and false Solomon's seal.

Microrelief: Negligible.

Depth To Water Table: At surface.

Described By: D. Olson and R. Graber on November 16, 1979.

Depth (cm)		
0-320	Layer	Depth (cm)
320+	Hemic Siltv	0-138 138+
	clay	

Location: 300 feet S and 945 feet W of the NE corner of Sec. 31, T.69N., R.24W. (NE1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 31, T.69N., R.24W.)

Location: 655 feet N and 540 feet E of the SW corner of Sec. 21, T.69N., R.24W.

Vegetation: Scattered black spruce; lush understory consists of swamp laurel,

leatherleaf and cotton grass with some sedges and bog rosemary; ground

cover consists mostly of Sphagnum mosses with false Solomon's seal and

Depth (cm)

0-274

274-295

295+

(SE1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 21, T.69N., R.24W.)

Described By: D. Olson and R. Graber on November 16, 1979.

Laver

Hemic

Sapric

Clav

some other mosses and cranberry.

Microrelief: 55 cm.

Depth To Water Table: 15 cm.

Vegetation: Scattered black spruce; lush understory consists mostly of grasses with bog rosemary and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on November 16, 1979.

Layer	Depth (cm				
Fibric	0-122				
Hemic	122-320				
Limnic	320+				
with					
silt					

Location: 2550 feet S and 410 feet W of the NE corner of Sec. 33, T.69N., R.24W. (SW1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 33, T.69N., R.24W.)

Vegetation: Scattered black spruce; lush understory consists of sedges, Labrador tea, leatherleaf, and some grasses, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 15 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and D. Haverkost on November 16, 1979.

clay

Location: 1880 feet N and 160 feet W of the SE corner of Sec. 36, T.69N., R.24W. (SE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 36, T.69N., R.24W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer	Depth (cm)		Layer	Depth (cm)
Fibric	0-122		Hemic	0-427
Hemic	122-442		Sapric	427-457+
Sandy	442+		Bottom	
clav			unknown	

Location: 875 feet N and 120 feet W of the SE corner of Sec. 13, T.70N., R.24W. (SE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 13, T.70N., R.24W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, bog rosemary, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 30 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-548+
Bottom	
unknown	

Location: 550 feet S and 1300 feet E of the NW corner of Sec. 22, T.70N., R.24W. (SW1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 22, T.70N., R.24W.)

Vegetation: Black spruce crown cover of about 95 percent; lush understory consists of leatherleaf, bog rosemary, cotton grass, Labrador tea, and some sedges; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer Depth (cm) 0-518 Hemic Clay 518+

Location: 1475 feet S and 220 feet W of the NE corner of Sec. 28, T.70N., R.24W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 28, T.70N., R.24W.)

Vegetation: Scattered black spruce; lush understory consists of leatherleaf, grasses, and some cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 27, 1979.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-442
Silty	442+
clay	

Location: 1000 feet N and 2700 feet W of the SE corner of Sec. 18, T.63N., R.25W.

Vegetation: Scattered ash, black spruce, and paper birch; lush understory consists mostly of grasses with some alder and raspberry; ground cover consists of

Microrellef: 10 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and D. Haverkost on May 20, 1980.

Depth (cm) Layer Hemic 0-213 213 +Loam

Location: 600 feet S and 1250 feet E of the NW corner of Sec. 10, T.63N., R.25W.

Vegetation: Black spruce crown cover of about 35 percent; sparse understory con-

sists of some leatherleaf, Labrador tea, swamp laurel, and grasses; ground

cover consists mostly of Sphagnum mosses with some cranberry and false

(SE1/4 of NE1/4 of NW1/4 of NW1/4, Sec. 10, T.63N., R.25W.)

Described By: H. Mooers and B. Sether on April 11, 1980.

(NE1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 18, T.63N., R.25W.)

some Sphagnum and other mosses.

 -y	. 10100013	unu	υ.	Havorkost	OII	way	20,	1300

Layer	Depth (cm)
Hemic Silty	0-335 335+
clay	0001

Location: 500 feet N and 955 feet W of the SE corner of Sec. 8, T.64N., R.25W. (NW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 8, T.64N., R.25W.)

Vegetation: Scattered tamarack; understory consists of some bog birch, grasses, leatherleaf, Labrador tea, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Solomon's seal.

Depth To Water Table: 30 cm.

Microrellef: 25 cm.

Depth To Water Table: Not recorded.

Described By: H. Mooers and B. Sether on April 10, 1980.

Depth (cm)
0- 61
61-274
274+

Location: 550 feet N and 1990 feet W of the SE corner of Sec. 9, T.64N., R.25W. (NE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 9, T.64N., R.25W.)

Vegetation: Scattered tamarack; understory consists of some grasses, leatherleaf, bog rosemary, swamp laurel, sedges, and bog birch; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 50 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-122
Fibric	122-396
Hemic	396+

Location: 500 feet N and 925 feet E of the SW corner of Sec. 32, T.70N., R.24W. (NW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 32, T.70N., R.24W.)

Vegetation: Black spruce and tamarack crown cover of about 85 percent; lush understory consists of speckled alder. Labrador tea, bog rosemary, leatherleaf. and some grasses; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 20 cm.

Depth To Water Table: Near surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm)
Fibric	0-229
Hemic	229-442
Sandy	442+
clay	

Location: 1800 feet N and 1620 feet W of the SE corner of Sec. 2, T.64N., R.25W. (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 2, T.64N., R.25W.)

Vegetation: Sparse understory consists of some grasses, black spruce, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 20 cm.

Depth To Water Table: 10 cm.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm)
Fibric	0-244
Hemic	244-396+
Bottom	
unknown	

Location: 1290 feet S and 885 feet E of the NW corner of Sec. 11, T.64N., R.25W. (SW1/4 of SE1/4 of NW1/4 of NW1/4, Sec. 11, T.64N., R.25W.)

Vegetation: Lush understory consists mostly of sedges with some tamarack, black spruce, bog birch, leatherleaf, and bog rosemary; ground cover consists of some Sphagnum mosses, other mosses, cranberry, and pitcher plant.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Fibric	0-213
Hemic	213-411
Silty	411+
clav	

Location: 910 feet N and 1550 feet W of the SE corner of Sec. 12, T.64N., R.25W. (NE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 12, T.64N., R.25W.)

Vegetation: Understory consists of some tamarack, bog birch, cotton grass, leatherleaf, and bog rosemary, ground cover consists of some Sphagnum and other mosses.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 11, 1980.

Hemic 0-183 Sand 183+ Location: 1120 feet N and 950 feet E of the SW corner of Sec. 18, T.64N., R.25W. (NW1/4 of NE1/4 of SW1/4 of SW1/4, Sec. 18, T.64N., R.25W.)

Vegetation: Tamarack crown cover of about 30 percent with about 30 percent crown cover of northern white cedar and scattered black spruce; understory consists of Labrador tea with some leatherleaf and bog birch; ground cover consists mostly of Sphagnum mosses with some feather mosses, cranberry, and false Solomon's seal.

Microreliet: 30 cm.

Depth To Water Table: 10 cm.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Hemic	0-183
Sapric	183-213
loam	213+

Location: 60 feet S and 290 feet E of the NW corner of Sec. 5, T.65N., R.25W. (Gov. Lot 4, Sec. 5, T.65N., R.25W.)

Vegetation: Tamarack crown cover of about 75 percent with about 25 percent black spruce; lush understory consists of speckled alder, Labrador tea. leatherleaf, bog rosemary, and sedges with some grasses and cattails, ground cover consists of Sphagnum mosses.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Layer	Depth (cm)
Hemic	0-152
Sandy	152+
clav	

Location: 1415 feet N and 1915 feet W of the SE corner of Sec. 31, T.64N., R.25W. (SW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 31, T.64N., R.25W.)

Vegetation: Scattered tamarack and black spruce; understory consists of some

Vegetation: Scattered tamarack and black spruce; understory consists of some bog birch, Labrador tea, sedges, grasses, and ferns; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 35 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm
Hemic	0- 91
Fibric	91-122
Hemic	122-305
Silt	305+
loam	

Location: 540 feet N and 150 feet E of the SW corner of Sec. 2, T.65N., R.25W. (NW1/4 of SW1/4 of SW1/4, Sec. 2, T.65N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 15 cm.

Depth To Water Table: Not visible.

Described By: D. Haverkost and R. Graber on November 28, 1979.

Hemic 0-305 Medium 305+ sand Location: 250 feet S and 1260 feet W of the NE corner of Sec. 7, T.65N., R.25W. (NE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 7, T.65N., R.25W.)

Vegetation: Black spruce crown cover of about 65 percent with about 25 percent tamarack and scattered aspen; lush understory consists of Labrador tea and grasses with some sedges and speckled alder; ground cover consists of Sphagnum mosses with some snowberry.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Layer	Depth (cm)
Hemic Medium sand with minor clay	0-213 213+

Location: 1290 feet S and 1840 feet W of the NE corner of Sec. 27, T.65N., R.25W. (NW1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 27, T.65N., R.25W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some sedges, grasses, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 25 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and B. Sether on April 10, 1980.

Fibric Hemic Fibric

Bottom

Location: 240 feet S and 200 feet E of the NW corner of Sec. 32, T.65N., R.25W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 32, T.65N., R.25W.)

Vegetation: Black spruce crown cover of about 60 percent; sparse understory consists of some Labrador tea, bog rosemary, bog birch, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 25 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and B. Sether on April 10, 1980.

Depth (cm)	Layer	Depth (cr
0- 61	Hemic	0-122
61- 91	Sand	122+
91-427+		

Location: 1850 feet N and 550 feet W of the SE corner of Sec. 35, T.65N., R.25W. (NW1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 35, T.65N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of some grasses, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrellef: 25 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-427+
Bottom	
unknown	

Location: 610 feet N and 2450 feet W of the SE corner of Sec. 6, T.66N., R.25W. (NW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 6, T.66N., R.25W.)

Vegetation: Black spruce crown cover of about 60 percent with about 30 percent crown cover of tamarack; lush understory consists of grasses with some leatherleaf, bog rosemary, Labrador tea, and speckled alder; ground cover consists of Sphagnum mosses.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Layer	Depth (cm)
Hemic	0-183
Medium	183+
and	
coarse	
sand	

Location: 1875 feet N and 90 feet W of the SE corner of Sec. 8, T.66N., R.25W. (NE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 8, T.66N., R.25W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack and aspen; understory consists of grasses and leatherleaf with some bog birch, cattails and Labrador tea; ground cover consists of Sphagnum mosses.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 29, 1979.

Layer	Depth (cm)
Hemic Sandy	0-137 137+
loam	

th (cm) Layer Depth (cm)

137 Fibric 0-91
- Hemic 91-122

Microrellef: 30 cm.

Location: 2000 feet N and 690 feet W of the SE corner of Sec. 9, T.66N., R.25W. (NE1/4 of SW1/4 of NE1/4 of SE1/4. Sec. 9, T.66N., R.25W.)

Vegetation: Black spruce crown cover of about 80 percent with scattered tamarack; understory consists of leatherleaf with some swamp laurel, bog rosemary, and grasses; ground cover consists of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Layer	Depth (cm)
Fibric	0 61
Hemic	61-228
Sapric	228-244
Very	244+
fine	
sand	

Location: 2550 feet S and 2550 feet E of the NW corner of Sec. 21, T.66N., R.25W. (SE1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 21, T.66N., R.25W.)

Vegetation: Black spruce crown cover of about 60 percent with about 25 percent crown cover of tamarack and scattered paper birch, black ash, and aspen; lush understory consists of Labrador tea with some speckled alder, grasses, and willow; ground cover consists of some Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Layer	Depth (cm)
Hemic	0-167
Medium	167+
sand	

(NW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 19, T.66N., R.25W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of

Location: 440 feet S and 1600 feet W of the NE corner of Sec. 9, T.66N., R.25W.

Vegetation: Scattered black spruce and tamarack; sparse understory consists of

some leatherleaf, swamp laurel, grasses, and sedges; ground cover consists

122+

(SE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 9, T.66N., R.25W.)

Described By: D. Mellem and P. Guertin on November 29, 1979.

Silty

clay

 egetation: Scattered black spruce and tamarack; sparse understory consists of leatherleaf with some sedges, swamp laurel, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some cranberry.

Location: 2300 feet S and 800 feet E of the NW corner of Sec. 29, T.66N., R.25W.

Microrellef: 30 cm.

Depth To Water Table: 30 cm.

mostly of Sphagnum mosses.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 11, 1979.

Layer	Depth (cm)
Hemic Coarse sand	0-274 274+

Location: 760 feet S and 1760 feet W of the NE corner of Sec. 31, T.66N., R.25W. (NW1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 31, T.66N., R.25W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, swamp laurel, and sedges; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: 183 cm.

Described By: D. Mellem and P. Guertin on November 29, 1979.

Laver

Hemic Coarse Location: 1650 feet S and 360 feet W of the NW corner of Sec. 33, T.66N., R.25W. (SE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 33, T.66N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of sedges with some swamp laurel and leatherleaf; ground cover consists mostly of false Solomon's seal.

Microrellef: 15 cm.

Depth To Water Table: 122 cm.

Described By: D. Mellem and P. Guertin on November 28, 1979.

	Depth (cm)	Layer	Depth (cm)
	0-183	Hemic	0-335
•	183+	Silt	335+
		loam	

Location: 2000 feet S and 625 feet W of the NE corner of Sec. 14, T.67N., R.25W. (NE1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 14, T.67N., R.25W.)

Vegetation: Scattered black spruce; sparse understory consists of some leatherleaf, swamp laurel, sedges, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 24 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 29, 1979.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-274
Clay	274+

Location: 1410 feet N and 1000 feet E of the SW corner of Sec. 17, T.67N., R.25W. (SE1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 17, T.67N., R.25W.)

Vegetation: Black spruce crown cover or about 75 percent with scattered tamarack, black ash, and paper birch; understory consists of speckled alder, leatherleaf, and bog rosemary with some grasses, Labrador tea, and swamp laurel; ground cover consists of Sphagnum mosses with some cranberry.

Microrelief: 25 cm.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Hemic 0-236 Sapric 236-244 Very 244+ fine sand Location: 100 feet N and 75 feet E of the SW corner of Sec. 30, T.68N., R.25W. (Gov. lot 4, Sec. 30, T.68N., R.25W.)

Vegetation: Scattered black spruce; understory consists of sedges with some leatherleaf, Labrador tea, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some snowberry and cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 28, 1977.

Layer	Depth (cm)
Fibric	0- 21
Hemic	21-295
Sand	295+

Location: 1350 feet S and 60 feet E of the NW corner of Sec. 31, T.68N., R.25W. (Gov. Lot 2, Sec. 31, T.68N., R.25W.)

Vegetation: Scattered black spruce; understory consists of leatherleaf with some sedges, bog rosemary, swamp laurel, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 28, 1977.

ayer_	Depth (cm)
ibric	0- 13
lemic	13-261
Silty	261+
lay	
oam	

Location: 2160 feet N and 50 feet E of the SW corner of Sec. 31, T.68N., R.25W. (Gov. Lot 3, Sec. 31, T.68N., R.25W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of some leatherleaf, sedge, Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 25 cm

Depth To Water Table: At surface.

Described By: D. Mellem on October 28, 1977.

 Layer
 Depth (cm)

 Fibric
 0- 12

 Hemic
 12-455

 Sand
 455+

Location: 775 feet N and 50 feet E of the SW corner of Sec. 31, T.68N., R.25W. (Gov. Lot 4, Sec. 31, T.68N., R.25W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of some sedges, leatherleaf, Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 28, 1977.

Layer	Depth (cm)
Fibric	0- 9
Hemic	9-480
Sand	480+

Location: 2160 feet S and 950 feet E of the NW corner of Sec. 32, T.68N., R.25W. (NW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 32, T.68N., R.25W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; understory consists of leatherleaf with some swamp laurel, bog rosemary, cotton grass, and grasses; ground cover consists of some Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-213
Silty	213+
clay	
loam	

Location: 2400 feet S and 2425 feet W of the NE corner of Sec. 2, T.69N., R.25W. (SW1/4 of SW1/4 of SW1/4 of NE1/4, Sec. 2, T.69N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of swamp laurel and cotton grass with some leatherleaf, bog rosemary, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, and cranberry.

Depth (cm)

0-213

213-477

477+

Microrelief: 35 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

Layer

Hemic

Sapric

Clay

Location: 1335 feet N and 900 feet E of the SW corner of Sec. 3, T.69N., R.25W. (SW1/4 of SE1/4 of NW1/4 of SW1/4. Sec. 3, T.69N., R.25W.)

Vegetation: Tamarack crown cover of about 70 percent with about 20 percent black spruce; lush understory consists mostly of Labrador tea with leatherleaf, bog rosemary, and grasses; ground cover consists of Sphagnum mosses with some feather mosses.

Microrellef: 20 cm.

Depth To Water Table: 10 cm.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-325
Clay	325+

Location: 1350 feet N and 1610 feet E of the SW corner of Sec. 4, T.69N., R.25W. (SE1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 4, T.69N., R.25W.)

Vegetation: Black spruce crown cover of about 30 percent with scattered tamarack; sparse understory consists of leatherleaf with some Labrador tea, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses and false Solomon's seal.

Microrellef: 50 cm.

Depth To Water Table: 10 cm.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-366
Silty	366+
clay	

Location: 2155 feet N and 1540 feet E of the SW corner of Sec. 10, T.69N., R.25W. (SW1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 10, T.69N., R.25W.)

Vegetation: Black spruce crown cover of about 75 percent with about 20 percent tamarack; understory consists of leatherleaf with bog rosemary and some Labrador tea and grasses; ground cover consists of Sphagnum mosses with some feather mosses.

Microrellet: 25 cm

Microrellef: 15 cm.

Depth To Water Table: At surface.

Depth To Water Table: Not recorded.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer	Depth (cm)
Hemic	0-335
Sapric	335-366
Clay	366+

Location: 625 feet S and 890 feet W of the NE corner of Sec. 13, T.69N., R.25W. (SE1/4 of NW1/4 of NE1/4 of NE1/4 of Sec. 13, T.69N., R.25W.)

Vegetation: Scattered black spruce: Jush understory consists mostly of grasses with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 15 cm.

Denth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on November 16, 1979.

Layer	Depth (cm)	
Fibric	0-547	
Hemic	547+	

Location: 1850 feet S and 315 feet E of the NE corner of Sec. 22, T.69N., R.25W.

Vegetation: Understory consists mostly of speckled alder with grasses, leatherleaf,

Microrellef: 10 cm

Depth To Water Table: At surface.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer	Depth (cm	
Hemic	0-274	
Sapric	274-305	
Silty	305+	
clav		

(SW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 22, T.69N., R.25W.)

and bog rosemary; ground cover consists of Sphagnum mosses.

Layer Depth (cm) Fibric 0-152 Hemic 152-411 Silty 411+ clay

Location: 2570 feet N and 2150 feet E of the SW corner of Sec. 14, T.69N., R.25W.

Vegetation: Scattered tamarack and black spruce: understory consists of sedges

cover consists mostly of Sphagnum mosses with some other mosses.

with some leatherleaf, swamp laurel, bog rosemary, and cotton grass; ground

(NW1/4 of NE1/4 of NE1/4 of SW1/4, Sec. 14, T.69N., R.25W.)

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

Location: 450 feet N and 1675 feet E of the SW corner of Sec. 23, T.69N., R.25W. (NE1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 23, T.69N., R.25W.)

Depth (cm)

0-152

152-285 285-305

305 +

Vegetation: Consists of black spruce, tamarack, Labrador tea, leatherleaf and bog birch; ground cover consists of Sphagnum mosses, and cranberry.

Microrelief: 30 cm.

Depth to Water Table: Not recorded.

Layer

Fibric

Hemic

Sapric

soil

Mineral

Described: On July 19, 1976.

Location: 2460 feet N and 2080 feet W of the SE corner of Sec. 23, T.69N., R.25W. (NE1/4 of NW1/4 of NW1/4 of SE1/4, Sec. 23, T.69N., R.25W.)

Vegetation: Consists of tamarack, Labrador tea, leatherleaf and bog birch; ground cover consists of Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described: On July 19, 1976.

Layer	Depth (cm
Hemic	0-305
Sapric	305+

Location: 50 feet S and 50 feet E of the NW corner of Sec. 13, T.69N., R.25W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 13, T.69N., R.25W.)

Vegetation: Scattered tamarack and black spruce: sparse understory consists of sedges with some bog birch, Labrador tea, leatherleaf, swamp laurel, and bog rosemary: ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrelief: 40 cm

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

ayer	Depth (cm)
emic	0-457
apric	457+

Location: 925 feet S and 940 feet W of the NE corner of Sec. 23, T.69N., R.25W. (NE1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 23, T.69N., R.25W.)

Vegetation: Consists of black spruce, tamarack, Labrador tea, bog rosemary, and leatherleaf.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described: On July 19, 1976.

Layer	Depth (cm)
Hemic Bottom	0-305+
unknown	

Location: 250 feet S and 2620 feet E of the NW corner of Sec. 24, T.69N., R.25W. (NE1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 24, T.69N., R.25W.)

Vegetation: Scattered tamarack and black spruce; lush understory consists mostly of bog birch with some grasses, Labrador tea, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer	Depth (cm)	
Hemic	0-315	
Clay	315+	

Location: 1800 feet S and 475 feet E of the NW corner of Sec. 26, T.69N., R.25W. (SE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 26, T.69N., R.25W.)

Vegetation: Consists of black spruce, tamarack, Labrador tea, bog rosemary, leatherleaf and swamp laurel; ground cover consists of Sphagnum mosses and cranberry.

Microrellef: 30 cm.

Depth To Water Table: 5 cm. Described: On July 19, 1976.

Hemic 0-305 Mineral 305+ soil . Location: 1560 feet S and 2550 feet E of the NW corner of Sec. 28, T.69N., R.25W. (NF1/4 of NE1/4 of SE1/4 of NW1/4, Sec. 28, T.69N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of sedges and leatherleaf with some Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, and cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

Layer	Depth (cm)	
Hemic	0-152	
Sapric	152-183	
Silty	183+	
clay		

Location: 145 feet S and 2075 feet W of the NE corner of Sec. 34, T.69N., R.25W. (NE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 34, T.69N., R.25W.)

Vegetation: Consists of black spruce, poplar, bog rosemary, leatherleaf, swamp laurel, bog birch, williow, and speckled alder; ground cover consists of Sphagnum mosses.

Microrellef: 45 cm.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Depth to Water Table: Not recorded.

Described: On July 19, 1976.

Layer	Depth (cm)	
Hemic	0- 76	
Sapric	76-188	
Clay	188+	
Sapric	76-188	

Location: 2475 feet S and 1200 feet W of the NE corner of Sec. 35, T.69N., R.25W. (SW1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 35, T.69N., R.25W.)

Vegetation: Tamarack and black spruce crown cover of about 95 percent; lush understory consists of leatherleaf and bog rosemary with some Labrador tea; ground cover consists of Sphagnum mosses with some feather mosses.

Microrellef: 40 cm.

Depth To Water Table: Near surface.

Described By: D. Olson and R. Graber on November 16, 1979.

.ayer	Depth (cm
Hemic	0-213
Clay	213+

Location: 1425 feet S and 2125 feet E of the NW corner of Sec. 34, T.151N., R.25W. (NW1/4 of NE1/4 of SE1/4 of NW1/4, Sec. 34, T.151N., R.25W.)

Vegetation: Northern white cedar crown cover of about 40 percent with scattered black spruce; understory consists of bog birch and grasses with some willow, Labrador tea, leatherleaf, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry, other mosses, and aster.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and N. Aaseng on December 10, 1979.

Layer	Depth (cn
Hemic Bottom	0-547+
unknown	

Location: 2450 feet N and 1865 feet W of the SE corner of Sec. 31, T.69N., R.25W. (NW1/4 of NE1/4 of NW1/4 of SE1/4, Sec. 31, T.69N., R.25W.)

Vegetation: Scattered black spruce; lush understory consists of Labrador tea and leatherleaf with some swamp laurel and grasses.

Microrelief: 15 cm.

Depth To Water Table: 15 cm.

Described By: D. Mellem and D. Haverkost on November 16, 1979.

Layer	Depth (cm)	
Fibric	0- 91	
Hemic	91-183	
Silty	183+	
clay		

Location: 655 feet N and 150 feet W of the SE corner of Sec. 27, T.70N., R.25W. (SE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 27, T.70N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists mostly of leatherleaf and swamp laurel with some cotton grass and grasses; ground cover consists of Sphagnum mosses with some feather mosses.

Microrelief: Not recorded.

Depth To Water Table: Below 235 cm.

Described By: D. Olson and R. Graber on November 16, 1979.

_ayer	Depth (cm)
ibric	0- 61
Hemic	61- 91
Sapric	91-235
Clav	235+

Location: 940 feet S and 875 feet E of the NW corner of Sec. 3, T.152N., R.25W. (Gov. Lot 4, Sec. 3, T.152N., R.25W.)

Vegetation: Consists mostly of grasses with some tamarack, black spruce, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: B. Sether and L. Severson on April 7, 1980.

Layer	Depth (cm)
Hemic	0-198
Sapric	198-244
Sandy	244+
clay	
loam	

Fibric 0-183 Hemic 183-335 Limnic 335-366 + Bottom unknown

Depth (cm)

Location: 100 feet N and 2500 feet W of the SE corner of Sec. 27, T.151N., R.25W.

Vegetation: Black spruce grown cover of about 60 percent; understory consists of

(SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 27, T.151N., R.25W.)

Labrador tea; ground cover consists mostly of mosses.

Layer

Described By: D. Mellem and D. Haverkost on December 10, 1979.

Location: 950 feet S and 900 feet E of the NW corner of Sec. 33, T.152N., R.25W. (NW1/4 of SE1/4 of NW1/4 of NW1/4, Sec. 33, T.152N., R.25W.)

Vegetation: Consists of grasses, speckled alder, and sedges with some dogwood, cattail, horsetail, and aster.

Microrellef: 5 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and N. Aaseng on December 10, 1979.

Layer Depth (cm)
Sapric 0-183
Medium 183+
to
fine
sand

Location: 2100 feet N and 1600 feet W of the SE corner of Sec. 34, T.152N., R.25W. (SE1/4 of NE1/4 of NW1/4 of SE1/4, Sec. 34, T.152N., R.25W.)

Vegetation: Scattered black spruce; understory consists mostly of grasses with willow.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on December 10, 1979.

_ayer	Depth (cm
lemic	0-91
Silt	91+
and	
sand	

Location: 625 feet S and 600 feet W of the NE corner of Sec. 18, T.153N., R.25W. (SW1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 18, T.153N., R.25W.)

Vegetation: Lush understory consists of sedges and grasses with some black spruce and leatherleaf; ground cover consists of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 30 cm.

Depth To Water Table: 10 cm.

Described By: B. Sether and S. Nelson on April 3, 1980.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-259
Sandy	259+
loam	

Location: 2000 feet S and 275 feet W of the NE corner of Sec. 19, T.153N., R.25W. (SE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 19, T.153N., R.25W.)

Vegetation: Tamarack crown cover of about 40 percent with scattered black spruce; lush understory consists mostly of sedges with some leatherleaf, swamp laurel, bog rosemary, and cotton grass; ground cover consists of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: 10 cm. above surface.

Described By: H. Mooers and S. Nelson on April 7, 1980.

Layer	Depth (cm)
Hemic Silt Ioam	0-244 244+

Location: 2175 feet S and 2475 feet E of the NW corner of Sec. 21, T.153N., R.25W. (NE1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 21, T.153N., R.25W.)

Vegetation: Lush understory consists mostly of grasses with some black spruce. leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some ouner mosses, cranberry, and pitcher plant.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and L. Severson on April 7, 1980.

Layer	Depth (cm)
Hemic	0-229
Medium	229+
to	
coarse	
cand	

Location: 810 feet N and 1425 feet E of the SW corner of Sec. 22, T.153N., R.25W. (SW1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 22, T.153N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of grasses with some leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some feather mosses, snowberry, and false Solomon's seal.

Microrelief: 50 cm.

Depth To Water Table: At surface.

Described By: B. Sether and L. Severson on April 7, 1980.

Layer	Depth (cm)
Hemic	0-183
Fibric	183-213
Medium	213+
sandy	
loam	

Location: 675 feet N and 425 feet E of the SW corner of Sec. 5, T.153N., R.25W. (SE1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 5, T.153N., R.25W.)

Vegetation: Black spruce crown cover of about 85 percent; sparse understory consists of some Labrador tea; ground cover consists mostly of Sphagnum mosses with some cranberry and snowberry.

Microrelief: 46 cm.

Depth To Water Table: 30 cm.

Described By: D. Mellem and L. Severson on April 7, 1980.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-152
Sapric	152-163
Silt	163+
loam	
with	
pebbles	

Location: 310 feet N and 300 feet E of the SW corner of Sec. 20, T.153N., R.25W. (NW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 20, T.153N., R.25W.)

Vegetation: Lush understory consists mostly of grasses with some black spruce, tamarack, alder, leatherleaf, bog rosemary, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and L. Severson on April 7, 1980.

Layer	Depth (cm)
Fibric	0-305
Hemic	305-335
Clay	335+
loam	

Location: 800 feet N and 110 feet W of the SE corner of Sec. 24, T.153N., R.25W. (SE1/4 of NE1/4 of SE1/4 of SE1/4. Sec. 24, T.153N., R.25W.)

Vegetation: Black spruce crown cover of about 60 percent; sparse understory consists of some Labardor tea, leatherleaf, swamp laurel, grasses, and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and S. Nelson on April 8, 1980.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-168
Loamy sand	168+

Location: 1150 feet N and 2620 feet E of the SW corner of Sec. 25, T.153N., R.25W. (NE1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 25, T.153N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of Labrador tea with some sedges, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, snowberry, and false Solomon's seal.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: B. Sether and D. Haverkost on April 8, 1980.

 Layer
 Depth (cm)

 Hemic
 0- 91

 Sapric
 91-122

 Silt
 122+

 loam
 122+

Location: 1090 feet S and 475 feet W of the NE corner of Sec. 29, T.153N., R.25W.

Vegetation: Scattered tamarack and black spruce; understory consists of

leatherleaf with some sedges, Labrador tea, swamp laurel, and bog rosemary;

ground cover consists of Sphagnum mosses with some feather mosses, pitcher

Depth (cm)

0- 61

61-320

320+

(SW1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 29, T.153N., R.25W.)

plant, cranberry, and false Solomon's seal.

Described By: H. Mooers and S. Nelson on April 7, 1980.

Laver

Fibric

Hemic

Silty

clav

Depth To Water Table: Not recorded.

Location: 2550 feet N and 825 feet E of the SW corner of Sec. 27, T.153N., R.25W. (NW1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 27, T.153N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of sedges with some leatherleaf. Labrador tea, swamp laurel, and grasses: ground cover consists mostly of Sphagnum mosses with some feather mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and S. Nelson on April 7, 1980.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-335
Sapric	335-381
Silty	381+
clay	
loam	

Location: 1825 feet S and 1860 feet W of the NE corner of Sec. 30, T.153N., R.25W. (SW1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 30, T.153N., R.25W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of sedges with some Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some feather mosses.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and S. Nelson on April 7, 1980.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-366
Sapric	366-381
Silty	381+
clay	
loam	

Location: 490 feet N and 175 feet W of the SE corner of Sec. 30, T.153N., R.25W.
 (NE1/4 of SE1/4 of SE1/4, Sec. 30, T.153N., R.25W.)
 Vegetation: Understory consists of grasses with some black spruce, bog birch,

Location: 1375 feet N and 2425 feet E of the SW corner of Sec. 28, T.153N., R.25W.

Vegetation: Lush understory consists of sedges and grasses with some black

spruce, tamarack, leatherleaf, and swamp laurel; ground cover consists mostly

Depth (cm)

0-198

198+

of Sphagnum mosses with some feather mosses and false Solomon's seal.

(SE1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 28, T.153N., R.25W.)

Described By: B. Sether and L. Severson on April 7, 1980.

Layer

Hemic

Sandv

clav

Vegetation: Understory consists of grasses with some black spruce, bog birch, leatherleaf, and bog rosemary; ground cover consists of some Sphagnum and feather mosses.

Microrellef: 15 cm.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: B. Sether and S. Nelson on April 3, 1980.

Layer	Depth (cm)
Hemic Sandy clay	0-244 244+

Location: 2200 feet S and 150 feet W of the NE corner of Sec. 31, T.153N., R.25W. (NE1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 31, T.153N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of some leatherleaf, sedges, grasses, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some feather mosses, cranberry, and pitcher plant.

Microrellef: 30 cm.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on April 3, 1980.

Layer Depth (cm)

Hemic 0-244

Sapric 244-274

Loamy 274+
sand

Location: 825 feet S and 160 feet E of the NW corner of Sec. 31, T.153N., R.25W. (Gov. Lot 1, Sec. 31, T.153N., R.25W.)

Vegetation: Scattered tamarack and black spruce; sparse understory consists of some leatherleaf, sedges, Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some feather mosses, snowberry, and pitcher plant.

Microrellef: 30 cm.

Depth To Water Table: 6 cm.

Described By: H. Mooers and D. Haverkost on April 3, 1980.

Layer	Depth (cm)
Hemic	0-335
Silty	335+
clay	
loam	

Location: 1725 feet S and 1610 feet W of the NE corner of Sec. 32, T.153N., R.25W. (SE1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 32, T.153N., R.25W.)

Vegetation: Scattered tamarack and black spruce; understory consists of sedges with some speckled alder, leatherleaf, Labrador tea, swamp laurel, bog rosemary, and grasses; ground cover consists mostly of Sphagnum mosses with some feather mosses, false Solomon's seal, and pitcher plant.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and S. Nelson on April 7, 1980.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-274
Loamy	274+
sand	

Location: 750 feet S and 90 feet E of the NW corner of Sec. 35, T.153N., R.25W. (NW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 35, T.153N., R.25W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; understory consists of some speckled alder, bog birch, Labrador tea, leatherleaf, swamp laurel, bog rosemary, sedges, grasses, and ferns; ground cover consists mostly of Sphagnum mosses with some feather mosses, snowberry, and pitcher plant.

Microrellef: 30 cm.

Depth To Water Table: 8 cm.

Described By: H. Mooers and S. Nelson on April 7, 1980.

Laver Depth (cm) Hemic 0-259 Loamy 259+ sand with pebbles

Location: 1660 feet N and 400 feet E of the SW corner of Sec. 7, T.154N., R.25W. (Gov. Lot 3, Sec. 7, T.154N., R.25W.)

Vegetation: Scattered black spruce, aspen, and tamarack; lush understory consists mostly of grasses with some bog birch, cattails, Labrador tea, leatherleaf, bog rosemary, blackberry, and ferns; ground cover consists mostly of Sphagnum mosses with some snowberry.

Microrellef: 23 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer	Depth (cm
Hemic	0-76
Silt	76+
oam	

Location: 125 feet N and 2510 feet W of the SE corner of Sec. 10, T.154N., R.25W. (SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 10, T.154N., R.25W.)

Vegetation: Consists of tamarack with bog birch, willow and speckled alder; ground cover consists of Sphagnum mosses.

Microrellef: Not recorded.

Vegetation: Not recorded.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 16, 1976.

Depth To Water Table: Not recorded. Described By: M. Domeier on July 16, 1976.

Layer	Depth (cm)
Hemic Sandy clay	0-122 122+

Location: 1160 feet S and 250 feet E of the NW corner of Sec. 13, T.154N., R.25W. (SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 13, T.154N., R.25W.)

Vegetation: Tamarack crown cover of about 65 percent with about 25 percent black spruce; understory consists of leatherleaf with some swamp laurel, grasses, and speckled alder; ground cover consists of Sphagnum mosses. Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)
Hemic	0-183
Sapric	183-198
Medium	198+
to	
fine	
sand	

Location: 175 feet S and 2125 feet W of the NE corner of Sec. 15, T.154N., R.25W.

(NE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 15, T.154N., R.25W.) Vegetation: Crown cover consists of black spruce, tamarack, and northern white cedar; understory consists of Labrador tea, bog rosemary, and leatherleaf.

Microrelief: Not recorded.

Depth To Water Table: Not recorded. Described By: M. Domeier on July 16, 1976.

Layer Depth (cm) Hemic 0-91 Sandy 91 +loam

Location: 500 feet S and 1790 feet W of the NE corner of Sec. 15, T.154N., R.25W.

(SW1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 15, T.154N., R.25W.)

Laver Depth (cm) Hemic 0-122 Loamy 122+ sand

Location: 660 feet N and 490 feet E of the SW corner of Sec. 10, T.154N., B.25W. (SE1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 10, T.154N., R.25W.)

Vegetation: Tamarack crown cover of about 65 percent with about 20 percent crown cover of black spruce and scattered paper birch; understory consists of leatherleaf and bog rosemary with some speckled alder, Labrador tea, and grasses; ground cover consists of some Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 11, 1979.

Layer	Depth (cm)
lemic	0-167
Sapric	167-183
ine	183+
and	

Location: 1140 feet S and 1110 feet W of the NE corner of Sec. 15, T.154N., R.25W. (SW1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 15, T.154N., R.25W.)

Vegetation: Not recorded. Microrellef: Not recorded. Depth To Water Table: Not recorded.

Described By: M. Domeier on July 16, 1976.

Layer	Depth (cm)
Hemic Sandy loam	0-122 122+

Location: 800 feet S and 1440 feet W of the NE corner of Sec. 15, T.154N., R.25W. (NE1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 15, T.154N., R.25W.)

Vegetation: Not recorded. Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 16, 1976.

Depth (cm)
0-122 122+

Location: 1860 feet S and 335 feet W of the NE corner of Sec. 15, T.154N., R.25W. (SW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 15, T.154N., R.25W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 16, 1976.

Layer Depth (cm)

Peat 0-122
Clay 122+
loam

Location: 1490 feet S and 710 feet W of the NE corner of Sec. 15, T.154N., R.25W. (NE1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 15, T.154N., R.25W.)

Vegetation: Not recorded.
Microrelief: 60 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 16, 1976.

Layer	Depth (cm)
Hemic	0-122
Sand	122+

Location: 2200 feet S and 25 feet W of the NE corner of Sec. 15, T.154N., R.25W. (NE1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 15, T.154N., R.25W.)

Vegetation: Crown cover consists of black spruce and tamarack; understory consists of Labrador tea, bog rosemary, bog birch, willow, and speckled alder.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 16, 1976.

_ayer	Depth (cm)
Peat	0-122
Sandy	122+
lay	
oam	

Location: 1775 feet S and 250 feet E of the NW corner of Sec. 15, T.154N., R.25W. (SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 15, T.154N., R.25W.)

Vegetation: Consists of balsam fir, poplar and ferns.

Microrellef: Negligible.

Depth To Water Table: Below 69 cm.

Described By: M. Domeier on September 20, 1976.

Layer	Depth (cm)
Hemic	0-58
Sapric	58-69
Mineral	69+
soil	

Location: 2550 feet S and 1825 feet W of the NE corner of Sec. 20, T.154N., R.25W. (SW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 20, T.154N., R.25W.)

Vegetation: Old clearcut area surrounded by northern white cedar and black spruce of over 20 feet; understory consists of grasses; ground cover consists of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer	Depth (cm)
Hemic	0-183
Fibric	183-305
Clay	305+
loam	

Location: 1300 feet S and 75 feet W of the NE corner of Sec. 21, T.154N., R.25W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 21, T.154N., R.25W.)

Vegetation: Stunted tamarack and northern white cedar crown cover of about 35 percent each with scattered black spruce; understory consists of grasses with some bog rosemary, leatherleaf, and Labrador tea; ground cover consists of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)
Fibric Hemic Medium to coarse sand with pebbles	0-122 122-213 213+

Location: 950 feet N and 1625 feet E of the SW corner of Sec. 24, T.154N., R.25W. (NE1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 24, T.154N., R.25W.)

Vegetation: Scattered tamarack, black spruce, and paper birch; understory consists of bog birch and leatherleaf with some bog rosemary and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 35 cm.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Depth (cm)
0- 91
91-244
244+

Location: 475 feet S and 245 feet E of the NW corner of Sec. 27, T.154N., R.25W. (SW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 27, T.154N., R.25W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists of sedges with some leatherleaf, bog rosemary, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer	Depth (cm)
Fibric	0-274
Hemic	274-517
Silty	517+
clay	

Location: 100 feet S and 2000 feet E of the NW corner of Sec. 1, T.155N., R.25W. (Gov. Lot 3, Sec. 1, T.155N., R.25W.)

Vegetation: Scattered black spruce; sparse understory consists of Labrador tea with some leatherleaf, swamp laurel, bog rosemary, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 15 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)
Hemic	0-167
Sapric	167-183
Fine	183+
sand	

Location: 1470 feet S and 240 feet W of the NE corner of Sec. 13, T.155N., R.25W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 13, T.155N., R.25W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered northern white cedar, balsam and paper birch; understory consists of bog birch with some Labrador tea, grasses, and blackberry; ground cover consists mostly of Sphaanum mosses.

Microrellef: 30 cm.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer Depth (cm)
Hemic 0-122
Silty 122+
clay

Location: 1050 feet N and 1670 feet W of the SE corner of Sec. 6, T.156N., R.25W.

bog birch, grasses, and forbs with some speckled alder, Labrador tea, and

Depth (cm)

0-91

91 +

Location: 1990 feet S and 2255 feet E of the NW corner of Sec. 31, T.156N., R.25W.

Vegetation: Old clear cut area surrounded by black spruce about 20 feet tall; lush

understory consists of Labrador tea and leatherleaf with some bog birch and

grasses; ground cover consists mostly of Sphagnum mosses with snowberry

(SW1/4 of NE1/4 of SE1/4 of NW1/4, Sec. 31, T.156N., R.25W.)

(NE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 6, T.156N., R.25W.)

Vegetation: Scattered black spruce and paper birch; understory consists mostly of

ferns: ground cover consists mostly of Sphagnum mosses.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Laver

Hemic Sandy

clav

Location: 300 feet S and 1710 feet E of the NW corner of Sec. 4, T.156N., R.25W. (NE1/4 of NW1/4 of NE1/4 of NW1/4. Sec. 4, T.156N., R.25W.)

Vegetation: Black spruce crown cover of about 45 percent; understory consists of Labrador tea; ground cover consists mostly of mosses with some Sphagnum mosses and snowberry.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)
Hemic	0-152
Medium	152+
to	
fine	
sand	

Location: 250 feet N and 1650 feet E of the SW corner of Sec. 24, T.156N., R.25W. (SE1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 24, T.156N., R.25W.)

Vegetation: Scattered black spruce; sparse understory consists of Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses and snowberry.

Microrellef: 15 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)
Hemic	0-137
Sandy	137+
clay	

Location: 1775 feet N and 375 feet W of the SE corner of Sec. 32, T.156N., R.25W. (NW1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 32, T.156N., R.25W.)

Vegetation: Scattered black spruce and paper birch; sparse understory consists of some speckled alder, grasses, Labrador tea, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 30 cm.

Depth To Water Table: 15 cm. above surface.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)
Hemic	0-122
Medium	122+
cond	

Location: 150 feet N and 575 feet E of the SW corner of Sec. 4, T.156N., R.25W. (SE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 4, T.156N., R.25W.)

Vegetation: Black spruce crown cover of about 85 percent; understory consists of sedges with some leatherleaf and Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrellef: 6 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 30, 1979.

.ayer	Depth (cm)
lemic	0-122
Sapric	122-152
Clay	152+
nam	

Location: 1150 feet N and 1960 feet W of the SE corner of Sec. 28, T.156N., R.25W. (NE1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 28, T.166N., R.25W.)

Vegetation: Scattered black spruce, tamarack, and paper birch; lush understory consists mostly of grasses and bog birch with some speckled alder, cattall, and blackberry; ground cover consists of some Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer	Depth (cm)
Hemic Silty	0-122 122+
clay	

Location: 1150 feet S and 2050 feet W of the NE corner of Sec. 35, T.156N., R.25W. (SE1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 35, T.156N., R.25W.)

Vegetation: Scattered black spruce, tamarack, and paper birch; lush understory consists of Labrador tea, bog birch, and leatherleaf with some speckled alder, swamp laurel, bog rosemary, and sedges; ground cover consists mostly of Sohagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer	Depth (cm)
Hemic	0-106
SIn	106+
and	
sand	

Depth To Water Table: 30 cm. Described By: D. Mellem and F

Microrellef: 30 cm.

and some other mosses.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer Depth (cm)
Hemic 0-61
Silty 61+
loam

Location: 750 feet S and 1415 feet W of the NE corner of Sec. 2, T.157N., R.25W. (Gov. Lot 2, Sec. 2, T.157N., R.25W.)

Vegetation: Black spruce crown cover of about 95 percent; understory consists of Labrador tea with some leatherleaf, swamp laurel, bog rosemary, and grasses; ground cover consists mostly of Sphagnum mosses with some cranberry and faise Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: 31 cm.

mosses and cranberry.

Depth To Water Table: At surface.

Microrellef: 20 cm.

Described By: B. Wakanabo on August 24, 1977.

Layer	Depth (cm)
Peat	0-232
Silt	232+
and	
very	
fine	
sand	

Location: 1450 feet N and 1000 feet E of the SW corner of Sec. 15, T.157N., R.25W.

Vegetation: Scattered black spruce and tamarack; understory consists of

leatherleaf and sedges with some swamp laurel, bog rosemary, and cotton

grass; ground cover consists mostly of Sphagnum mosses with some other

Depth (cm)

0-213

213+

(SW1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 15, T.157N., R.25W.)

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer

Hemic

Clay

Location: 650 feet S and 2575 feet E of the NW corner of Sec. 2, T.157N., R.25W. (Gov. Lot 3, Sec. 2, T.157N., R.25W.)

Vegetation: Black spruce crown cover of about 70 percent with a tamarack crown cover of about 25 percent; lush understory consists mostly of leatherleaf and bog birch with some Labrador tea, willow, swamp laurel, grasses, and bog rosemary; ground cover consists of Sphagnum mosses with some cranberry.

Microrelief: 42 cm

Depth To Water Table: 23 cm.

Described By: B. Wakanabo on August 24, 1977.

Layer	Depth (cm)
Hemic Silty clay	0-210 210+

Location: 1050 feet S and 1530 feet W of the NE corner of Sec. 16, T.157N., R.25W. (SE1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 16, T.157N., R.25W.)

Vegetation: Black spruce crown cover of about 40 percent; understory consists of Labrador tea, leatherleaf, and sedges with some swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrellef: 40 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer	Depth (cm)
Hemic	0-177
Clay	177+

Location: 1865 feet N and 2500 feet E of the SW corner of Sec. 21, T.157N., R.25W.

(NE1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 21, T.157N., R.25W.)

Location: 2150 feet S and 1800 feet W of the NE corner of Sec. 15, T.157N., R.25W.

Vegetation: Scattered black spruce and tamarack; sparse understory consists of

leatherleaf, swamp laurel, bog rosemary, sedges, and grasses; ground cover

consists mostly of Sphagnum mosses with some other mosses and cranberry.

Depth (cm)

0-228

228-305

305-320

320+

(NW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 15, T.157N., R.25W.)

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer

Fibric

Hemic

Sapric

Silty

clay

Vegetation: Scattered black spruce and tamarack; sparse understory consists of Labrador tea, leatherleaf, swamp laurel, bog rosemary and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer	Depth (cr
Fibric	0-122
Hemic	122-274
Sapric	274-335
Silty	335+
clay	

Location: 665 feet N and 2525 feet E of the SW corner of Sec. 22, T.157N., R.25W. (SE1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 22, T.157N., R.25W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack; sparse understory consists of Labrador tea with some leatherleaf, swamp laurel, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and lingen berry.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer	Depth (cm)
Fibric	0- 45
Hemic	45-122
Clay	122+

Location: 800 feet S and 1150 feet W of the NE corner of Sec. 28, T.157N., R.25W. (NW1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 28, T.157N., R.25W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, bog rosemary, and sedges: ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-274
Very	274+
fine	
sand	

Location: 50 feet S and 100 feet W of the NE corner of Sec. 32, T.157N., R.25W. (NE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 32, T.157N., R.25W.)

Vegetation: Black spruce crown cover of about 30 percent with scattered tamarack; understory consists of Labrador tea with some leatherleaf, swamp laurel, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-305 305+

Location: 500 feet N and 550 feet W of the SE corner of Sec. 36, T.157N., R.25W. (NW1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 36, T.157N., R.25W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists of Labrador tea, leatherleaf, and swamp laurel with some sedges; ground cover consists mostly of Sphagnum mosses.

Microrelief: 61 cm.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Depth-To Water Table: 45 cm.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-167
Silt	167+
and	
sand	

Location: 675 feet S and 1375 feet E of the NW corner of Sec. 4, T.158N., R.25W. (Gov. Lot 3, Sec. 4, T.158N., R.25W.)

Vegetation: Black spruce crown cover of about 90 percent; sparse understory consists of Labrador tea with some sedges; ground cover consists mostly of Sphagnum and other mosses with some lingen berry and cranberry.

Microrellef: 40 cm.

Depth To Water Table: Below 198 cm.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-198 198+

Location: 2280 feet N and 1775 feet W of the SE corner of Sec. 16, T.158N., R.25W. (SW1/4 of NE1/4 of NW1/4 of SE1/4, Sec. 16, T.158N., R.25W.)

Vegetation: Scattered black spruce and tamarack; understory consists of sedges with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry and pitcher plant.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Layer Depth (cm) 0-238 Hemic Clay 238 +

Location: 25 feet S and 1425 feet W of the NE corner of Sec. 16, T.158N., R.25W.

Vegetation: Scattered black spruce; sparse understory consists of cotton grass

mostly of Sphagnum mosses with some other mosses and cranberry.

with some Labrador tea, leatherleaf, and swamp laurel; ground cover consists

(NE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 16, T.158N., R.25W.)

Described By: D. Olson and N. Aaseng on November 15, 1979.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer	Depth (cm)
Hemic Fine	0-259 259+
sand	

(NE1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 17, T.158N., R.25W.)

Vegetation: Black spruce crown cover of about 40 percent with about 40 percent crown cover of tamarack; lush understory consists mostly of leatherleaf. swamp laurel, bog rosemary, and grasses with some horsetall; ground cover consists of some Sphagnum and feather mosses.

Location: 925 feet S and 825 feet W of the NE corner of Sec. 17, T.158N., R.25W.

Location: 15 feet S and 2225 feet E of the NW corner of Sec. 15, T.158N., R.25W.

Vegetation: Tamarack crown cover of about 60 percent with scattered black

spruce; lush understory consists of Labrador tea, leatherleaf, swamp laurel,

and bog rosemary with some grasses; ground cover consists of Sphagnum and

Depth (cm)

0- 91 91-183

183-213

213 +

(NW1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 15, T.158N., R.25W.)

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer Fibric

Hemic Sapric

Fine

sand

Microrellef: 40 cm.

feather mosses.

Depth To Water Table: Near surface.

Microrellet: 50 cm

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-305 305+

Location: 950 feet N and 1735 feet E of the SW corner of Sec. 19, T.158N., R.25W. (SE1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 19, T.158N., R.25W.)

Vegetation: Scattered black spruce, tamarack, and paper birch: sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, sedges, and grasses; ground cover consists mostly of feather mosses with some Sphagnum mosses and snowberry.

Microrellef: 45 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Clay

loam

Layer Depth (cm) Hemic 0-106 106+

Vegetation: Scattered black spruce and tamarack; sparse understory consists of sedges with some leatherleaf, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Location: 1825 feet N and 500 feet E of the SW corner of Sec. 20, T.158N., R.25W.

(NE1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 20, T.158N., R.25W.)

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm)
Hemic Silty	0-381 381+
clay	

Location: 950 feet N and 2135 feet W of the SE corner of Sec. 20, T.158N., R.25W. (SE1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 20, T.158N., R.25W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of grasses, bog rosemary, and leatherleaf with some speckled alder and sedges; ground cover consists mostly of feather mosses with some Sphagnum mosses.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Hemic	0- 91
Fibric	91-183
Hemic .	183-416
Silty	416+
clay	

Location: 1100 feet N and 1640 feet W of the SE corner of Sec. 21, T.158N., R.25W. (NE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 21, T.158N., R.25W.)

Vegetation: Scattered black spruce and tamarack: understory consists of sedges with some bog rosemary, swamp laurel, and leatherleaf; ground cover consists of some cranberry.

Microrellef: 25 cm.

Microrellet: 20 cm

Depth To Water Table: At surface.

Described By: B. Wakanabo on August 25, 1977.

Layer

Fibric Hemic

Silty

clay

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng of November 15, 1979.

Layer	Depth (cm
Fibric-	0- 91
Hemic	91-264
Clay	264+

Location: 2600 feet S and 855 feet W of the NE corner of Sec. 25, T.158N., R.25W.

Vegetation: Understory consists mostly of grasses with some leatherleaf and bog

Depth (cm)

0- 65

65-206

206+

(SE1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 25, T.158N., R.25W.)

rosemary; ground cover consists of Sphagnum mosses.

Location: 2175 feet S and 1990 feet E of the NW corner of Sec. 22, T.158N., R.25W. (NW1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 22, T.158N., R.25W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists of grasses, leatherleaf, and swamp laurel; ground cover consists of Sphagnum mosses with some pitcher plant.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-305
Silty	305+
clay	

Location: 2500 feet S and 475 feet W of the NE corner of Sec. 25, T.158N., R.25W. (SW1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 25, T.158N., R.25W.)

Vegetation: Black spruce crown cover of about 90 percent; lush understory consists of grasses, leatherleaf, and Labrador tea with some swamp laurel and bog rosemary; ground cover consists of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellef: 35 cm.

Layer	Depth (cm)
Fibric	0- 59
Hemic	59-222
Silty	222+
clay	

Location: 975 feet S and 2550 feet W of the NE corner of Sec. 23, T.158N., R.25W. Vegetation: Scattered black spruce and tamarack; lush understory consists mostly

(NW1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 23, T.158N., R.25W.)

of Labrador tea with some leatherleaf and swamp laurel; ground cover consists mostly of mosses with some lingen berry.

Microrelief: 25 cm.

Depth To Water Table: Not recorded

Described By: D. Olson and N. Aaseng on October 31, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-122
Clay	122+

Depth To Water Table: 10 cm.

Described By: B. Wakanabo on August 25, 1977.

Layer	Depth (cm)
Fibric	0- 59
Hemic	59-222
Silty	222+
clay	

Vegetation: Scattered black spruce and tamarack; understory consists mostly of Labrador tea; ground cover consists mostly of mosses with some lingen berry. Microrellef: 25 cm.

(NE1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 27, T.158N., R.25W.)

Location: 1245 feet N and 1875 feet E of the SW corner of Sec. 27, T.158N., R.25W.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on October 31, 1979.

Layer	Depth (cm)
Fibric	0- 76
Hemic	76-157
Clay	157+
loam	

Location: 650 feet S and 625 feet W of the NE corner of Sec. 36, T.158N., R.25W. (SW1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 36, T.158N., R.25W.)

Vegetation: Black spruce grown cover of about 80 percent with a tamarack grown cover of about 25 percent; lush understory consists of Labrador tea and leatherleaf with some bog rosemary, grasses, and swamp laurel: ground cover consists of Sphagnum mosses with some cranberry.

Microrelief: 28 cm.

Depth To Water Table: 20 cm.

Described By: B. Wakanabo on August 24, 1977.

Location: 225 feet N and 610 feet E of the SW corner of Sec. 36, T.158N., R.25W. (SE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 36, T.158N., R.25W.)

Vegetation: Black spruce crown cover of about 60 percent with a tamarack crown cover of about 35 percent; lush understory consists of bog birch, leatherleaf, and grasses with some bog rosemary and swamp laurel; ground cover consists of some Sphagnum mosses and false Solomon's seal.

Microrellef: 25 cm.

Depth To Water Table: 20 cm.

Described By: B. Wakanabo on August 24, 1977.

Layer	Depth (cm)
Hemic	0-223
Silt	223+

Location: 575 feet N and 1950 feet E of the SW corner of Sec. 36, T.158N., R.25W. (NE1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 36, T.158N., R.25W.)

Vegetation: Black spruce crown cover of about 80 percent; lush understory consists of Labrador tea and leatherleaf with some swamp laurel, bog rosemary, and sedges; ground cover consists of false Solomon's seal with some Sphagnum mosses.

Microrelief: 10 cm.

Depth To Water Table: 30 cm.

Described By: B. Wakanabo on August 24, 1977.

Layer	Depth (cm)
Fibric	0-41
Hemic	41-90
Silt	90+
loam	

Layer Depth (cm) 0- 24 Hemic Fibric 24- 42 Hemic 42-287 Silty 287+

clay

plant.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Location: 575 feet N and 2125 feet W of the SE corner of Sec. 36, T.158N., R.25W. (NE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 36, T.158N., R.25W.)

Vegetation: Black spruce crown cover of about 40 percent; lush understory consists of leatherleaf and sedges with some Labrador tea, bog rosemary, and swamp laurel; ground cover consists of Sphagnum mosses with some false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: 10 cm.

Described By: B. Wakanabo on August 24, 1977.

Location: 2290 feet N and 1510 feet W of the SE corner of Sec. 19, T.159N., R.25W.

Vegetation: Scattered tamarack; understory consists of bog birch and swamp

laurel with some leatherleaf. Phragmites, fern, buck bean and horsetail; ground

cover consists mostly of Sphagnum mosses with some cranberry and pitcher

Depth (cm

0- 76

76-350

350 +

(SE1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 19, T.159N., R.25W.)

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer

Fibric

Hemic

Clay

Location: 600 feet N and 725 feet W of the SE corner of Sec. 36, T.158N., R.25W. (NE1/4 of SW1/4 of SE1/4 of SE1/4. Sec. 36, T.158N., R.25W.)

Vegetation: Scattered black spruce; understory consists of some sedges, Labrador tea, and bog rosemary; ground cover consists of some false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: 5 cm.

Described By: D. Olson on August 23, 1977.

Layer	Depth (cm)
Fibric	0- 24
Hemic	24-352
Silty	352+
clay	

Location: 2425 feet S and 2100 feet E of the NW corner of Sec. 22, T.159N., R.25W. (SW1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 22, T.159N., R.25W.)

Vegetation: Scattered tamarack and black spruce; sparse understory consists of some leatherleaf, Labrador tea, swamp laurel, sedges, and cotton grass; ground cover consists of some cranberry and false Solomon's seal.

Microrellef: 15 cm.

Depth To Water Table: 10 cm.

Described By: D. Olson and N. Aaseng on November 15, 1979.

91
21

Location: 565 feet S and 2225 feet W of the NE corner of Sec. 28, T.159N., R.25W. (SE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 28, T.159N., R.25W.)

Location: 560 feet S and 1275 feet E of the NW corner of Sec. 19, T.159N., R.25W.

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists of bog rosemary, grasses, cattails, speckled alder, and reeds; ground

cover consists mostly of Sphagnum and feather mosses with some pitcher

Depth (cm)

0-167

167-244

244-305

305-411

411+

(SE1/4 of NE1/4 of NW1/4 of NW1/4, Sec. 19, T.159N., R.25W.)

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer

Fibric

Hemic

Fibric

Hemic

Clav

Vegetation: Tamarack crown cover of about 60 percent with scattered black spruce; lush understory consists mostly of Labrador tea, leatherleaf, and grasses with swamp laurel and some sedges; ground cover consists of some Sphagnum mosses.

Microrelief: 30 cm.

plant.
Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Depth To Water Table: Near surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer	Depth (cm)
Fibric	0-106
Hemic	106-167
Sapric	167-183
Silty	183+
clay	

Location: 25 feet N and 1125 feet E of the SW corner of Sec. 28, T.159N., R.25W. (SE1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 28, T.159N., R.25W.)

Vegetation: Scattered black spruce; lush understory consists mostly of bog rosemary, swamp laurel, and leatherleaf with Labrador tea and some grasses; around cover consists of some Sphagnum mosses.

Microrellef: 20 cm.

Depth To Water Table: Near surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Fibric 0- 91
Hemic 91-122
Loam 122+

Location: 1725 feet N and 2540 feet W of the SE corner of Sec. 28, T.159N., R.25W. (NW1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 28, T.159N., R.25W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some sedges, swamp laurel, leatherleaf, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrelief: 50 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Fibric 0-91 Hemic 91-96 Clay 96+ Location: 2200 feet S and 150 feet W of the NE corner of Sec. 5, T.63N., R.26W. (SW1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 5, T.63N., R.26W.)

Vegetation: Scattered black spruce and paper birch; understory consists of sedges and grasses with some leatherleaf and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm
Hemic	0-61
Sapric	61-91
Sand	91+

Location: 110 feet S and 230 feet E of the NW corner of Sec. 5, T.63N., R.26W. (Gov. Lot 4, Sec. 5, T.63N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; understory consists of leatherleaf with some bog birch, bog rosemary, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 10 cm.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Fibric Hemic Fibric Hemic Loamy sand	0- 91 91-122 122-152 152-350 350+

Location: 100 feet S and 690 feet W of the NE corner of Sec. 9, T.63N., R.26W. (NE1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 9, T.63N., B.26W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; lush understory consists of leatherleaf with some bog birch, willow, sedges, grasses, swamp laurel, bog rosemary, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses.

Denth To Water Table: 10 cm. above surface.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Hemic	0-213
Sapric	213-244
Hemic	244-366
Loam	366+

Location: 75 feet N and 150 feet E of the SW corner of Sec. 11, T.63N., R.26W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 11, T.63N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; understory consists of some bog birch, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses. Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Hemic Sandy	0-305 305+
loam	

Location: 2050 feet S and 150 feet E of the NW corner of Sec. 13, T.63N., R.26W. (NW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 13, T.63N., R.26W.)

Vegetation: Black spruce grown cover of about 85 percent; sparse understory consists of some Labrador tea, leatherleaf, bog rosemary, sedges, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm)
Hemic	0-335
Sapric	335-366
Loamy	366+
sand	

Location: 475 feet N and 700 feet E of the SW corner of Sec. 19, T.63N., R.26W. (Gov. Lot 4, Sec. 19, T.63N., R.26W.)

Vegetation: Scattered black spruce and tamarack; understory consists of some Labrador tea, leatherleaf, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 40 cm.

Depth To Water Table: 10 cm.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm
Fibric	0- 61
Hemic	61- 91
Silty	91+
clay	
loam	

Location: 1200 feet S and 325 feet W of the NE corner of Sec. 23, T.63N., R.26W. (SE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 23, T.63N., R.26W.) Vegetation: Black spruce crown cover of about 35 percent with scattered

tamarack; understory consists of some bog birch, willow, leatherleaf, sedges, grasses, Labrador tea, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and B. Sether on April 11, 1980.

Layer	Depth (cm
Fibric	0- 61
Hemic	61-427+
Bottom	0. 121
unknown	

Location: 625 feet S and 1400 feet W of the NE corner of Sec. 30, T.63N., R.26W. (SE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 30, T.63N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; understory consists of some Labrador tea, leatherleaf, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses and blueberry.

Microrellet: 30 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Hemic Silty clay	0-122 122+

Location: 1155 feet N and 1275 feet E of the SW corner of Sec. 2, T.64N., R.26W. (NE1/4 of NE1/4 of SW1/4 of SW1/4, Sec. 2. T.64N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered paper birch: sparse understory consists of some black spruce, Labrador tea, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and snowberry.

Microrelief: 15 cm.

Depth To Water Table: 30 cm.

Described By: B. Sether and D. Haverkost on April 8, 1980.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-229
Silty	229+
clav	

Location: 175 feet S and 2075 feet W of the NE corner of Sec. 13. T.64N., R.26W. (NE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 13, T.64N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; understory consists of leatherleaf with some bog birch and bog rosemary; ground cover consists of some Sphagnum and feather mosses. Microrelief: 25 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Hemic Silty clay loam	0-152 152+

Location: 2150 feet S and 60 feet E of the NW corner of Sec. 19, T.64N., R.26W. (Gov. Lot 2, Sec. 19, T.64N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered paper birch; sparse understory consists of Labrador tea with some speckled alder and grasses; ground cover consists mostly of Sphagnum mosses with some snowberry and bunchberry.

Microrellef: 25 cm.

mosses.

Microrelief: 20 cm.

Microrelief: 46 cm.

Depth To Water Table: 30 cm.

Depth To Water Table: At surface.

Depth To Water Table: 10 cm.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm
Peat	0- 30
Clay	30+
loam	

Location: 450 feet N and 200 feet W of the SE corner of Sec. 36, T.64N., R.26W.

Vegetation: Understory consists of some sedges, grasses, bog birch, aider, willow.

black spruce, and ferns; ground cover consists of some Sphagnum and other

Location: 2410 feet N and 150 feet E of the SW corner of Sec. 7, T.65N., R.26W.

Vegetation: Scattered black spruce and tamarack; sparse understory consists of

cover consists mostly of Sphagnum mosses with some cranberry.

Described By: D. Mellem and P. Guertin on November 30, 1979.

leatherleaf, Labrador tea, swamp laurel, bog rosemary, and sedges; ground

(NE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 36, T.64N., R.26W.)

Described By: H. Mooers and B. Sether on April 11, 1980.

Laver

Hemic

Sapric

Silty

clav

(Gov. Lot 3, Sec. 7, T.65N., R.26W.)

Location: 980 feet N and 300 feet E of the SW corner of Sec. 30, T.64N., R.26W. (Gov. Lot 4, Sec. 30, T.64N., R.26W.)

Vegetation: Scattered black spruce; understory consists of some leatherleaf, swamp laurel, black spruce, and grasses; ground cover consists mostly of Sphagnum mosses with some feather mosses and cranberry.

Microrelief: 25 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Hemic	0-183
Loam	183 +

(Gov. Lot 3, Sec. 2, T.65N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of Labrador tea with some speckled alder and grasses; ground cover consists mostly of mosses with some Sphagnum mosses and snowberry.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Depth (cm) mic 0-30 pric 30-46 nd 46+

Location: 1200 feet S and 1850 feet E of the NW corner of Sec. 2, T.65N., R.26W.

Microrelief: 60 cm.

Depth To Water Table: Not visible.

Depth (cm)	Lay
0-61	Hen
61-91	Sap
91+	San

Location: 135 feet S and 2310 feet W of the NE corner of Sec. 19, T.65N., R.26W. (NW1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 19, T.65N., R.26W.)

Vegetation: Scattered tamarack and black spruce: lush understory consists mostly of sedges with some cotton grass, grasses, Labrador tea, and leatherleaf; around cover consists mostly of Sphagnum mosses with some other mosses. Microrellef: 35 cm.

Dopth To Water Table: 15 cm.

Described By: H. Mooers and S. Nelson on April 8, 1980.

Layer Depth (cm) Hemic 0-138 Sandy 138+ clay and fine sand

Depth (cm) Laver Hemic 0-138 Loamy 138 +sand

Location: 875 feet S and 2100 feet E of the NW corner of Sec. 31, T.64N., R.26W. (NE1/4 of SW1/4 of NE1/4 of NW1/4, Sec. 31, T.64N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some swamp laurel, bog rosemary, Labrador tea, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrelief: 35 cm.

Depth To Water Table: 20 cm.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Hemic	0-198
Sand	198+

Location: 2600 feet N and 2600 feet E of the SW corner of Sec. 4, T.65N., R.26W. (NE1/4 of NE1/4 of NE1/4 of SW1/4, Sec. 4, T.65N., R.26W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; understory consists of Labrador tea, leatherleaf, swamp laurel and some grasses; ground cover consists of Sphagnum mosses with some cranberry.

Microrelief: 25 cm.

Depth To Water Table: Below surface.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)
Hemic	0-198
Sapric	198-213
Medium	213+
sand	

Location: 115 feet N and 2410 feet E of the SW corner of Sec. 29, T.65N., R.26W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 29, T.65N., R.26W.)

Vegetation: Scattered black spruce; understory consists of leatherleaf with some grasses, alder, poplar, tamarack, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: B. Sether and D. Haverkost on April 8, 1980.

Layer	Depth (cm)
Hemic	0-30
Sapric	30-61
Medium	61+
sand	

Location: 270 feet N and 310 feet E of the SW corner of Sec. 30, T.65N. R.26W. (Gov. Lot 4, Sec. 30, T.65N., R.26W.)

Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf with some grasses, cotton grass, willow, Labrador tea, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some feather mosses.

Microrelief: 60 cm.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: B. Sether and D. Haverkost on April 8, 1980.

Layer Depth (cm)
Fibric 0- 91
Hemic 91-244
Sandy 244+
clay
loam

Location: 2575 feet N and 140 feet E of the SW corner of Sec. 5, T.66N., R.26W.

Vegetation: Northern white cedar crown cover of about 75 percent; sparse un-

Depth (cm)

0- 61

derstory consists of some grasses, bog birch, Labrador tea, and ferns.

(NW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 5, T.66N., R.26W.)

Described By: D. Meilem and P. Guertin on November 29, 1979.

Laver

Hemic

Clay

Location: 2275 feet N and 1150 feet E of the SW corner of Sec. 33. T.65N.. R.26W. (NE1/4 of NE1/4 of NW1/4 of SW1/4. Sec. 33. T.65N.. R.26W.)

Vegetation: Black spruce crown cover of about 65 percent: understory consists of leatherleaf with some black spruce. Labrador tea, and grasses: ground cover consists mostly of Sphagnum mosses with some feather mosses, cranberry, and bilberry.

Microrelief: 30 cm.

Depth To Water Table: 8 cm.

Described By: H. Mooers and S. Nelson on April 8, 1980.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-107
Loam	107+

Location: 75 feet N and 1225 feet E of the SW corner of Sec. 8, T.66N.. R.26W. (SE1/4 of SE1/4 of SW1/4, of SW1/4, Sec. 8, T.66N.. R.26W.)

Vegetation: Tamarack crown cover of about 85 percent with scattered black spruce and northern white cedar; understory consists of Labrador tea. leatherleaf, black ash, bog rosemary and grasses with some speckled alder and willow; ground cover consists of some Sphagnum and other mosses.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 27, 1979.

Layer	Depth (cm
Hemic Sandy	0-168 168+
clav	

Location: 700 feet S and 1560 feet E of the NW corner Sec. 11, T.66N., R.26W. (NW1/4 of SW1/4 of NE1/4 of NW1/4, Sec. 11, T.66N., R.26W.)

Location: 175 feet N and 230 feet W of the SE corner of Sec. 4 T.66N., R.26W.

Vegetation: Tamarack crown cover of about 30 percent; lush understory consists

Depth (cm)

0-548

548+

mostly of bog birch and grasses with leatherleaf and bog rosemary.

(SF1/4 of SF1/4 of SF1/4 of SF1/4, Sec. 4, T.66N., B.26W.)

Described By: D. Mellem and P. Guertin on November 29, 1979.

Laver

Hemic

Silt

over

clay

sandv

Vegetation: Scattered black spruce; lush understory consists mostly of sedges with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Microrelief: Negligible.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 29, 1979.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-274
Sandy	274+
clav	

Location: 100 feet S and 2580 feet W of the NE corner of Sec. 12, T.66N., R.26W. (NW1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 12, T.66N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent with scattered tamarack; sparse understory consists of speckled alder, bog birch and Labrador tea with some grasses; ground cover consists of some Sphagnum mosses.

Microrellef: 8 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 29 1979

Layer Depth (cm)
Hemic 0-183
Sandy 183 + clay

Location: 1550 feet S and 145 feet W of the NE corner of Sec. 14, T.66N., R.26W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 14, T.66N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some bog birch and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrelief: 8 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and P. Guertin on November 29, 1979.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-305
Clay	305+

Location: 175 feet S and 310 feet W of the NE corner of Sec. 16, T.66N., R.26W. (NW1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 16, T.66N., R.26W.)

Vegetation: Scattered black spruce, paper birch, and tamarack; sparse understory consists of grasses, speckled alder, Labrador tea, and leatherleaf; ground cover consists mostly of mosses.

Microrelief: 15 cm.

Depth To Water Table: 15 cm. above surface.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Layer	Depth (c
Hemic	0-335
Sapric	335-366
Medium	366+
to	
fine	
sand	

Location: 2000 feet S and 1040 feet W of the NE corner of Sec. 35, T.66N., R.26W. (SW1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 35, T.66N., R.26W.)

Vegetation: Scattered black spruce and paper birch; sparse understory consists of Labrador tea, leatherleaf, speckled alder, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

(Gov. Lot 1, Sec. 4, T.67N., R.26W.)

Depth To Water Table: At surface.

Sphagnum mosses with some other mosses.

Described By: D. Mellem on October 18, 1977.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Laver Depth (cm) 0-107 Hemic Mineral 107 +soil

Location: 575 feet N and 810 feet E of the SW corner of Sec. 3. T.67N., R.26W. (NW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 3, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 90 percent; sparse understory consists of some grasses and leatherleaf; ground cover consists mostly of Sphagnum mosses with some cranberry and other mosses.

Microrelief: 30 cm

Depth To Water Table: At surface.

Described By: D. Mellem on October 18, 1977.

Layer	Depth (cm)
Hemic	0-178
Clay	178+
loam	

Location: 2075 feet S and 250 feet E of the NW corner of Sec. 13, T.67N., R.26W. (NW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 13, T.67N., R.26W.)

Vegetation: Scattered black spruce; understory consists of sedges with some leatherleaf, cranberry, bog rosemary, and swamp laurel; ground cover consists

Depth To Water Table: At surface.

Laver Depth (cm) Fibric 0-15 15-30 Hemic Sapric 30-32 Clay 32 +loam

Location: 700 feet S and 1115 feet W of the NE corner of Sec. 4, T.67N., R.26W.

Vegetation: Black spruce crown cover of about 40 percent; sparse understory con-

sists of some Labrador tea and sedges; ground cover consists mostly of

mostly of Sphagnum mosses. Microrelief: 30 cm.

Described By: D. Mellem on October 20, 1977.

Layer	Depth (cm)
Fibric Hemic Sand	0- 17 17-269 269+
Odila	2031

Described By: D. Mellem on October 27, 1977.

Layer	Depth (cm)
Fibric	0- 25
Hemic	25-254
Sand	254+

Location: 2125 feet N and 490 feet W of the SE corner of Sec. 13, T.67N., R.26W.

Vegetation: Scattered black spruce; understory consists of leatherleaf and sedges

with some bog rosemary; ground cover consists mostly of Sphagnum mosses

(SW1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 13, T.67N., R.26W.)

Location: 2050 feet S and 450 feet W of the NE corner of Sec. 4, T.67N., R.26W.

Vegetation: Scattered black spruce; sparse understory consists of some

leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of

Depth (cm)

0- 50

50-115

115+

(SW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 4, T.67N., R.26W.)

Sphagnum mosses with some cranberry.

Described By: D. Mellem on October 18, 1977.

Laver

Fibric

Hemic

Clav

loam

Depth To Water Table: At surface.

with some cranberry.

Depth To Water Table: At surface.

Microrellef: 45 cm.

Microrellef: 30 cm.

Location: 2060 feet S and 1550 feet W of the NE corner of Sec. 14, T.67N., R.26W. (NE1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 14, T.67N., R.26W.)

Vegetation: Scattered black spruce; understory consists mostly of grasses with some swamp laurel and leatherleaf; ground cover consists mostly of Sphagnum mosses with some cranberry.

Depth (cm)

0-411

411+

Microrellef: 30 cm.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 19, 1977.

Layer

Peat

soil

Mineral

Location: 1800 feet S and 610 feet E of the NW corner of Sec. 14, T.67N., R.26W. (SE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 14, T.67N., R.26W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges with some leatherleaf, Labrador tea, bog rosemary, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 20, 1977.

Layer	Depth (cm)
Fibric	0- 47
Hemic	47-371
Sapric	371-418
Sand	418+

Location: 2200 feet N and 2150 feet E of the SW corner of Sec. 14, T.67N., R.26W. (SW1/4 of NE1/4 of NE1/4 of SW1/4, Sec. 14, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists mostly of sedges with some Labrador tea, bog rosemary, swamp laurel, and leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 20, 1977.

Layer	Depth (cm)
Fibric	0- 39
Hemic	39-365
Loamy	365+
sand	

Location: 90 feet N and 675 feet E of the SW corner of Sec. 14, T.67N., R.26W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 14, T.67N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some sedges, swamp laurel, bog rosemary, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 20, 1977.

Location: 2200 feet N and 2100 feet W of the SE corner of Sec. 14, T.67N., R.26W. (SE1/4 of NW1/4 of NW1/4 of SE1/4, Sec. 14, T.67N., R.26W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges with some leatherleaf, Labrador tea, bog rosemary, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 35 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 20, 1977.

Layer	Depth (cm)
Fibric	0- 24
Hemic	24-316
Loamy	316-322
sand	
Silty	322+
clay	

Location: 2240 feet N and 1670 feet W of the SE corner of Sec. 15, T.67N., R.26W. (SW1/4 of NE1/4 of NW1/4 of SE1/4, Sec. 15, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent; understory consists of sedges and leatherleaf with some bog rosemary and swamp laurel; ground cover consists mostly of Sobagnum mosses with some crapherry.

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 20, 1977.

Layer	Depth (cm)
Fibric	0- 37
Hemic	37-370
Mineral	370+
soil	

Location: 800 feet N and 1680 feet W of the SE corner of Sec. 15, T.67N., R.26W. (SE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 15, T.67N., R.26W.)

Vegetation: Scattered black spruce; sparse understory consists of some leatherleaf, sedges, bog rosemary, Labrador tea, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 20, 1977.

Layer	Depth (cm)
Fibric	0- 27
Hemic	27-274
Loamy	274+
eand	

Location: 775 feet N and 680 feet W of the SE corner of Sec. 15, T.67N., R.26W. (SE1/4 of NW1/4 of SE1/4 of SE1/4, Sec. 15, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 45 percent; sparse understory consists of some sedges, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 20, 1977.

Layer	Depth (cm)
Fibric	0- 22
Hemic	22- 79
Fibric	79-100
Hemic	100-333
Loamy	333+
sand	

Location: 625 feet N and 575 feet E of the SW corner of Sec. 17, T.67N., R.26W. (NE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 17, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 85 percent; understory consists of Labrador tea with some leatherleaf, sedges, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry, false Solomon's seal, and reindeer moss.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Wakanabo on October 19, 1977.

Layer	Depth (cm
Hemic	0-130
Sandy clay	130+
loam	

Location: 650 feet N and 1440 feet W of the SE corner of Sec. 18, T.67N., R.26W. (SE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 18, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 85 percent; understory consists of leatherleaf with some sedges, bog rosemary, swamp laurel, and bog birch; ground cover consists mostly of Sphagnum mosses with cranberry and some reindeer moss.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: R. Wakanabo on October 19, 1977.

Layer	Depth (cm)
Hemic Loamy sand	0-140 140+

Location: 675 feet N and 430 feet W of the SE corner of Sec. 18, T.67N., R.26W. (SW1/4 of NE1/4 of SE1/4 of SE 1/4, Sec. 18, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 95 percent; understory consists of some bog rosemary, leatherleaf, bog birch, and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry and reindeer moss.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: R. Wakanabo on October 19, 1977.

Layer	Depth (cm)
Hemic	0-130
Clay	130+
loam	

Location: 660 feet S and 1950 feet W of the NE corner of Sec. 19, T.67N., R.26W. (SW1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 19, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 80 percent; understory consists of leatherleaf with some bog rosemary, swamp laurel, bog birch, and sedges; ground cover consists mostly of Sphagnum mosses with cranberry and other mosses.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: R. Wakanabo on October 19, 1977.

Layer	Depth (cm)
Hemic	0-170
Limnic	170-200
Mineral	200+
soil	

Location: 675 feet S and 2225 feet W of the NE corner on Sec. 19, T.67N., R.26W. (SE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 19, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 95 percent; understory consists of Labrador tea with some leatherleaf, sedges, and bog rosemary; ground cover consists mostly of Sphagnum mosses with cranberry.

Microrellef: 40 cm.

Depth To Water Table: 5 cm.

Described By: R. Wakanabo on October 19, 1977

Layer Depth (cm) Hemic 0-50 Sand 50+

Location: 625 feet S and 1110 feet E of the NW corner of Sec. 21, T.67N., R.26W. (SE1/4 of NE1/4 of NW1/4 of NW1/4, Sec. 21 T.67N., R.26W.)

Vegetation: Scattered black spruce; understory consists of sedges with some leatherleaf and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 26, 1977.

Layer	Depth (cm)
Fibric	0- 29
Hemic	29-309
Sand	309-321
Silty	321+
clay	

Location: 25 feet N and 2010 feet E of the SW corner of Sec. 22, T.67N., R.26W. (SW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 22, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 95 percent; understory consists of some leatherleaf, bog rosemary, swamp laurel, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Described By: T. Malterer on October 27, 1977.

Depth (cm
0- 45
45-410
410+

Location: 2500 feet N and 2360 feet W of the SE corner of Sec. 23, T.67N., R.26W. (NW1/4 of NW1/4 of NW1/4 of SE1/4, Sec. 23, T.67N., R.26W.)

Vegetation: Black spruce crown cover of about 50 percent with scattered tamarack; lush understory consists mostly of leatherleaf and sedges with some Labrador tea, swamp laurel, bog rosemary, and grasses; ground cover consists of false Solomon's seal and Sphagnum mosses with some other mosses. Microrelief: 25 cm.

Depth To Water Tabie: At surface.

Described By: B. Wakanabo on October 26, 1977.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-210
Sand	210+

Location: 1425 feet N and 50 feet W of the SE corner of Sec. 32, T.67N., R.26W. (SE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 32, T67N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of speckled alder with some leatherleaf, cotton grass, ferns, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Described By: D. Olson on October 25, 1977.

Layer	Depth (cm)
Hemic	0-210
Silty	210+
clay	

Location: 2440 feet N and 1975 feet W of the SE corner of Sec. 2, T.68N., R.26W. (NE1/4 of NW1/4 of NW1/4 of SE1/4, Sec. 2, T.68N., R.26W.)

Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf, Labrador tea, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses. cranberry, lingen berry, and false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: 5 cm.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

.ayer	Depth (cm)
ibric	0- 91
lemic	91-305
ilty	305+
lay	

Location: 2050 feet S and 875 feet E of the NW corner of Sec. 8, T.68N., R.26W. (NW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 8, T.68N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of Labrador tea with some swamp laurel, sedges, and balsam fir: ground cover consists of some cranberry, snowberry, lingen berry, and blueberry.

Microrellef: 20 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Olson on November 15, 1979.

Hemic 0-229	Layer	Depth (cm)
loam	Clay	0-229 229+

Location: 1820 feet S and 80 feet W of the NE corner of Sec. 10, T.68N., R.26W. (SE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 10, T.68N., R.26W.)

Vegetation: Scattered northern white cedar, tamarack, and black spruce: understory consists of bog birch, Labrador tea and leatherleaf with some dogwood and grasses; ground cover consists mostly of Sphagnum mosses with some pitcher plant.

Microrelief: 35 cm.

Depth To Water Table: At surface.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer	Depth (cm)
Hemic	0-138
Sapric	138-152
Silty	152+
clav	

Location: 2100 feet S and 1775 feet E of the NW corner of Sec. 14, T.68N., R.26W. (NE1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 14, T.68N., R.26W.)

Vegetation: Scattered black spruce, northern white cedar and tamarack, lush understory consists of bog birch, dogwood, Labrador tea, leatherleaf, sedges, ferns, aster, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses and false Solomon's seal.

Microrelief: 20 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

Layer	Depth (cm)
Hemic Sapric Silty clay and very fine	0-274 274-290 290+
sand	

Location: 775 feet S and 2350 feet W of the NE corner of Sec. 18, T.68N., R.26W. (NW1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 18, T.68N., R.26W.)

Vegetation: Scattered black spruce, tamarack, and paper birch; sparse understory consists of some Labrador tea and grasses; ground cover consists mostly of feather mosses with some Sphagnum mosses.

Microrellef: Negligible.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm
Hemic Silty clay	0-91 91+

Location: 1625 feet N and 1700 feet W of the SE corner of Sec. 28, T.68N., R.26W. (SW1/4 of SW1/4 of NE1/4 of SE1/4, Sec. 28, T.68N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent; understory consists of Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 19, 1977.

Layer	Depth (cm)
Fibric	0- 7
Hemic	7- 65
Sapric	65-118
Sandy	118+
loam	

Location: 1205 feet N and 1725 feet W of the SE corner of Sec. 28, T.68N., R.26W. (NW1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 28, T.68N., R.26W.) Vegetation: Scattered black spruce; understory consists of some Labrador tea;

regetation: Scattered black spruce; understory consists of some Labrador tea; ground cover consists mostly of Sphagnum mosses with some cramberry.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 19, 1977.

Layer	Depth (cm)
Fibric	0- 10
Hemic	10-215
Sandy	215+
clav	
loam	

Location: 10 feet N and 2225 feet W of the SE corner of Sec. 29, T.68N., R.26W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 29, T.68N., R.26W.)

Vegetation: Logged area with some Labrador tea, grasses, and cranberry. Microrellef: 25 cm.

Depth To Water Table: 2 cm.

Described By: D. Mellem on October 19, 1977.

Layer	Depth (cm
Fibric	0- 8
Hemic	8-101
Silt	101+
loam	

Location: 10 feet N and 1200 feet W of the SE corner of Sec. 29, T.68N., R.26W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 29, T.68N., R.26W.)

Vegetation: Scattered black spruce; understory consists mostly of Labrador tea with some sedges; ground cover consists mostly of Sphagnum mosses.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 19, 1977.

Layer	Depth (cm)
Fibric Hemic Silty clay loam	0- 10 10-130 130+

Location: 2220 feet S and 1025 feet E of the NW corner of Sec. 30, T.68N., R.26W. (Gov. Lot 2, Sec. 30, T.68N., R.26W.)

Vegetation: Black spruce crown cover of about 80 percent; understory consists of some Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses and snowberry.

Microrelief: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 29, 1979.

Layer	Depth (cm)
Hemic	0-107
Clay	107+

Location: 990 feet S and 2230 feet W of the NE corner of Sec. 32, T.68N., R.26W. (NE1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 32, T.68N., R.26W.)

Vegetation: Black spruce crown cover of about 90 percent, sparse understory consists of some Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 19, 1977.

Layer	Depth (cm)
Fibric	0-10
Hemic	10-60
Silty	60+
clay	
ioam	

Location: 10 feet N and 675 feet W of the SE corner of Sec. 32, T.68N., R.26W. (SE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 32, T.68N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of sedges with Labrador tea and some leatherleaf, cotton grass, and grasses; ground cover consists of Sphagnum mosses with some cranberry, false Solomon's seal, and other mosses.

Microrelief: 20 cm.

Depth To Water Table: 5 cm.

Described By: D. Olson and B. Wakanabo on October 5, 1977.

Layer	Depth (cm)
Peat	0-79
Mineral	79+
soil	

Location: 1250 feet S and 2300 feet E of the NW corner of Sec. 33, T.68N., R.26W. (SW1/4 of SE1/4 of NE1/4 of NW1/4, Sec. 33, T.68N., R.26W.)

Vegetation: Not recorded.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Olson on October 18, 1977.

.ayer	Depth (cm)
ibric	0- 10
lemic	10-120
Coarse	120-125
and	
Clay	125+
oam	

Location: 30 feet N and 50 feet E of the SW corner of Sec. 33, T.68N., R.26W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 33, T.68N., R.26W.)

Vegetation: Black spruce crown cover of about 90 percent; lush understory consists of grasses. Labrador tea, and leatherleaf with some cotton grass and sedges; ground cover consists of false Solomon's seal and Sphagnum mosses with some other mosses and cranberry.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Olson and B. Wakanabo on October 5, 1977.

Layer	Depth (cm)
Peat	0-76
Mineral	76+
soil	

Location: 1900 feet N and 2250 feet W of the SE corner of Sec. 33, T.68N., R.26W.

(NE1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 33, T.68N., R.26W.) Vegetation: Black spruce crown cover of about 95 percent; understory consists of some Labrador tea, leatherleaf, swamp laurel, bog rosemary, sedges and grasses; ground cover consists mostly of false Solomon's seal with some Sphagnum and feather mosses.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: T. Malterer on October 18, 1977.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-330
Sapric	330-345
Limnic	345-355
Loamy	355+
sand	

Location: 10 feet N and 2500 feet W of the SE corner of Sec. 34, T.68N., R.26W. (SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 34, T.68N., R.26W.)

Vegetation: Tamarack crown cover of about 40 percent with a black spruce crown cover of about 30 percent; lush understory consists mostly of leatherleaf with grasses, bog rosemary, and some Labrador tea, swamp laurel, sedges, and cotton grass; ground cover consists of cranberry and Sphagnum mosses with some false Solomon's seal.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Described By: D. Olson and B. Wakanabo on October 5, 1977.

Layer	Depth (cm)
Peat	0-189
Clay	189+
with	
pebbles	

Location: 10 feet N and 1010 feet E of the SW corner of Sec.33, T.68N., R.26W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 33, T.68N., R.26W.)

Vegetation: Black spruce crown cover of about 55 percent with scattered tamarack; lush understory consists mostly of cotton grass with Labrador tea, sedges and some bog rosemary and grasses; ground cover consists of Sphagnum mosses with some false Solomon's seal.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Olson and B. Wakanabo on October 5, 1977.

Layer	Depth (cm)
Peat	0-114
Mineral	114+
nail .	

Location: 20 feet N and 1510 feet W of the SE corner of Sec. 33., T.68N., R.26W. (SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 33, T.68N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of leatherleaf with some Labrador tea, bog rosemary, and sedges; ground cover consists of Sphagnum mosses with some other mosses and cranberry. Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: T. Malterer on October 18, 1977.

Layer	Depth (cm)
Fibric Hemic Clay loam	0- 25 25-100 100+

Location: 15 feet N and 675 feet E of the SW corner of Sec. 35, T.68N., R.26W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 35, T.68N., R.26W).

Vegetation: Black spruce crown cover of about 90 percent; understory consists of grasses with some Labrador tea, swamp laurel, bog rosemary, and cotton grass; ground cover consists of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: B. Wakanabo on October 5, 1977.

Layer	Depth (cm)
Peat	0-330
Mineral	330+
eoil	

Location: 2550 feet N and 2505 feet W of the SE corner of Sec. 33, T.68N., R.26W. (NW1/4 of NW1/4 of NW1/4 of SE1/4, Sec. 33, T.68N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of leatherleaf with some Labrador tea, sedges, grasses, bog rosemary, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on October 18, 1977.

Layer	Depth (cm
Fibric Hemic Sapric Sandy	0- 40 40-190 190-195 195+
loam	

Location: 2300 feet S and 725 feet W of the NE corner of Sec. 34, T.68N., R.26W. (NE1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 34, T.68N., R.26W.)

Vegetation: Northern white cedar crown cover of about 40 percent with a black spruce crown cover of about 35 percent and scattered tamarack; lush understory consists of Labrador tea and cotton grass with some leatherleaf. swamp laurel, bog rosemary, and grasses; ground cover consists of cranberry with some Sphagnum and other mosses.

Microrellef: 35 cm.

Depth To Water Table: 10 cm.

Described By: B. Wakanabo on October 5, 1977.

Layer	Depth (cm
Peat Sandy clay loam with pebbles	0-335 335+

Location: 1000 feet N and 1600 feet W of the SE corner of Sec. 2, T.69N., R.26W. (NE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 2, T.69N., R.26W.)

Vegetation: Scattered black spruce and jack pine; lush understory consists mostly of sedges with some Labrador tea, leatherleaf, swamp laurel, and bog rosemary: ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 15 cm.

Depth To Water Table: 5 cm.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

ayer	Depth (cm
lemic	0-290
Silty	290+
clay	

Location: 1675 feet S and 175 feet E of the NW corner of Sec. 3, T.69N., R.26W. (SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 3, T.69N., R.26W.)

Vegetation: Black spruce and tamarack crown cover of about 35 percent; lush understory consists mostly of Labrador tea, swamp laurel, and leatherleaf with some grasses; ground cover consists mostly of cranberry and feather mosses with some Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Near surface.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer Depth (cm)
Hemic 0-213
Silty 213+
clay

Graber of November 16, 1915.

 Layer
 Depth (cm)

 Hemic
 0-213

 Sapric
 213-229

 Clay
 229+

Location: 400 feet S and 1975 feet W of the NE corner of Sec. 10, T.69N., R.26W.

Vegetation: Understory consists mostly of sedges and cotton grass with some

swamp laurel, Labrador tea, and bog rosemary; ground cover consists mostly

Depth (cm)

0-290

290+

(SW1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 10, T.69N., R.26W.)

of Sphagnum mosses with some other mosses and cranberry.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer

Hemic

Silty

Location: 2430 feet S and 1850 feet E of the NW corner of Sec. 3, T.69N., R.26W.

Vegetation: Scattered tamarack and black spruce; understory consists of leatherleaf, cotton grass, Labrador tea, and swamp laurel; ground cover con-

sists mostly of Sphagnum mosses with some other mosses and cranberry.

(SE1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 3. T.69N., R.26W.)

Described By: D. Olson and R. Graber on November 16, 1979.

Microrellef: 30 cm.

Depth To Water Table: 40 cm.

Location: 1350 feet N and 1590 feet E of the SW corner of Sec. 9, T.69N., R.26W. (SW1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 9, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent with a tamarack crown cover of about 30 percent; lush understory consists mostly of Labrador tea and leatherleaf with grasses; ground cover consists mostly of cranberry with false Solomon's seal.

Microrellef: 25 cm.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 26, 1977.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-148
Silty	148+
clay	
with	
calcareous	
pebbles	

Location: 725 feet S and 1700 feet W of the NE corner of Sec. 15, T.69N., R.26W. (NW1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 15, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf with some Labrador tea and swamp laurel; ground cover consists mostly of Sphagnum mosses and false Solomon's seal with cranberry.

Microrellef: 35 cm.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 26, 1977.

loam

 Layer
 Depth (cm)

 Hemic
 0-198

 Sapric
 198-213

 Clay
 213+

Location: 2255 feet S and 125 feet W of the NE corner of Sec. 14, T.69N., R.26W.

Vegetation: Scattered black spruce; lush understory consists mostly of sedges with

some swamp laurel and Labrador tea; ground cover consists mostly of

(NE1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 14, T.69N., R.26W.)

Described By: D. Olson and R. Graber on November 16, 1979.

Sphagnum mosses with some other mosses.

Layer Depth (cm)
Peat 0-234
Sandy 234+
clay

Location: 2605 feet S and 700 feet E of the NW corner of Sec. 6, T.69N., R.26W. (Gov. Lot 5, Sec. 6, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of poplar, sedges, leatherleaf, grasses, and horsetall; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 15 cm.

Depth To Water Table: 10 cm. above surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer	Depth (cm)	
Fibric	0- 91	
Hemic	91-122	
Medium	122+	
sand		

Location: 650 feet N and 1510 feet W of the SE corner of Sec. 11, T.69N., R.26W. (SE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 11, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some swamp laurel, Labrador tea, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

Layer	Depth (cm)	
Fibric	0- 61	
Hemic	61-198	
Silty	198+	
clay		

Location: 2500 feet S and 1550 feet W of the NE corner of Sec. 15, T.69N., R.26W. (SE1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 15, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of Labrador tea with some leatherleaf and grasses; ground cover consists mostly of Sphagnum mosses with cranberry and some false Solomon's seal and other mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 26, 1977.

Layer	Depth (cm	
Peat	0-125	
Sandy	125+	
clay		
loam		

Location: 10 feet S and 2580 feet E of the NW corner of Sec. 15, T.69N., R.26W. (NF1/4 of NF1/4 of NF1/4 of NW1/4. Sec. 15, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 50 percent with scattered tamarack; lush understory consists mostly of leatherleaf with Labrador tea, grasses, and some swamp laurel; ground cover consists mostly of Sphagnum mosses and cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 26, 1977.

Layer Depth (cm)
Peat 0-230
Mineral 230+
soil

Location: 2590 feet N and 1400 feet W of the SE corner of Sec. 15, T.69N., R.26W. (NE1/4 of NE1/4 of NW1/4 of SE1/4, Sec. 15, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent; understory consists of Labrador tea with swamp laurel; ground cover consists of cranberry, Sphagnum mosses, and other mosses.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 26, 1977.

Layer	Depth (cm)
Peat	0-119
Sandy	119+
clay	
oam	

Location: 10 feet S and 2215 feet W of the NE corner of Sec. 16, T.69N., R.26W. (NE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 16, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with a tamarack crown cover of about 35 percent; lush understory consists mostly of leatherleaf and grasses with some swamp laurel; ground cover consists mostly of cranberry and Sphagnum mosses.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 26, 1977.

_ayer	Depth (cm)
Peat	0-231
Silt	231+

Location: 1725 feet E and 10 feet S of the NW corner of Sec. 16, T.69N., R.26W. (NE1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 16, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent; lush understory consists mostly of Labrador tea and leatherleaf with some grasses; ground cover consists of Sphagnum mosses and cranberry.

Microrelief: 20 cm

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 26, 1977.

Layer	Depth (cm
Hemic	0-55
Sandy	55+
clay	
loam	

Location: 20 feet N and 1875 feet W of the SE corner of Sec. 16, T.69N., R.26W. (SW1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 16, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; understory consists mostly of cotton grass with some sedges, Labrador tea, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microsellet: 30 cm

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Olson on November 15, 1979.

Layer	Depth (cm)	
Fibric	0-152	
Hemic	152-381	
Clay	381+	

Location: 810 feet S and 1600 feet W of the NE corner of Sec. 17, T.69N., R.26W. (NW1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 17, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and snowberry.

Microrelief: 30 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Depth (cm)
0-122
122-152
152+

Location: 1575 feet S and 1860 feet E of the NW corner of Sec. 19, T.69N., R.26W. (NE1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 19, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 90 percent; understory consists mostly of Labrador tea with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 20 cm.

Depth To Water Table: 25 cm.

Described By: D. Olson on August 23, 1977.

Layer Depth (cm)
Hemic 0-100
Clay 100+

Location: 1590 feet S and 675 feet E of the NW corner of Sec. 19, T.69N., R.26W. (Gov. Lot 2, Sec. 19, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 95 percent; lush understory consists mostly of leatherleaf and Labrador tea; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 35 cm.

Depth To Water Table: 25 cm.

Described By: D. Olson on August 23, 1977.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-223
Silty	223+
clay	

Location: 2275 feet S and 125 feet E of the NW corner of Sec. 22, T.69N., R.26W. (NW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 22, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; lush understory consists mostly of grasses with leatherleaf and some Labrador tea and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 28, 1977.

.ayer	Depth (cm
Peat Sandy Slay oam	0-270 270+

Location: 1040 feet N and 200 feet E of the SW corner of Sec. 22. T.69N.. R.26W. (NW1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 22, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of leatherleaf and grasses with Labrador tea and some bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry and other mosses.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 28, 1977.

Depth (cm) Layer Peat 0-297 Clay 297+

Location: 25 feet N and 250 feet E of the SW corner of Sec. 22, T.69N., R.26W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 22, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists of leatherleaf and swamp laurel with some bod rosemary, grasses, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some

Microrelief: 20 cm.

Depth To Water Table: At surface

Described By: B. Wakanabo on September 28, 1977.

Layer	Depth (cm)
Peat	0-270
Clay	270+

Location: 2450 feet S and 2225 feet F of the NW corner of Sec. 23, T.69N., R26W. (SW1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 23, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists of sedges with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: Negligible.

Depth To Water Table: Not recorded.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer	Depth (cm
Fibric	0-138
Hemic	138-229
Loam	229+

Location: 1375 feet S and 1175 feet E of the NW corner of Sec. 24, T.69N., R.26W. (SE1/4 of SE1/4 of NW1/4 of NW1/4, Sec. 24, T.69N., R.26W.)

Vegetation: Scattered black spruce; understory consists of cotton grass and Labrador tea with some leatherleaf, sedges, swamp laurel, and bog rosemary: ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

Location: 1950 feet S and 1840 feet W of the NE corner of Sec. 25, T.69N., B.26W. (SW1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 25, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 45 percent; understory consists of Labrador tea and grasses with some speckled alder; ground cover consists mostly of Sphagnum mosses with some cranberry and other mosses.

Depth (cm)

0-31

31+

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: D. Olson on October 3, 1977.

Layer

Hemic

Silty

clav

Location: 1060 feet S and 525 feet E of the NW corner of Sec. 25, T.69N., R.26W.
(SE1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 25, T.69N., R.26W.)
Vegetation: Understory consists of black spruce and leatherleaf with some
tamarack ewamp laural had recemeny and cotton grees; ground cover con-

tamarack, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of false Solomon's seal with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Wakanabo on October 3, 1977.

Layer	Depth (cm)
Peat	0-230
Silty	230+
clay	

Layer	Depth (cm)
Hemic	0- 91
Sapric	91-138
Silty	138+
clay	

Location: 1650 feet S and 2425 feet E of the NW corner of Sec. 25, T.69N., R.26W.

Vegetation: Black spruce crown cover of about 65 percent; lush understory con-

sists mostly of Labrador tea and leatherleaf; ground cover consists mostly of

(NE1/4 of NE1/4 of SE1/4 of NW1/4, Sec. 25, T.69N., R.26W.)

Location: 1430 feet S and 1450 feet E of the NW corner of Sec. 25, T.69N., R.26W. (NW1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 25, T.69N., R.26W.)

Vegetation: Scattered black spruce; lush understory consists mostly of leatherleaf with some grasses; ground cover consists of some cranberry and false Solomon's seal.

Microrelief: Not recorded.

Depth To Water Table: At surface. Described By: D. Olson on October 3, 1977.

Layer	Depth (cm)
Sapric	0-137
Mineral	137+
soil	

Location: 675 feet S and 250 feet W of the NE corner of Sec. 26, T.69N., R.26W. (SE1/4 of NE1/4 of NE1/4, Sec. 26, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; understory consists of Labrador tea and leatherleaf with some swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry. Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on October 3, 1977.

.ayer	Depth (cm)
Peat	0-300
Silt	300+

Layer Depth (cm) Sapric Mineral 123+

soil

Sphagnum mosses with some cranberry.

Described By: D. Olson on October 3, 1977.

Microrellef: Not recorded.

Depth To Water Table: At surface

Location: 675 feet S and 1210 feet W of the NE corner of Sec. 26, T.69N., R.26W. (SW1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 26, T.69N., R.26W.)

Vegetation: Not recorded.
Microrellet: Not recorded.

Depth To Water Table: Not recorded.

Described By: B. Wakanabo on October 3, 1977.

 Layer
 Depth (cm)

 Hemic
 0-266

 Silt
 266+

Location: 675 feet S and 2200 feet W the NE corner of Sec. 26, T.69N., R.26W. (SE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 26, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 50 percent with scattered tamarack; understory consists of grasses with some leatherleaf, Labrador tea, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on October 3, 1977.

Layer	Depth (cm)
Peat	0-312
Silt	312+

Location: 1975 feet N and 875 feet E of the SW corner of Sec. 26, T.69N., R.26W. (SW1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 26, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack; lush understory consists mostly of leatherleaf with Labrador tea, grasses, and some swamp laurel and bog rosemary; ground cover consists mostly of Sphagnum mosses with cranberry, and some other mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: B. Wakanabo on September 27, 1977.

Layer	Depth (cm)
Peat	0-340
Silt	340+

Location: 650 feet N and 900 feet W of the SE corner of Sec. 26, T.69N., R.26W. (NF1/4 of SW1/4 of SF1/4 of SF1/4. Sec. 26, T.69N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of Labrador tea, leatherieaf, and swamp laurel with grasses; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 27, 1977.

Layer	Depth (cm)
Peat	0-158
Sandy	158+
clay	
loam	

Location: 1975 feet S and 1960 feet W of the NE corner of Sec. 27, T.69N., R.26W. (SW1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 27, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack: lush understory consists mostly of leatherleaf with Labrador tea, swamp laurel, and some grasses; ground cover consists mostly of Sphagnum mosses with some crapherry and other mosses.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 28, 1977.

Layer	Depth (cm)
Peat	0-347
Clay	347+

Location: 1010 feet S and 255 feet E of the NW corner of Sec. 27, T.69N., R.26W. (NW1/4 of SW1/4 of NW1/4, Sec. 27, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; lush understory consists of Labrador tea, leatherleaf, and grasses with some cotton grass, swamp laurel, and bog rosemary; ground cover consists of Sphagnum mosses with some cranberry and other mosses.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 28, 1977.

Layer	Depth (cm)
Peat Sandy clay loam	0-267 267+

Location: 1950 feet S and 300 feet E of the NW corner of Sec. 27 T.69N., R.26W. (SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 27, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack; lush understory consists of grasses and leatherleaf with Labrador tea, swamp laurel, bog rosemary, and some sedges; ground cover consists mostly of Sohagnum mosses with some pitcher plant.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 28, 1977.

Layer	Depth (cm)
Peat	0-238
Clay	238+

Location: 2025 feet N and 975 feet E of the SW corner of Sec. 27, T.69N., R.26W. (SE1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 27, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 30 percent with scattered tamarack; understory consists of grasses and leatherleaf with Labrador tea and some swamp laurel and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry and other mosses.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on October 27, 1977.

Layer	Depth (cm)
Peat	0-283
Silty	283+
clav	

Location: 2050 feet N and 20 feet E of the SW corner of Sec. 27, T.69N., R.26W. (SW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 27, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 75 percent; lush understory consists of Labrador tea and leatherleaf with swamp laurel and grasses; ground cover consists of Sphagnum mosses with some cranberry and other mosses. Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 27, 1977.

Layer	Depth (cm)
Peat	0-230
Silty	230+
clay	

Location: 675 feet N and 2120 feet E of the SW corner of Sec. 27, T.69N., R.26W. (SW1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 27, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 50 percent; lush understory consists mostly of Labrador tea with swamp laurel and grasses; ground cover consists mostly of Sphagnum mosses and cranberry with some false Solomon's seal and other mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: B. Wakanabo on September 27, 1977.

Layer Depth (cm)
Hemic 0-79
Clay 79+

Location: 1990 feet N and 140 feet W of the SE corner of Sec. 27, T.69N., R.26W. (SE1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 27, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 50 percent; lush understory consists of Labrador tea and leatherleaf with some swamp laurel, grasses, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellef: 40 cm.

Depth To Water Table: 5 cm.

Described By: B. Wakanabo on September 27, 1977.

Layer	Depth (cm)
Peat Silty clay	0-358 358+

Location: 450 feet S and 20 feet E of the NW corner of Sec. 34, T.69N., R.26W. (SE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 34, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 50 percent with scattered tamarack; lush understory consists mostly of Labrador tea with some swamp laurel and sedges; ground cover consists mostly of Sphagnum mosses with other mosses and some cranberry.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Olson on October 4, 1977.

Layer	Depth (cm)
Hemic Silty	0-152 152+
clay	

Location: 1800 feet S and 310 feet W of the NE corner of Sec. 34, T.69N., R.26W. (SE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 34, T.69N., R.26W.)

Vegetation: Scattered tamarack and black spruce; lush understory consists of Labrador tea and leatherleaf with some sedges; ground cover consists of some cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Oison on October 4, 1977.

Layer	Depth (cm)
Hemic Mineral soil	0-131 131+

Location: 20 feet N and 20 feet E of the SW corner of Sec. 34, T.69N., R.26W. (SW1/4 of SW1/4 of SW1/4, Sec. 34, T.69N., R.26W.)
Vecetation: Black spruce crown cover of about 75 percent; understory consists of

Vegetation: Black spruce crown cover of about 75 percent; understory consists of Labrador tea with some leatherleaf, swamp laurel, cotton grass, and bog rosemary; ground cover consists of Sphagnum mosses with some cranberry. Microrellef: 35 cm.

Location: 2060 feet N and 1010 feet W of the SE corner of Sec. 28, T.69N., R.26W.

Vegetation: Black spruce crown cover of about 75 percent with scattered

tamarack; lush understory consists mostly of Labrador tea with bog rosemary

and some leatherleaf and swamp laurel; ground cover consists of snowberry

Depth (cm)

0-130

130+

(SW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 28, T.69N., R.26W.)

with some cranberry and mosses.

Described By: B. Wakanabo on September 27, 1977.

Layer

Peat

Silty

clay

Depth To Water Table: At surface.

Microrellef: 15 cm.

Depth To Water Table: 5 cm.

Described By: R. Wakanabo on October 4, 1977.

Layer	Depth (cm)
Peat	0-112
Silt	112+

Location: 20 feet N and 690 feet E of the SW corner of Sec. 34, T.69N., R.26W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 34, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent; understory consists of Labrador tea with some cotton grass, bog rosemary, swamp laurel, and leatherleaf; ground cover consists of some cranberry.

Microrellef: 25 cm.

Depth To Water Table: 5 cm.

Described By: R. Wakanabo on October 4, 1977.

Layer	Depth (cm)
Peat Mineral soil	0-183 183+

Location: 975 feet N and 2510 feet E of the SW corner of Sec. 34, T.69N., R.26W. (SE1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 34, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack; understory consists of Labrador tea with some leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 35 cm.

Depth To Water Table: 5 cm.

Described By: R. Wakanabo on October 4, 1977.

Layer	Depth (cm)
Peat	0-160
Clay	160+

Location: 2025 feet S and 1860 feet W of the NE corner of Sec. 36, T.69N., R.26W. (NW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 36, T.69N., R.26W.)

Vegetation: Black spruce crown cover of about 30 percent; lush understory consists mostly of leatherleaf with some Labrador tea, cotton grass, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 25 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and R. Graber on November 16, 1979.

Layer	Depth (cm)
Hemic	0-381
Sapric	381-396
Silty	396+
clay	

Location: 1425 feet N and 1090 feet E of the SW corner of Sec. 36, T.70N., R.26W. (SE1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 36, T.70N., R.26W.)

Vegetation: Scattered black spruce; understory consists of some cotton grass, leatherleaf, Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, false Solomon's seal, lingen berry, and lichens.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

mostly of Sphagnum mosses.

Depth To Water Table: Not visible

Microrellef: Negligible.

Described By: N. Aaseng and D. Haverkost on November 16, 1979.

Layer Depth (cm)
Fibric 0- 91
Hemic 91-198
Silty 198+
clay

Location: 1875 feet S and 1915 feet E of the NW corner of Sec. 18, T.152N., R.26W.

Vegetation: Black spruce crown cover of about 80 percent; lush understory con-

sists of Labrador tea and leatherleaf with some grasses; ground cover consists

Depth (cm)

0-152

152+

(SE1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 18, T.152N., R.26W.)

Described By: D. Mellem and D. Haverkost on December 10, 1979.

Layer

Hemic

Sandy

clay

Location: 2280 feet N and 100 feet E of the SW corner of Sec. 36, T.151N., R.26W. (NW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 36, T.151N., R.26W.)

Vegetation: Scattered northern white cedar and paper birch; lush understory consists mostly of bog birch and sedges with some Labrador tea. speckled alder, willow, dogwood, grasses, fern, iris, and swamp cinquefoil; ground cover consists of Sphagnum mosses and some lichens.

Microrelief: 35 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and N. Aaseng on December 10, 1979.

Layer	Depth (cm)
Hemic	0-64
Sapric	64-96
Silty	96+
clay	
loam	

Location: 1350 feet N and 1800 feet E of the SW corner of Sec. 23, T.152N., R.26W. (SE1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 23, T.152N., R.26W.)

Vegetation: Scattered black spruce, tamarack, and northern white cedar; lush understory consists of dogwood and grasses with some Labrador tea, willow, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: 60 cm.

Described By: D. Mellem and D. Haverkost on December 10, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-122
Fine	122+
sandy	
clay	

Location: 975 feet S and 670 feet W of the NE corner of Sec. 30, T.152N., R.26W. (NE1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 30, T.152N., R.26W.)

Vegetation: Scattered black spruce, tamarack, northern white cedar, and balsam

Location: 1100 feet S and 575 feet W of the NE corner of Sec. 8, T.152N., R.26W.

Vegetation: Lush understory consists mostly of grasses with some willow, alder,

Depth (cm)

61-91

91+

(SW1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 8, T.152N., B.26W.)

Described By: B. Sether and S. Nelson on April 3, 1980.

Laver

Hemic

Sapric

sandy

clay loam

Medium

vegetation: Scattered black spruce, tamarack, northern white cedar, and balsam fir; understory consists of bog birch, speckled alder, willow, Labrador tea, leatherleaf, bog rosemary, sedges, lingen berry, and swamp cinquefoll; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 20 cm.

and cattail.

Depth To Water Table: At surface.

Microrelief: 5 cm.

Depth To Water Table: Not recorded

Described By: T Malterer and N. Aaseng on December 10, 1979.

Layer	Depth (cm)
Hemic	0-152
Sapric	152-229
Loam	229+

Location: 1375 feet S and 775 feet W of the NE corner of Sec. 33, T.152N., R.26W. (NE1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 33, T.152N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent; understory consists mostly of Labrador tea with some leatherleaf, swamp laurel. bog rosemary. sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and N. Aaseng on December 10, 1979.

 Layer
 Depth (cm)

 Fibric
 0- 18

 Hemic
 18-313

 Silty
 313+

clay loam Location: 545 feet N and 1175 feet W of the SE corner of Sec. 33, T.152N., R.26W. (NW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 33, T.152N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent; understory consists mostly of leatherleaf with some Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: 60 cm.

Described By: D. Mellem and D. Haverkost on December 10, 1979.

Layer	Depth (cm)
Hemic	0-137
Silty	137+
clay	

Location: 1425 feet N and 1530 feet W of the SE corner of Sec. 20. T.153N., R.26W. (SE1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 20, T.153N., R.26W.)

Vegetation: Northern white cedar crown cover of about 85 percent with scattered balsam fir; ground cover consists of some feather and Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 25, 1976.

Layer	Depth (cm)
Hemic	0- 10
Fine	10+
2024	

Location: 355 feet S and 1875 feet E of the NW corner of Sec. 29, T.153N., R.26W. (SE1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 29, T.153N., R.26W.)

Vegetation: Cut over area: jush understory consists mostly of shrubs and grasses with some willows, paper birch, black spruce, balsam fir, hazel, and thistles.

Microrellef: Negligible.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 25, 1976.

Depth (cm) Layer Hemic 0- 30 Fine 30+ sand

Location: 100 feet N and 2275 feet W of the SE corner of Sec. 2, T.154N., R.26W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 2, T.154N., R.26W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; understory consists of some Labrador tea, horsetail, grasses, fern. willow, and speckled alder; ground cover consists of some Sphagnum and other mosses and cranberry.

Microrellef: 15 cm.

Sphagnum mosses. Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 6, 1976.

Layer	Depth (cm)
Sapric Sandy	0-46 46+
loam	40±

Location: 40 feet N and 1210 feet E of the SW corner of Sec. 4, T.154N., R.26W.

Vegetation: Black spruce crown cover of about 30 percent with scattered

tamarack; lush understory consists mostly of bog birch with Labrador tea.

leatherleaf, willow, and some grasses; ground cover consists mostly of

(SE1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 4, T.154N., R.26W.)

Location: 950 feet S and 1425 feet E of the NW corner of Sec. 31, T.153N., R.26W.

Vegetation: Northern white cedar crown cover of about 85 percent; lush understory

consists mostly of grasses with some balsam fir. fern, and Labrador tea;

Depth (cm)

0- 10

10- 51

51+

(SW1/4 of SW1/4 of NE1/4 of NW1/4, Sec. 31, T.153N., B.26W.)

ground cover consists of some false Solomon's seal.

Layer

Hemic

Sapric

Clav

(SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 3, T.154N., R.26W.) Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; understory consists of speckled alder with some bog birch and grasses; ground cover consists of some Sphagnum mosses.

Location: 75 feet N and 800 feet E of the SW corner of Sec. 3, T.154N., R.26W.

Microrelief: 45 cm.

Microrelief: Negligible.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 23, 1976.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 6, 1976.

Layer	Depth (cm)
Sapric	0-127
Loam	127+

Location: 50 feet N and 1675 feet W of the SE corner of Sec. 4, T.154N., R.26W. (SW1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 4, T.154N., R.26W)

Vegetation: Not recorded. Microrelief: Not recorded. Depth To Water Table: Not recorded.

Described By: R. Zarth on September 6, 1976.

Described By: R. Zarth on September 6, 1976. Layer Depth (cm) Peat 0-132 Mineral 132+ Depth (cm) soil

Location: 75 feet N and 525 feet F of the SW corner of Sec. 2, T.154N., R.26W. (SE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 2, T.154N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; lush understory consists mostly of grasses and speckled alder with some Labrador tea; ground cover consists of some Sphagnum and other mosses.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 6, 1976.

Layer	Depth (cm)
Hemic	0- 97
Sandy loam	97+
loam	

Location: 50 feet N and 1960 feet W of the SE corner of Sec. 3, T.154N., R.26W. (SW1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 3, T.154N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; lush understory consists mostly of leatherleaf with some grasses, Labrador tea, swamp laurel, and bog birch; ground cover consists of some cranberry and false Solomon's seal.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 6, 1976.

Layer	Depth (cm
Hemic Medium to	0-145 145+
coarse sand	

Location: 50 feet N and 1250 feet W of the SE corner of Sec. 5, T.154N., R.26W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 5, T.154N., R.26W.)

Vegetation: Consists mostly of grasses and willow with leatherleaf and bog birch.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 6, 1976.

Layer	Depth (cm)
Peat	0-213
Mineral	213+
soil	

Layer Hemic 0-183 Sandy 183 +loam

Location: 1120 feet S and 2450 feet E of the NW corner of Sec. 16, T.154N., R.26W. (SE1/4 of SE1/4 of NE1/4 of NW1/4, Sec. 16, T.154N., R.26W.)

Vegetation: Tamarack crown cover of about 90 percent with scattered black spruce; understory consists mostly of grasses with some bog rosemary and cattalls; ground cover consists of some Sphagnum mosses and pitcher plant. Microrellef: 22 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer Depth (cm) Hemic 0-305 305+ Fine sand

Location: 1200 feet N and 2050 feet W of the SE corner of Sec. 31, T.154N., R.26W. (NF1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 31, T.154N., R.26W.) Vegetation: Not recorded.

Microrelief: Not recorded

Depth To Water Table: Not recorded. Described By: Not recorded.

Laver Depth (cm) Peat 0-15 Mineral 15 +soil

Location: 75 feet N and 700 feet W of the SE corner of Sec. 31, T.154N., R.26W. (SE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 31, T.154N., R.26W.)

Vegetation: Not recorded.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Layer	Depth (cm
Peat Mineral	0-76 76+
soil	

Location: 1335 feet S and 60 feet E of the NW corner of Sec. 31, T.154N., R.26W. (Gov. Lot 2, Sec. 31, T.154N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack and northern white cedar; lush understory consists mostly of speckled alder with some bog birch.

Microrellef: Not recorded. Depth To Water Table: Not recorded. Described By: Not recorded.

> Layer Depth (cm) Hemic 0-84 Sandy 84+

Location; 2425 feet S and 265 feet E of the NW corner of Sec. 4, T.155N., R.26W. (SW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 4, T.155N., R.26W.)

Vegetation: Scattered black spruce and tamarack; understory consists of sedges with some Labrador tea, bog birch, willow, and fern; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellef: 60 cm.

Depth To Water Table: At surface.

Described By: R. Zarth and R. Fox on August 16, 1976.

Layer	Depth (cm)
Hemic	0- 58
Medium	58+

Location: 1300 feet N and 685 feet E of the SW corner of Sec. 4, T.155N., R.26W. (NW1/4 of NE1/4 of SW1/4 of SW1/4, Sec. 4, T.155N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered northern white cedar, tamarack, and paper birch; understory consists of some Labrador tea, speckled alder, sedges, grasses, and fern; ground cover consists of some Sphagnum and other mosses and false Solomon's seal.

Microrelief: 60 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on August 17, 1976.

Layer	Depth (cm)
Hemic	0-46
Sandy	46+
clay	
loam	

Location: 90 feet N and 1010 feet E of the SW corner of Sec. 4, T.155N., R.26W. (SE1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 4, T.155N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with a tamarack crown cover of about 35 percent; lush understory consists mostly of Labrador tea with bog birch and some bog rosemary, leatherleaf, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Zarth and R. Fox on August 17, 1976.

Layer	Depth (cm)
Hemic	0-221
Coarse	221+
eend	

Location: 1450 feet S and 1450 feet E of the NW corner of Sec. 9, T.155N., R.26W. (NW1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 9, T.155N., R.26W.)

Vegetation: Black spruce crown cover of about 85 percent; jush understory consists mostly of leatherleaf with some cotton grass, Labrador tea, bog rosemary, and swamp laurel; ground cover consists of some false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Zarth and R. Fox on August 17, 1976.

Layer	Depth (cm)
Hemic Clay Ioam	0-279 279+

Location: 2420 feet S and 1690 feet E of the NW corner of Sec. 9, T.155N., R.26W. (SE1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 9, T.155N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of leatherleaf with bog rosemary, swamp laurel, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on August 17, 1976.

Layer	Depth (cm)
Hemic	0-259
Coarse	259+
sand	

Location: 100 feet N and 2425 feet E of the SW corner of Sec. 9, T.155N., R.26W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 9, T.155N., R.26W.)

Vegetation: Black spruce crown cover of about 50 percent; understory consists of Labrador tea and leatherleaf with some bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth and R. Fox on August 16, 1976.

Hemic 0-173 Medium 173+ sand Location: 1055-feet S and 2375 feet W of the NE corner of Sec. 16, T.155N., R.26W. (SW1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 16, T.155N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent; lush understory consists of Labrador tea, cotton grass, and leatherleaf with some swamp laurel and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on August 16, 1976.

Layer	Depth (cm)
Hemic	0-173
Coarse sand	173+
to	
gravel	

Location: 1800 feet N and 1775 feet W of the SE corner of Sec. 16, T.155N., R.26W. (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 16, T.155N., R.26W.)

Vegetation: Black spruce crown cover of about 85 percent; lush understory consists mostly of sedges with some Labrador tea; ground cover consists of mosses with some Sohagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on August 16, 1976.

Layer	Depth (cm)
Hemic	0-66
Sapric	66-71
Sandy	71+
loam	

Location: 1475 feet S and 925 feet W of the NE corner of Sec. 21, T.155N., R.26W. (NW1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 21, T.155N., R.26W.)

Vegetation: Tamarack crown cover of about 85 percent with scattered black spruce and paper birch; lush understory consists mostly of Labrador tea with some grasses, balsam fir, and speckled alder; ground cover consists mostly of Sphagnum mosses.

Microrelief: 60 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on August 17, 1976.

Layer	Depth (cm)
Hemic	0-196
Sandy	196+
loam	

Location: 1075 feet N and 265 feet W of the SE corner of Sec. 21, T.155N., R.26W. (NW1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 21, T.155N., R.26W.)

Vegetation: Black spruce crown cover of about 85 percent with scattered tamarack; understory consists mostly of sedges with some Labrador tea, bog rosemary, and leatherleaf; ground cover consists of mosses with some Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on August 16, 1976.

Layer	Depth (cm)
Hemic	0-162
Sapric	162-165
Pebbly	165+
sandy	
loam	

Location: 75 feet N and 25 feet W of the SE corner of Sec. 21, T.155N., R.26W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 21, T.155N., R.26W.)

Vegetation: Tamarack crown cover of about 85 percent with scattered black spruce; lush understory consists mostly of speckled alder with leatherleaf and some bog birch, balasm fir, Labrador tea, and sedges; ground cover consists mostly of Sphagnum mosses.

Microrelief: 60 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on August 16, 1976.

Layer	Depth (cm
Hemic	0-86
Sapric	86-97
Coarse	97+
pebbly	
sand	

Location: 2150 feet S and 1360 feet W of the NE corner of Sec. 6, T.156N., R.26W. (SE1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 6, T.156N., R.26W.)

Vegetation: Black spruce crown cover of about 70 percent; understory consists of Labrador tea, leatherleaf, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 27, 1979.

Layer Depth (cm)

Hemic 0-213

Fine 213+
and medium
sand

Location: 2600 feet N and 500 feet W of the SE corner of Sec. 6, T.156N., R.26W. (NW1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 6, T.156N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of grasses and Labrador tea with some leatherleaf, swamp laurel, and speckled alder; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 27, 1979.

Layer	Depth (cm)
Hemic	0-165
Silty	165+
clay	

Location: 1810 feet S and 400 feet E of the NW corner of Sec. 8, T.156N., R.26W. (SE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 8, T.156N., R.26W.)

Vegetation: Scattered black spruce, northern white cedar, paper birch, and tamarack; lush understory consists mostly of grasses with some speckled alder; ground cover consists of some Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 27, 1979.

Layer	Depth (cm)
Hemic Sandy clay	0-90 90+

seal.

Microrellef: 15 cm.

Depth To Water Table: At surface:

Location: 1950 feet N and 475 feet W of the SE corner of Sec. 9, T.156N., R.26W. (NW1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 9, T.156N., R.26W.)

Vegetation: Northern white cedar crown cover of about 65 percent with about 30 percent crown cover of tamarack and scattered black spruce; understory consists of Labrador tea with some grasses, speckled alder, sedges, and cattail; ground cover consists of some feather and Sphagnum mosses, cranberry, and snowberry.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer Depth (cm) Sapric 0-61 Medium 61 +coarse sand

Location: 2375 feet S and 625 feet W of the NE corner of Sec. 19, T.156N., R.26W.

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly

of sedges with some Labrador tea, leatherleaf, swamp laurel, and cotton grass;

ground cover consists mostly of Sphagnum mosses with some false Solomon's

Depth (cm)

0- 60

60-335

335+

(SE1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 19, T.156N., R.26W.)

Described By: D. Meliem and D. Haverkost on September 26, 1979.

Layer

Fibric

Hemic

Sandy

loam

Location: 2210 feet N and 1600 feet F of the SW corner of Sec. 17 T 156N R 26W (SW1/4 of NE1/4 of NE1/4 of SW1/4, Sec. 17, T.156N., R.26W.)

Vegetation: Scattered northern white cedar, black spruce, and tamarack; lush understory consists mostly of grasses with some balsam fir, willow, and alder; ground cover consists of some mosses.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 27, 1979.

Layer	Depth (cr
Sapric Sandy	0-15 15+
loam	

Location: 2425 feet S and 25 feet W of the NE corner of Sec. 19, T.156N., R.26W. (SE1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 19, T.156N., R.26W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem and D. Haverkost on September 26, 1979.

Layer Depth (cm) 0- 90 Fibric 90-273 Hemic Sandv 273 +loam

Location: 750 feet S and 1975 feet E of the NW corner of Sec. 19, T.156N., R.26W. (NW1/4 of SE1/4 of NE1/4 of NW1/4, Sec. 19, T.156N., R.26W.)

Location: 550 feet N and 115 feet F of the SW corner of Sec. 18, T.156N., R.26W.

Vegetation: Scattered black spruce: understory consists of some Labrador tea.

Described By: D. Mellem and D. Haverkost on September 26, 1979.

Laver

Fibric

Hemic

Coarse

sand

leatherleaf, and sedges; ground cover consists mostly of Sphagnum mosses.

Depth (cm)

0-120

120-304

304+

(Gov. Lot 4, Sec. 18, T.156N., R.26W.)

Depth To Water Table: Not recorded.

Microrellef: Not recorded.

Vegetation: Scattered black spruce; understory consists of leatherleaf with some sedges. Labrador tea, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 15 cm

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 26, 1979.

Layer	Depth (cm
Fibric	0-150
Hemic	150-304
Fine	304+
sand	

Location: 2110 feet S and 125 feet E of the NW corner of Sec. 19, T.156N., R.26W (Gov. Lot 2, Sec. 19, T.156N., R.26W.)

Vegetation: Scattered black spruce; understory consists of leatherleaf with some Labrador tea and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 26, 1979.

Layer Depth (cm) Fibric 0- 90 Hemic 90-354 Sapric 354-364 Silt 364+ with calcareous pebbles

Location: 1775 feet N and 125 feet E of the SW corner of Sec. 19, T.156N., R.26W. (Gov. Lot 3, Sec. 19, T.156N., R.26W.)

Vegetation: Scattered black spruce: understory consists of sedges with some swamp laurel, Labrador tea, leatherleaf, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 26, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-312
Sapric	312-317
Silt	317+
and	
fine	
eand	

Location: 2400 feet S and 765 feet E of the NE corner of Sec. 20, T.156N., R.26W. (SW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 20, T.156N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some bog birch, willow, Labrador tea, bog rosemary, leatherleaf, and ferns: ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 26, 1979.

Layer	Depth (cm)
Hemic Sandy Ioam	0-211 211+

Location: 2125 feet N and 675 feet W of the SE corner of Sec. 28, T.156N., R.26W. (SW1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 28, T.156N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with about 30 percent crown cover of tamarack and scattered paper birch; understory consists of some Labrador tea, leatherleaf, swamp laurel, bog rosemary, grasses and speckled alder; ground cover consists of some Sphagnum and feather mosses.

Microrellef: 25 cm.

Depth To Water Table: 15 cm.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Location: 2625 feet S and 1125 feet E of the NW corner of Sec. 1, T.157N., R.26W.

Vegetation: Black spruce crown cover of about 40 percent; understory consists of

Labrador tea with some leatherleaf, swamp laurel, and sedges; ground cover

consists of Sphagnum and other mosses with some lingen berry and cranberry.

Depth (cm)

0-183

183+

(SE1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 1, T.157N., R.26W.)

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer

Clay

Location: 950 feet S and 75 feet E of the NW corner of Sec. 30, T.156N., R.26W. (Gov. Lot 1, Sec. 30, T.156N., R.26W.)

Vegetation: Scattered black spruce; sparse understory consists of some Labrador tea, sedges, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 26, 1979.

Layer	Depth (cm
Fibric	0- 60
Hemic	60-195
Sandy	195+
clay	

Location: 2400 feet N and 1325 feet W of the SE corner of Sec. 3, T.157N., R.26W. (NW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 3, T.157N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some Labrador tea, alder, leatherleaf, and sedges; ground cover consists mostly of Sphagnum mosses with some feather mosses.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 15, 1979.

 Layer
 Depth (cm)

 Hemic
 0-152

 Silt
 152+

Location: 1250 feet S and 400 feet E of the NW corner of Sec. 15, T.157N., R.26W.

Location: 2450 feet S and 120 feet E of the NW corner of Sec. 30, T.156N., R.26W.

Vegetation: Scattered black spruce and tamarack; understory consists of grasses

Described By: D. Mellem and D. Haverkost on September 26, 1979.

Laver

Hemic

Sandy

clay

with some alder, Labrador tea, and balsam fir; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Depth (cm)

0-100

100 +

(Gov. Lot 2, Sec. 30, T.156N., R.26W.)

Depth To Water Table: At surface.

Microrelief: 20 cm.

(SE1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 15, T.157N., R.26W.)

Vegetation: Black spruce crown cover of about 30 percent with scattered tamarack, paper birch, and aspen; understory consists of Labrador tea with some cotton grass, swamp laurel, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, lingen berry, and

blueberry.
Microrelief: 20 cm.

Depth To Water Table: Not visible.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer	Depth (cm
Sapric	0- 91
Clay	91+
loam	

Location: 725 feet S and 365 feet E of the NW corner of Sec. 16, T.157N., R.26W. (NE1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 16, T.157N., R.26W.)

Vegetation: Scattered black spruce and northern white cedar; sparse understory consists of some grasses, willow, and Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrellef: 15 cm.

Microrellef: 35 cm

Depth To Water Table: Not recorded.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Sapric 0-81
Medium 81+
sand

Location: 550 feet S and 210 feet E of the NW corner of Sec. 19, T.157N., R.26W. (SW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 19, T.157N., R.26W.)

Vegetation: Scattered black spruce; sparse understory consists of some leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with false Solomon's seal and some other mosses, cranberry, and pitcher plant.

Microrelief: 60 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 4, 1979.

Layer	Depth (cm)
Fibric	0-165
Hemic	165-272
Sapric	272-287
Sandy	287+
clav	

Location: 675 feet N and 285 feet E of the SW corner of Sec. 19, T.157N., R.26W. (SW1/4 of NW1/4 of SW1/4, Sec. 19, T.157N., R.26W.)

Vegetation: Scattered black spruce; sparse understory consists of some sedges, leatherleaf, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses and some false Solomon's seal.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 4, 1979.

Layer	Depth (cm)
Fibric	0-211
Hemic	211-302
Sandy clay	302+

Location: 500 feet N and 410 feet W of the SE corner of Sec. 27, T.157N., R.26W. (NW1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 27, T.157N., R.26W.)

Vegetation: Scattered black spruce, paper birch, and northern white cedar; sparse understory consists of some willow, poplar, ferns, cattails, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 7 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

 Layer
 Depth (cm)

 Hemic
 0-107

 Clay
 107+

Location: 700 feet S and 2200 feet W of the NE corner of Sec. 30, T.157N., R.26W. (NE1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 30, T.157N., R.26W.)

Vegetation: Tamarack crown cover of about 25 percent with scattered black spruce; understory consists mostly of sedges with some bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 3, 1979.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-335
Sapric	335-348
Sandy	348+
clay	
with	
calcareous	
pebbles	

Location: 1900 feet N and 215 feet E of the SW corner of Sec. 30, T.157N., R.26W. (Gov. Lot 3, Sec. 30, T.157N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some bog birch, bog rosemary, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 4, 1979.

Layer	Depth (cm)
Fibric	0- 90
Hemic	90-212
Sandy	212+
clay	

Location: 650 feet S and 200 feet E of the NW corner of Sec. 31, T.157N., R.26W. (Gov. Lot 1, Sec. 31, T.157N., R.26W.)

Vegetation: Tamarack crown cover of about 35 percent; lush understory consists mostly of sedges with some bog birch, willow, bog rossemary, Labrador tea, and cattalis; ground cover consists of some Sphagnum mosses.

Microrellef: 10 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 4, 1979.

Layer	Depth (cm
Hemic	0-210
Sandy	210+
clay	

Location: 1765 feet N and 1020 feet W of the SE corner of Sec. 2, T.158N., R.26W. (NW1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 2, T.158N., R.26W.)

Vegetation: Scattered tamarack and northern white cedar; sparse understory consists of some speckled alder, reeds, leatherleaf, sedges, and ferns; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Fibric	0-213
Hemic	213-274
Sapric	274-290
Clay	290+
loam	

Location: 2500 feet N and 125 feet E of the SW corner of Sec. 3, T.158N., R.26W. (NW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 3, T.158N., R.26W.)

Vegetation: Scattered tamarack; sparse understory consists of some sedges, reeds, leatherleaf, and bog rosemary; ground cover consists of some Sphagnum mosses.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Fibric	0-229
Hemic	229-274
Sapric	274-305
Clay	305+
loam	

Location: 2175 feet N and 50 feet W of the SE corner of Sec. 3, T.158N., R.26W. (SE1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 3, T.158N., R.26W.)

Vegetation: Scattered tamarack; sparse understory consists of some sedges, tamarack, speckled alder, and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: B. Sether and D. Haverkost on March 13, 1980.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-335
Sapric	335-350
Clay	350+
loam	

Location: 985 feet N and 1190 feet W of the SE corner of Sec. 5, T.158N., R.26W. (SW1/4 of NW1/4 of SE1/4 of SE1/4, Sec. 5, T.158N., R.26W.)

Vegetation: Scattered tamarack; lush understory consists mostly of sedges with some bog rosemary, bog birch, leatherleaf, horsetails, and buckbean; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 10 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm
Hemic	0-152
Fibric	152-244
Hemic	244-396
Clay	396+
loam	

Location: 850 feet S and 590 feet W of the NE corner of Sec. 6, T.158N., R.26W. (Gov. Lot 1, Sec. 6, T.158N., R.26W.)

Vegetation: Scattered tamarack; understory consists of sedges with some bog rosemary, leatherleaf, cotton grass, and bog birch; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 10 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

.ayer	Depth (cm)
lemic Clay cam	0-180 180+

Location: 1825 feet N and 175 feet E of the SW corner of Sec. 6, T.158N., R.26W. (Gov. Lot 6, Sec. 6, T.158N., R.26W.)

Vegetation: Scattered tamarack; sparse understory consists of some sedges, reeds, willow, leatherleaf, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Fibric	0-168
Hemic	168-350
Silty	350+
clay	

Location: 25 feet S and 1150 feet W of the NE corner of Sec. 7, T.158N., R.26W. (NW1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 7, T.158N., R.26W.)

Vegetation: Understory consists of some grasses, tamarack, speckled alder, leatherlear, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: H. Mooers, B. Sether, J. Dahl, and D. Haverkost on March 6, 1980.

Layer	Depth (cm
Fibric	0-250
Hemic	250-330
Sapric	330-345
Clay	345+
loam	

Location: 1925 feet N and 1510 feet W of the SE corner of Sec. 7, T.158N., R.26W. (NE1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 7, T.158N., R.26W.)

Vegetation: Scattered tamarack; understory consists of sedges with some leatherleaf, bog rosemary, cotton grass, and bog birch; ground cover consists mostly of Sphagnum mosses.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm
Fibric	0-274
Hemic	274-305
Clay	305+

Location: 60 feet S and 1550 feet E of the NW corner of Sec. 9, T.158N., R.26W. (NW1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 9, T.158N., R.26W.)

Vegetation: Lush understory consists mostly of grasses with some tamarack, sedges, bog rosemary, leatherleaf, and reeds.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: H. Mooers and D. Haverkost on March 11, 1980.

Layer	Depth (cm)
Hemic Silty clay	0-345 345+

Location: 450 feet S and 950 feet W of the NE corner of Sec. 10, T.158N., R.26W. (SE1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 10, T.158N., R.26W.)

Vegetation: Scattered tamarack; sparse understory consists of some sedges, reeds, leatherleaf, bog rosemary, and speckled alder; ground cover consists of some Sphaanum mosses.

Microrelief: Negligible.

Depth To Water Table: 15 cm. above surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm
Fibric	0-213
Hemic	213-274
Clay	274+
loam	

Location: 25 feet S and 2475 feet W of the NE corner of Sec. 10, T.158N., R.26W. (NW1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 10, T.158N., R.26W.)

Vegetation: Understory consists of some grasses, tamarack, leatherleaf, and bog rosemary; ground cover was not recorded.

Microrelief: 18 cm.

Depth To Water Table: Not recorded.

Described By: H. Mooers and D. Haverkost on March 11, 1980.

Layer	Depth (cm)
Hemic	0-260
Sapric	260-285
Clay	285+
loam	
with	
pebbles	

Location: 1810 feet N and 275 feet E of the SW corner of Sec. 10, T.158N., R.26W. (NW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 10, T.158N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 45 cm.

Depth To Water Table: 15 cm. above surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Fibric	0-183
Hemic	183-381
Sapric	381-411
Silty	411+
clay	

Location: 50 feet S and 2160 feet E of the NW corner of Sec. 11, T.158N., R.26W. (NW1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 11, T.158N., R.26W.)

Vegetation: Sparse understory consists of some tamarack, speckled alder, swamp laurel, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses

Microrelief: 15 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and D. Haverkost on March 12, 1980.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-244
Fibric	244-305
Hemic	305-350
Clay	350+
loam	

Location: 1900 feet N and 1975 feet W of the SE corner of Sec. 11, T.158N., R.26W. (NE1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 11, T.158N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent; understory consists of Labrador tea with some sedges, leatherleaf, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, snowberry, lingen berry, and false Solomon's seal.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm)
Hemic Clav	0-315 315+
loam	3137

Location: 1775 feet N and 20 feet W of the SE corner of Sec. 12, T.158N., R.26W. (NE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 12, T.158N., R.26W.)

Vegetation: Scattered black spruce; sparse understory consists of leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, craphery, and snowberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

 Layer
 Depth (cm)

 Fibric
 0-138

 Hemic
 138-259

 Sapric
 259-274

 Sility
 274+

 clay

Location: 850 feet S and 75 feet E of the NW corner of Sec. 19, T.158N., R.26W.

Vegetation: Scattered black spruce; understory consists of some sedges.

leatherleaf, swamp laurel, Labrador tea, and cotton grass; ground cover con-

Depth (cm)

0-305

305-396

396+

sists mostly of Sphagnum mosses with some cranberry and pitcher plant.

(SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 19, T.158N., R.26W.)

Described By: N. Aaseng and R. Graber on November 13, 1979.

Layer

Fibric

Hemic

Silty

clay

Location: 2275 feet S and 1015 feet W of the NE corner of Sec. 15, T.158N., R.26W. (SW1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 15, T.158N., R.26W.)

Vegetation: Scattered black spruce, tamarack, and northern white cedar; understory consists of some bog birch, willow, leatherleaf, bog rosemary, sedges, and five finger cinquefoil; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm)
Hemic	0-173
Clay	173+

Location: 1800 feet S and 320 feet E of the NW corner of Sec. 21, T.158N., R.26W. (SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 21, T.158N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, swamp laurel, bog rosemary, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Fibric 0-213 Hemic 213-305 Silty 305+ clay Location: 350 feet S and 1850 feet E of the NW corner of Sec. 22, T.158N., R.26W. (NE1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 22, T.158N., R.26W.)

Location: 2375 feet S and 1205 feet E of the NW corner of Sec. 15, T.158N., R.26W.

Vegetation: Scattered black spruce and tamarack; understory consists of sedges

mostly of Sphagnum mosses with some cranberry and pitcher plant.

with some leatherleaf, swamp laurel, and bog rosemary; ground cover consists

Depth (cm)

0-375

375+

(SE1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 15, T.158N., R.26W.)

Described By: D. Olson and N. Aaseng on November 14, 1979.

Laver

Hemic

Clay

loam

Vegetation: Scattered tamarack and black spruce; understory consists of bog rosemary with some sedges, cattails, Phragmites, leatherleaf, and bog birch; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: Not recorded.

Microrellef: 30 cm

Depth To Water Table: At surface.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-229
Clay	229+

Location: 125 feet N and 100 feet E of the SW corner of Sec. 26, T.158N., R.26W. (SW1/4 of SW1/4 of SW1/4, Sec. 26, T.158N., R.26W.)

Vegetation: Black spruce crown cover of about 75 percent; sparse understory consists of some Labrador tea and sedges; ground cover consists mostly of mosses with some Sphagnum mosses, lingen berry, blueberry, and snowberry.

Microrelief: 40 cm.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Depth To Water Table: 2 cm.

Described By: D. Olson and N. Aaseng on November 15, 1979.

 Layer
 Depth (cm)

 Hemic
 0-229

 Sapric
 229-242

 Clay
 242+

Location: 25 feet N and 110 feet W of the SE corner of Sec. 26, T.158N., R.26W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 26, T.158N., R.26W.)

Vegetation: Black spruce crown cover of about 35 percent; sparse understory consists of some Labrador tea and sedges; ground cover consists mostly of Sphagnum and other mosses with some lingen berry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on October 31, 1979.

Layer	Depth (cm
Hemic	0- 91
Sapric	91-118
Clav	118+

Location: 1825 feet S and 75 feet E of the NW corner of Sec. 1, T.159N., R.26W. (SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 1, T.159N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent; understory consists of Labrador tea and grasses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem on November 16, 1979.

Layer	Depth (cm)
Hemic	0-229
Sandy clay	229+

Location: 1450 feet N and 225 feet E of the SW corner of Sec. 3, T.159N., R.26W. (SW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 3, T.159N., R.26W.)

Vegetation: Aspen crown cover of about 80 percent; understory consists of grasses with some speckled alder and willow.

Microrelief: 20 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem on November 16, 1979.

Layer Depth (cm) Hemic 183+ Clay

Location: 1910 feet N and 2010 feet E of the SW corner of Sec. 18, T.159N., R.26W.

Vegetation: Scattered black spruce and tamarack: sparse understory consists of

some leatherleaf, swamp laurel, bog rosemary, sedges, and cotton grass;

ground cover consists mostly of Sphagnum mosses with some other mosses,

Depth (cm)

0-138

138-259

259-305 305+

(NW1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 18, T.159N., R.26W.)

Described By: R. Graber and D. Haverkost on November 14, 1979.

Laver Fibric

Hemic

Sapric

Clav loam with pebbles Location: 1675 feet N and 110 feet W of the SE corner of Sec. 5, T.159N., R.26W. (NE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 5, T.159N., R.26W.)

Venetation: Black spruce crown cover of about 35 percent with scattered aspen and tamarack; understory consists of grasses and some Labrador tea; ground cover consists of some mosses.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem on November 16, 1979.

Layer	Depth (cm)
Hemic	0-168
Silt	168+

Location: 1725 feet N and 1450 feet E of the SW corner of Sec. 20, T.159N., R.26W. (NW1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 20, T.159N., R.26W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Fibric	0-137
Hemic	137-411
Clay	411+

Location: 2075 feet N and 60 feet W of the SE corner of Sec. 23, T.159N., R.26W. (SE1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 23, T.159N., R.26W.)

Location: 1225 feet N and 1950 feet W of the SE corner of Sec. 7, T.159N., R.26W.

Vegetation: Scattered black spruce and tamarack; Jush understory consists mostly

of Labrador tea. leatherleaf, and sedges with some swamp laurel and cotton

Depth (cm)

0- 91

91-457

457+

(NW1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 7, T.159N., R.26W.)

grass; ground cover consists of Sphagnum mosses.

Laver

Fibric

Hemic

Sapric

Bottom unknown

Described By: R. Graber and D. Haverkost on November 14, 1979.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Vegetation: Black spruce crown cover of about 30 percent; lush understory consists mostly of leatherleaf with some Labrador tea, bog rosemary, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses. Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 21, 1979.

Layer	Depth (cm)
Peat Mineral soil	0-244 244+

Location: 2100 feet N and 1300 feet W of the SE corner of Sec. 23, T.159N., R.26W. (SW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 23, T.159N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered cedar; lush understory consists mostly of Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

cranberry, and pitcher plant.

Depth To Water Table: At surface.

Microrelief: 30 cm.

Described By: M. Domeier on July 21, 1976.

Layer	Depth (cm)
Sapric Mineral soil	0- 66 66+

Location: 2075 feet N and 975 feet W of the SE corner of Sec. 23, T.159N., R.26W. (SE1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 23, T.159N., R.26W.)

Vegetation: Black spruce crown cover of about 40 percent; lush understory consists mostly of Labrador tea with leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 21, 1976.

Layer	Depth (cm)
Fibric	0- 35
Sapric	35-157
Sandy	157+
clay	
loam	

Location: 750 feet S and 975 feet W of the NE corner of Sec. 24, T.159N., R.26W. (NW1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 24, T.159N., R.26W.)

Vegetation: Black spruce crown cover of about 75 percent; understory consists of some swamp laurel, sedges, cotton grass, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry. Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm)
Fibric	0-106
Hemic	106-455
Silty	455+
clay	

Location: 2620 feet S and 1875 feet E of the NW corner of Sec. 24, T.159N., R.26W. (SE1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 24, T.159N., R.26W.)

Vegetation: Scattered tamarack; lush understory consists mostly of grasses with some bog rosemary, swamp laurel, and bog birch.

Microrellef: 45 cm.

Depth To Water Table: 10 cm. above surface.

Described By: M. Domeier on July 21, 1976.

Layer Depth (cm)
Hemic 0-244
Mineral 244+
soil

Location: 2375 feet N and 1440 feet E of the SW corner of Sec. 24, T.159N., R.26W. (NW1/4 of NW1/4 of NE1/4 of SW1/4, Sec. 24, T.159N., R.26W.)

Vegetation: Scattered tamarack; lush understory consists mostly of cotton grass with bog birch and some bog rosemary; ground cover consists of Sphagnum masses.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 21, 1976.

Layer	Depth (cm
Hemic	0-244
Mineral	244+
soil	

Location: 2060 feet N and 1030 feet E of the SW corner of Sec. 24, T.159N., R.26W. (SE1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 24, T.159N., R.26W.)

Vegetation: Black spruce crown cover of about 65 percent; lush understory consists mostly of Labrador tea with some leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 21, 1976.

Layer	Depth (cm)
Hemic	0-183
Mineral	183+
enil	

Location: 350 feet N and 600 feet W of the SE corner of Sec. 24, T.159N., R.26W. (NE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 24, T.159N., R.26W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of grasses and leatherleaf with speckled alder and some bog rosemary, sedges, and cattallis; ground cover consists of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-183
Fibric	183-274
Sapric	274-442
Clay	442+

Location: 675 feet S and 2050 feet E of the NW corner of Sec. 25, T.159N., R.26W. (NW1/4 of SE1/4 of NE1/4 of NW1/4, Sec. 25, T.159N., R.26W.)

Vegetation: Scattered black spruce; understory consists of sedges with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry and pitcher plant.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm)
Fibric Hemic Fibric Hemic Bottom unknown	0- 91 91-366 366-457 457+

Location: 2100 feet N and 25 feet E of the SW corner of Sec. 25, T.159N., R.26W. (SW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 25, T.159N., R.26W.)

Vegetation: Scattered black spruce; lush understory consists mostly of leatherleaf, swamp laurel, bog rosemary, and sedges with Labrador tea and some cotton grass; ground cover consists of some Sphagnum and feather mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-396
Sapric	396-442
Silty	442+
clay	

Location: 1925 feet N and 790 feet W of the SE corner of Sec. 25, T.159N., R.26W. (NE1/4 of SW1/4 of NE1/4 of SE1/4, Sec. 25, T.159N., R.26W.)

Vegetation: Black spruce crown cover of about 60 percent with about 30 percent crown cover of northern white cedar with scattered tamarack; lush understory consists mostly of bog rosemary, grasses, and speckled alder with some Labrador tea, leatherleaf, horsetail, and ferns; ground cover consists mostly of feather and Sphaanum mosses with some pitcher plant.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-366
Silty	366+
clay	

Location: 250 feet S and 1500 feet E of the NW corner of Sec. 28, T.159N., R.26W. (NW1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 28, T.159N., R.26W.)

Vegetation: Scattered black spruce and tamarack: understory consists of sedges with some bog birch, Labrador tea. bog rosemary, and grasses: ground cover consists of some mosses and bitcher plant.

Microrelief: 5 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on November 14. 1979.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-213
Fibric	213-427
Sapric	427-457
Silt	457+

Location: 1910 feet N and 75 feet E of the SW corner of Sec. 34, T.159N., R.26W. (NW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 34, T.159N., R.26W.)

Vegetation: Scattered black spruce; lush understory consists mostly of leatherleaf and bog rosemary with grasses and some speckled alder and reeds: ground cover consists mostly of mosses including some Sphagnum mosses and pitcher plant.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)
Hemic	0-183
Fibric	183-274
Silty	274+
clav	

Location: 1025 feet S and 350 feet E of the NW corner of Sec. 35, T.159N., R.26W. (SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 35, T.159N., R.26W.)

Vegetation: Scattered black spruce; understory consists of some leatherleaf, sedges, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

Layer	Depth (cm)
Hemic Clay to silty clay	0-340 340+

Location: 1020 feet N and 125 feet E of the SW corner of Sec. 33, T.160N., R.26W. (NW1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 33, T.160N., R.26W.)

Vegetation: Scattered aspen; lush understory consists mostly of grasses and speckled alder.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: D. Mellem on November 16, 1979.

cm)
3
3

Location: 1750 feet S and 260 feet E of the NW corner of Sec. 1, T.64N., R.27W. (SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 1, T.64N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent; sparse understory consists of some grasses, Labrador tea, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some feather mosses, cranberry, and false Solomon's seal.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and S. Nelson on April 8, 1980.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-183
Sand	183+

Location: 1420 feet N and 975 feet E of the SW corner of Sec. 2, T.64N., R.27W. (SW1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 2, T.64N., R.27W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack and paper birch; sparse understory consists of some black spruce and Labrador tea; ground cover consists mostly of Sphagnum mosses with some cranberry and snowberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: B. Sether and D. Haverkost on April 8, 1980.

Layer	Depth (cm
Hemic	0-122
Sapric	122-152
Silt	152+
loam	
with	
pebbles	

Location: 2200 feet S and 500 feet E of the NW corner of Sec. 34, T.64N., R.27W. (Gov. Lot 2, Sec. 34, T.64N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some sedges, cotton grass, Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and N. Aaseng on December 10, 1979.

Layer	Depth (cm)
Fibric Hemic	0- 61 61-137
Sapric Sandy Ioam	137-145 145-150
Medium fine sand	150+

Location: 50 feet N and 2500 feet E of the SW corner of Sec. 36, T.64N., R.27W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 36, T.64N., R.27W.)

Vegetation: Scattered black spruce and tamarack; understory consists of sedges and leatherleaf with some alder, grasses, and Labrador tea; ground cover was not observed due to ice cover on surface.

Microrelief: 5 cm.

Depth To Water Table: 10 cm. above surface.

Described By: H. Mooers and B. Sether on April 10, 1980.

Layer	Depth (cm)
Hemic	0-152
Sapric	152-183
Sand	183+

Location: 2600 feet S and 1240 feet E of the NW corner of Sec. 1, T.65N., R.27W. (SE1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 1, T.65N., R.27W.)

Vegetation: Tamarack crown cover of about 60 percent with scattered black spruce; sparse understory consists of grasses with some swamp laurel and leatherleaf; ground cover consists of Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)		
Hemic Medium sand	0-244 244+		

Location: 50 feet N and 300 feet W of the SE corner of Sec. 14, T.65N., R.27W. (Gov. Lot 4, Sec. 14, T.65N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of leatherleaf with some grasses, black spruce, tamarack, Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some feather mosses and false Solomon's seal.

Microrelief: 40 cm.

Depth To Water Table: 5 cm.

Described By: H. Mooers and S. Nelson on April 8, 1980.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-213
Loamy	213+
sand	
with	
gravel	

Location: 1960 feet N and 1800 feet W of the SE corner of Sec. 23, T.65N., R.27W. (Gov. Lot 3, Sec. 23, T.65N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of black spruce with some Labrador tea, leatherleaf, swamp laurel, bog rosemary, and grasses; ground cover consists mostly of Sphagnum mosses with some feather mosses, cranberry, and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: 30 cm.

Described By: B. Sether and D. Haverkost on April 8, 1980.

_ayer	Depth (cm
Hemic Fine and medium sand	0-152 152+

Location: 2325 feet N and 650 feet W of the SE corner of Sec. 24, T.65N., R.27W. (Gov. Lot 3, Sec. 24, T.65N., R.27W.)

Vegetation: Scattered black spruce and paper birch; understory consists of alder with some grasses; ground cover consists mostly of Sphagnum mosses with some other mosses and snowberry.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: B. Sether and D. Haverkost on April 8, 1980.

Laver Depth (cm) 0-122 Sapric Sandy 122+ loam

Location: 250 feet S and 880 feet W of the NE corner of Sec. 2, T.67N., R.27W. (Gov. Lot 1, Sec. 2, T.67N., R.27W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack; sparse understory consists of some sedges, Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 23 cm.

Microrellef: 25 cm.

Depth To Water Table: Not recorded.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer	Depth (cm)	
Fibric	0-122	
Hemic	122-167	
Sandy clay	167+	

Location: 1590 feet S and 1910 feet E of the NW corner of Sec. 13, T.68N., R.27W. (NE1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 13, T.68N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some poplar, Labrador tea, leatherleaf, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses with some cranberry and snowberry.

Microrellef: 15 cm.

Layer Depth (cm) 0-45 Hemic Sapric 45-89 Clay 89+

Location: 1600 feet S and 2590 feet W of the NE corner of Sec. 12, T.68N., R.27W.

Vegetation: Black spruce crown cover of about 75 percent; understory consists of

leatherleaf; ground cover consists mostly of Sphagnum mosses with some

(NW1/4 of NW1/4 of SW1/4 of NE1/4, Sec. 12, T.68N., R.27W.)

other mosses, lingen berry, blueberry, snowberry, and cranberry.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Location: 2000 feet S and 600 feet E of the NW corner of Sec. 25, T.65N., R.27W. (NE1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 25, T.65N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent; understory consists of Labrador tea with some grasses, black spruce, and swamp laurel; ground cover consists of Sphagnum mosses with some other mosses, bilberry, cranberry, false Solomon's seal, and lichens.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: H. Mooers and S. Nelson on April 8, 1980.

Depth (cm)
0- 61
61-198
198+

Location: 640 feet N and 2010 feet W of the SE corner of Sec. 25, T.67N., R.27W. (SE1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 25, T.67N., R.27W.)

Vegetation: Scattered black spruce and poplar; understory consists of some Labrador tea, leatherleaf, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and snowberry.

Depth To Water Table: Not visible.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)
Hemic	0-183
Medium to	183+
fine	
sand	

Location: 2600 feet S and 2290 feet W of the NE corner of Sec. 26, T.67N., R.27W.

(Gov. Lot 2, Sec. 26, T.67N., R.27W.) Vegetation: Clear cut area with scattered black spruce and aspen surrounded by

Location: 2000 feet S and 650 feet E of the NW corner of Sec. 26, T.65N., R.27W.

Vegetation: Black spruce crown cover of about 85 percent with scattered

mosses, cranberry, snowberry, bilberry, and false Solomon's seal.

tamarack; understory consists of some leatherleaf, swamp laurel, grasses, and sedges; ground cover consists mostly of Sphagnum mosses with some feather

Depth (cm)

0- 91

91-244

244+

(Gov. Lot 2, Sec. 26, T.65N., R.27W.)

Described By: H. Mooers and S. Nelson on April 8, 1980.

Layer

Fibric

Hemic

Sand

black spruce crown cover; understory consists of Labrador tea with some swamp laurel, speckled aider, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Microrellef: 30 cm.

Depth To Water Table: 15 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and P. Guertin on November 30, 1979.

Layer	Depth (cm)	
Hemic	0-244	
Sand	244+	

Location: 270 feet N and 200 feet E of the SW corner of Sec. 26, T.68N., R.27W. (Gov. Lot 4, Sec. 26, T.68N., R.27W.)

Vegetation: Black spruce crown cover of about 75 percent; sparse understory consists of some Labrador tea; ground cover consists mostly of mosses including some Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: R. Graber and D. Haverkost on November 30, 1979.

Layer	Depth (cm)		
Hemic	0-152		
Sandy clay	152+		

Depth 10	wateri	able: N	Ot VISID	ie.			
Described	By: R.	Graber	and D.	Haverkost	on November	15,	1979.

Layer	Depth (cm	
Hemic	0-69	
Coarse	69+	
sand		

Location: 50 feet N and 1100 feet W of the SE corner of Sec. 36, T.68N., R.27W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 36, T.68N., R.27W.)

Vegetation: Black spruce crown cover or about 95 percent; sparse understory consists of leatherleaf with some Labrador tea and swamp laurel; ground cover consists of Sphagnum mosses.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 29, 1979.

Layer	Depth (cm	
Fibric	0-106	
Sapric	106-122	
Silty	122+	
clav		

Location: 2430 feet S and 160 feet W of the NE corner of Sec. 2, T.69N., R.27W. (Gov. Lot 2, Sec. 2, T.69N., R.27W.)

Vegetation: Scattered black spruce, poplar, tamarack, and paper birch; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses with some snow-berry.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 15, 1979.

Layer	Depth (cm)
Hemic	0-76
Sapric	76-91
Fine	91+
sand	

Location: 1810 feet N and 480 feet W of the SE corner of Sec. 11, T.69N., R.27W. (Gov. Lot 3, Sec. 11, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; lush understory consists mostly of grasses with leatherleaf and some swamp laurel and Labrador tea; ground cover consists of Sphagnum mosses with some cranberry.

Microrellef: 25 cm.

Depth To Water Table: 10 cm.

Described By: B. Wakanabo on August 25, 1977.

Layer	Depth (cm)
Hemic	0-223
Muck	223-232
Silty	232+
clav	

Location: 2000 feet S and 840 feet E of the NW corner of Sec. 12, T.69N., R.27W. (NW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 12, T.69N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some sedges, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry, false Solomon's seal, and other mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 15, 1979.

Layer	Depth (cm)
Hemic	0-208
Clay	208+

Location: 10 feet N and 2030 feet W of the SE corner of Sec. 12, T.69N., R.27W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 12, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; understory consists of Labrador tea with some grasses, bog rosemary, leatherleaf, and swamp laurel; ground cover consists mostly of cranberry with Sphagnum mosses.

Microrellef: 35 cm.

Depth To Water Table: 20 cm.

Described By: B. Wakanabo on August 25, 1977.

Layer	Depth (cm)
Fibric	0- 25
Hemic	25-178
Loam	178+

Location: 20 feet S and 20 feet W of the NE corner of Sec. 13, T.69N., R.27W. (NE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 13, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent with scattered tamarack; understory consists of leatherleaf with some swamp laurel, grasses, and bog birch; ground cover consists of mosses with some cranberry, blueberry, false Solomon's seal, and Sphagnum mosses.

Microrelief: 20 cm.

Depth To Water Table: Below 45 cm.

Described By: B. Wakanabo on August 25, 1977.

Layer	Depth (c
Fibric	0- 16
Sapric	16- 31
Sandy	31+
loam	

Location: 1500 feet S and 460 feet W of the NE corner of Sec. 14, T.69N., R.27W. (Gov. Lot 2, Sec. 14, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent; lush understory consists mostly of Labrador tea and leatherleaf with grasses and some bog rosemary and swamp laurel; ground cover consists mostly of Sphagnum mosses with cranberry.

Microrellef: 40 cm.

Depth To Water Table: 15 cm.

Described By: B. Wakanabo on August 25, 1977.

Layer	Depth (cm)
Hemic	0- 18
Fibric	18- 30
Hemic	30-337
Muck	337-344
Silty	344+
clay	
loam	

Location: 1700 feet S and 1080 feet W of the NE corner of Sec. 23, T.69N., R.27W. (Gov. Lot 2, Sec. 23, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 75 percent; lush understory con-

Vegetation: Black spruce crown cover of about 75 percent; lush understory consists mostly of sedges with some leatherleaf and Labrador tea; ground cover consists of some cranberry.

Microrellef: 35 cm.

Depth To Water Table: 5 cm.

Described By: B. Wakanabo on August 23, 1977.

Layer	Depth (cm)
Fibric	0- 24
Hemic	24-329
Silt	329+

Location: 1650 feet S and 2020 feet W of the NE corner of Sec. 24, T.69N., R.27W. (SE1/4 of NW1/4 of SW1/4 of NE1/4, Sec. 24, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 95 percent; lush understory consists mostly of Labrador tea with some sedges; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: 20 cm.

Described By: D. Olson on August 23, 1977.

Layer	Depth (cm)
Fibric	0-115
Hemic	115-265
Silty	265+
clay	

Location: 1600 feet S and 670 feet W of the NE corner of Sec. 24, T.69N., R.27W. (NE1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 24, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 95 percent; understory consists of sedges and Labrador tea with some swamp laurel; ground cover consists of cranberry.

Microrellef: 40 cm.

Depth To Water Table: 25 cm.

Described By: B. Wakanabo on August 23, 1977.

Laver Depth (cm) 0- 84 Fibric Hemic 84-193 Silty 193+ clay

Location: 1680 feet S and 1950 feet E of the NW corner of Sec. 24, T.69N., R.27W. (SE1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 24, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent; just understory consists mostly of sedges with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 35 cm.

Depth To Water Table: 10 cm.

Described By: B. Wakanabo on August 23, 1977.

ayer_	Depth (cm
ibric	0- 74
lemic .	74-256
Clay	256+
nam	

Location: 2210 feet S and 2500 feet E of the NW corner of Sec. 14, T.152N., R.27W. (NE1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 14, T.152N., R.27W.)

Vegetation: Scattered black spruce; understory consists mostly of grasses with some leatherleaf, Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrelief: Negligible.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Depth To Water Table: Not visible.

Described By: T. Malterer and N. Aaseng on December 10, 1979.

Layer	Depth (cm)
Fibric	0- 91
Hemic	91-366
Sandy	366+
clay	
with	
pebbles	

Location: 1500 feet N and 250 feet E of the SW corner of Sec. 10, T.153N., R.27W.

Vegetation: Tamarack crown cover of about 65 percent; understory consists of

grasses with some plack spruce, sedges, Phragmites, cattails, Labrador tea,

and bog rosemary; ground cover consists of some mosses and pitcher plant.

(SW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 10, T.153N., R.27W.)

Described By: D. Mellem and S. Nelson on April 3, 1980.

Location: 500 feet S and 2580 feet E of the NW corner of Sec. 22, T.152N., R.27W. (SE1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 22, T.152N., R.27W.)

Vegetation: Northern white cedar crown cover of about 30 percent with some scattered black spruce and tamarack; sparse understory consists of some bog birch, willow, Labrador tea, speckled alder, leatherleaf, bog rosemary, grasses. and swamp cinquefoil; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: T. Malterer and N. Aaseng on December 10, 1979.

Layer	Depth (cm)
Hemic	0-218
Sapric	218-292
Loamy	292+
sand	

Location: 1400 feet N and 150 feet W of the SE corner of Sec. 11, T.153N., R.27W. (SE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 11, T.153N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some alder, leatherleaf, sedges, grasses, ferns, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal. Microrellef: 30 cm

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on April 3, 1980.

Layer	Depth (cm)
Fibric	0-183
Hemic	183-290
Loamy	290+
sand	
with	
calcareous	
pebbles	

Layer Depth (cm) Hemic 0-183 Sapric 183-244 Silty 244+ clav loam

Location: 500 feet N and 550 feet E of the SW corner of Sec. 24, T.69N., R.27W. (NE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 24, T.69N., R.27W.)

Vegetation: Black spruce crown cover of about 40 percent; Jush understory consists mostly of Labrador tea with grasses and some swamp laurel; ground cover consists of Sphagnum mosses and false Solomon's seal.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: B. Wakanabo on August 23, 1979.

Layer	Depth (cm)
Hemic Siltv	0-104 104+
clay	

Location: 1200 feet N and 1125 feet E of the SW corner of Sec. 3, T.153N., R.27W. (NE1/4 of NE1/4 of SW1/4 of SW1/4, Sec. 3, T.153N., R.27W.)

Vegetation: Northern white cedar crown cover of about 35 percent with scattered black spruce and tamarack; sparse understory consists of some leatherleaf. Labrador tea, alder, and sedges; ground cover consists mostly of Sphagnum mosses with some reindeer moss, snowberry, and false Solomon's seal.

Microrellef: 30 cm

Depth To Water Table: 7 cm.

Described By: H. Mooers and D. Haverkost on April 3, 1980.

Layer	Depth (cm)
Hemic	0-61
Sapric	61-91
Sand	91+

Location: 2365 feet N and 400 feet W of the SE corner of Sec. 36, T.153N., R.27W. (NW1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 36, T.153N., R.27W.)

Vegetation: Northern white cedar crown cover of about 85 percent with scattered balsam fir; sparse understory consists of some shrubs and grasses; ground cover consists of some mosses.

Microrellef: Negligible.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 25, 1976.

Layer	Depth (cm)
Sapric Clay	0-86 86+
oam	

Location: 120 feet N and 625 feet E of the SW corner of Sec. 1, T.154N., R.27W. (SE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 1, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 85 percent; sparse understory consists of balsam fir and tamarack; ground cover consists of Sphagnum mosses and feather mosses.

Microrellef: 30 cm.

Depth To Water Table: 15 cm.

Described By: T. Malterer on August 17, 1976.

Layer	Depth (cm
Hemic Sandy clay loam	0-137 137+

Location: 110 feet N and 1075 feet E of the SW corner of Sec. 1, T.154N., R.27W. (SE1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 1, T.154N., R.27W.)

Vegetation: Not recorded. Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: T. Malterer on August 17, 1976.

Layer	Depth (cm
Sapric Sandy clay	0-51 51+
loam	

Location: 150 feet S and 100 feet E of the NW corner of Sec. 1, T.154N., R.27W. (Gov. Lot 4, Sec. 1, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered paper birch; lush understory consists mostly of Labrador tea with cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 40 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm)
Hemic	0- 81
Sand	81+

Location: 2030 feet S and 870 feet E of the NW corner of Sec. 2, T.154N., T.27W. (NW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 2, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 40 percent; sparse understory consists of some grasses. Labrador tea, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, snowberry, and pitcher plant.

Microrelief: 20 cm.

Depth To Water Table: 30 cm.

Described By: B. Sether and D. Haverkost on March 27, 1980.

Layer	Depth (cm)
Fibric	0-152
Hemic	152-213
Fibric	213-274
Hemic	274-305
Sapric	305-320
Sandy	320+
clay	
with	
pebbles	

Location: 75 feet N and 685 feet E of the SW corner of Sec. 2, T.154N., R.27W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 2, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of leatherleaf and cotton grass with some bog rosemary and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm
Fibric	0- 46
Hemic	46-290
Sandy	290+
clay	
loam	

Location: 100 feet N and 2065 feet W of the SE corner of Sec. 2, T.154N., R.27W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 2, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 65 percent with scattered tamarack; lush understory consists mostly of Labrador tea and leatherleaf with cotton grass and some bog rosemary and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm
Fibric	0- 46
Hemic	46-254
Loamy sand	254+

Location: 100 feet N and 1575 feet W of the SE corner of Sec. 2, T.154N., R.27W. (SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 2, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; understory consists of some leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some snowberry.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and L. Severson on March 25, 1979.

Depth (cm) Layer Fibric 0- 61 61- 91 Hemic Fibric 91-122 Hemic 122-152 Fibric 152-183 Sapric 183-213 Hemic 213-274 Sapric 274-290 290+ Sandy

Location: 125 feet N and 860 feet W of the SE corner of Sec. 2, T.154N., R.27W. (SE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 2, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 40 percent with scattered tamarack; jush understory consists mostly of leatherleaf with some swamp laurel, bog rosemary, cotton grass, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some pitcher plant.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm)
Hemic Sandy	0-244 244+
clay	

Location: 50 feet S and 2000 feet W of the NE corner of Sec. 2, T.154N., R.27W. (Gov. Lot 2, Sec. 2, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of Labrador tea; ground cover consists mostly of Sphagnum mosses. Microrellef: 25 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm
Hemic	0-183
Sandy.	183+
clay	
loam	

Location: 800 feet S and 1990 feet E of the NW corner of Sec. 2, T.154N., R.27W. (Gov. Lot 3, Sec. 2, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 15 percent; understory consists of black spruce with some Labrador tea, leatherleaf, swamp laurel, bog rosemary, and sedges; ground cover consists mostly of sphagnum mosses with some other mosses, cranberry, and snowberry.

Microrellef: 5 cm.

Depth To Water Table: 30 cm.

Described By: B. Sether and D. Haverkost on March 26, 1980.

Depth (cm
0-122
122-152
152-244
244-290
290+

Location: 2275 feet N and 750 feet E of the SW corner of Sec. 3, T.154N., R.27W. (SW1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of leatherleaf with cotton grass and some Labrador tea; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.
Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm)
Peat	0-244
Sandy	244+
loam	

Location: 75 feet N and 65 feet E of the SW corner of Sec. 3, T.154N., R.27W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 50 percent with a tamarack crown cover of about 40 percent; lush understory consists mostly of leatherleaf with Labrador tea, swamp laurel and some cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 15 cm.

Described By: M. Domeier on August 16, 1976.

Layer	Depth (cm)
Fibric	0- 18
Hemic	18-127
Sandy	127+
clay	
loom	

Location: 225 feet N and 700 feet E of the SW corner of Sec. 3, T.154N., R.27W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of some Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: 25 cm.

Described By: H. Mooers and L. Severson on March 27, 1980.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-213
Sapric	213-244
Medium	244+
sand	

Location: 100 feet N and 700 feet E of the SW corner of Sec. 3, T.154N., R.27W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 3, T.154N., R.27W.)

Vegetation: Lush understory consists mostly of leatherleaf and Labrador tea with some swamp laurel, black spruce, and tamarack; ground cover consists of Sphagnum mosses and other mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 16, 1976.

Depth (cm)
0- 18
18-163
163+

Location: 100 feet N and 2200 feet E of the SW corner of Sec. 3, T.154N., R.27W. (SW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; lush understory consists mostly of leatherleaf with cotton grass and some swamp laurel and bog rosemary; ground cover consists mostly of Sphagnum mosses with some pitcher plant.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 16, 1976.

Layer	Depth (cm)
Fibric	0- 23
Hemic	23-249
Clay	249+
loam	

Location: 100 feet N and 2575 feet W of the SE corner of Sec. 3, T.154N., R.27W. (SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 3, T.154N., R.27W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of cotton grass and swamp laurel with Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrellef: 40 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 16, 1976.

Layer	Depth (cm)
Hemic	0-259
Medium	259+
sand	

Location: 100 feet N and 675 feet W of the SE corner of Sec. 3, T.154N., R.27W. (SW1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists of Labrador tea, swamp laurel, and cotton grass with some bog rosemary and leatherleaf; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrelief: 45 cm.

Depth To Water Table: 15 cm.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm)
Hemic	0- 30
Fibric	30- 50
Hemic	50-310
Sandy	310+
clav	
loam	

Location: 100 feet N and 75 feet W of the SE corner of Sec. 3, T.154N., R.27W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of some Labrador tea, bog rosemary, leatherleaf, and grasses; ground cover consists mostly of Sphagnum mosses with some feather mosses, cranberry, and snowberry.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: H. Mooers and L. Severson on March 27, 1980.

.ayer	Depth (cm)
ibric	0- 61
lemic	61- 91
ibric	91-122
lemic	122-152
ibric	152-183
łemic	183-335
.oam	335+

Location: 800 feet S and 600 feet W of the NE corner of Sec. 3, T.154N., R.27W. (Gov. Lot 1, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of Labrador tea with some cotton grass, swamp laurel, leatherleaf, and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm
Fibric	0- 20
Hemic	20-249
Sandy	249+
clay	
loam	

Location: 25 feet S and 1690 feet W of the NE corner of Sec. 3, T.154N., R.27W. (Gov. Lot 2, Sec. 3, T.154N., R.27W.)

Vegetation: Lush understory consists mostly of black spruce with some Labrador tea, leatherleaf, swamp laurel, and cotton grass; ground cover consists mostly of Sohannum mosses with some cranberry and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 26, 1976.

Layer	Depth (cm)
Fibric	0- 76
Hemic	76-213
Sapric	213-244
Clay	244+
loam	

Location: 840 feet S and 2550 feet W of the NE corner of Sec. 3, T.154N., R.27W. (Gov. Lot 2, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; sparse understory consists of some Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 17, 1976.

Layer	Depth (cm)
Peat	0-249
Sandy	249+
loam	

Location: 25 feet S and 2075 feet E of the NW corner of Sec. 3, T.154N., R.27W. (Gov. Lot 3, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of some Labrador tea, leatherleaf, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, and cranberry.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth and R. Fox on August 26, 1976

Layer	Depth (cm)
Fibric	0- 10
Hemic	10- 51
Sapric	51-244
Medium	244+
sand	

Location: 25 feet S and 755 feet E of the NW corner of Sec. 3, T.154N., R.27W. (Gov. Lot 4, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of Labrador tea, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 19, 1976.

Layer	Depth (cm)
Fibric	0- 20
Hemic	20-198
Mineral	198+
soil	

Location: 875 feet S and 825 feet E of the NW corner of Sec. 3, T.154N., R.27W. (Gov. Lot 4, Sec. 3, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of leatherleaf with cotton grass and some Labrador tea; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier and T. Malterer on August 17, 1976.

Layer	Depth (cm)
Peat	0-183
Loamy	183+
sand	

Location: 50 feet N and 2150 feet E of the SW corner of Sec. 4, T.154N., R.27W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 4, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of Labrador tea and horsetail with some willow, balsam fir, and paper birch; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 16, 1976.

Layer	Depth (cm)
Hemic Clay loam	0-51 51+

Location: 30 feet N and 2610 feet W of the SE corner of Sec. 4, T.154N., R.27W. (SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 4, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 85 percent; sparse understory consists of some cotton grass, Labrador tea, and Indian pipes; ground cover consists mostly of Sphagnum and other mosses.

Microrelief: 50 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 16, 1976.

Layer	Depth (cm)
Hemic	0-76
Sandy	76+
clay	
loam	

Location: 50 feet N and 1650 feet W of the SE corner of Sec. 4, T.154N., R.27W. (SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 4, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of swamp laurel with Labrador tea and some cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 16, 1976.

.ayer	Depth (cm)
ibric	0- 25
lemic	25-122
Sandy	122+
lay	
oam	

Location: 75 feet N and 660 feet W of the SE corner of Sec. 4, T.154N., R.27W. (SE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 4, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of leatherleaf with Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 16, 1976.

Layer	Depth (cm
Fibric	0-25
Hemic	25-61
Medium sand	61+

Location: 75 feet S and 325 feet W of the NE corner of Sec. 4, T.154N., R.27W. (Gov. Lot 1, Sec. 4, T.154N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth and R. Fox on August 20, 1976.

Depth (cm)
0- 33
33-102
102+

Location: 50 feet S and 550 feet W of the NE corner of Sec. 4, T.154N., R.27W. (Gov. Lot 1, Sec. 4, T.154N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth and R. Fox on August 20, 1976.

Layer	Depth (cm)
Hemic	0- 33
Sapric	33-101
Clay	101+
loam	

Location: 50 feet S and 850 feet W of the NE corner of Sec. 4, T.154N., R.27W. (Gov. Lot 1, Sec. 4, T.154N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth and R. Fox on August 20, 1976.

Layer	Depth (cm
Hemic	0-76
Clay	76+
loam	

Location: 75 feet S and 1500 feet W of the NE corner of Sec. 4, T.154N., R.27W. (Gov. Lot 2, Sec. 4, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; sparse understory consists of horsetail with some Labrador tea, sedges, and balsam fir; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 45 cm.

Depth To Water Table: Not recorded. Described By: M. Domeier on August 24, 1976.

Depth (cm)
0- 10
10- 22
22- 40
40+

Location: 75 feet S and 1800 feet W of the NE corner of Sec. 4, T.154N., R.27W.

(Gov. Lot 2, Sec. 4, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of Labrador tea and horsetail, with some bog birch, sedges, and ferns; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 24, 1976.

Layer	Depth (cm)
Fibric	0- 8
Sapric	8-15
Fibric	15-23
Hemic	23-28
Mineral	28+
soil	

Location: 2225 feet S and 1750 feet E of the NW corner of Sec. 5, T.154N., R.27W. (NW1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 5, T.154N., R.27W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of leatherleaf with swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox on September 13, 1976.

Depth (cm)
0- 30
30-162
162-198
198+

Location: 125 feet N and 290 feet E of the SW corner of Sec. 5, T.154N., R.27W. (SW1/4 of SW1/4 of SW1/4, Sec. 5, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 85 percent; understory consists of Labrador tea with some cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox on September 9, 1976.

Layer	Depth (cm)
Fibric	0- 20
Hemic	20-142
Sapric	142-216
Silt	216+
loam	

Location: 150 feet N and 1650 feet E of the SW corner of Sec. 5, T.154N., R.27W. (SW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 5, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; lush understory consists mostly of leatherleaf with some sedges and speckled alder; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 13, 1976.

Layer	Depth (cm)
Fibric	0- 25
Hemic	25-106
Clay	106+
loam	

Location: 75 feet S and 1860 feet E of the NW corner of Sec. 5, T.154N., R.27W. (Gov. Lot 3, Sec. 5, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 50 percent; understory consists of leatherleaf and cotton grass with some Labrador tea and swamp laurel; ground cover consists of Sohagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 16, 1976.

 Layer
 Depth (cm)

 Hemic
 0-198

 Sapric
 198-269

 Sand
 269+

Location: 100 feet S and 120 feet E of the NW corner of Sec. 5, T.154N., R.27W. (Gov. Lot 4, Sec. 5, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of leatherleaf and cotton grass with some Labrador tea and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellet: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 16, 1976.

Layer	Depth (cm)
Hemic	0-283
Coarse	283+
sandy	
loam	
with	
limestone	
pebbles	

Location: 100 feet N and 1690 feet E of the SW corner of Sec. 6, T.154N., R.27W. (SW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 6, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 50 percent; lush understory consists mostly of Labrador tea with some leatherleaf, swamp laurel, and cotton grass; ground cover consists of some false Solomon's seal.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 13, 1976.

Layer	Depth (cm)
Fibric	0- 18
Hemic	18-198
Sand	198+

Location: 2215 feet N and 635 feet W of the SE corner of Sec. 6, T.154N., R.27W. (SW1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 6, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 65 percent; lush understory consists mostly of leatherleaf with grasses and some Labrador tea and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 28, 1976.

Layer	Depth (cm
Fibric	0- 30
Hemic	30-335
Mineral	335+
soil	

Location: 125 feet N and 725 feet W of the SE corner of Sec. 6, T.154N., R.27W. (SE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 6, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of leatherleaf with some Labrador tea and swamp laurel; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrellef: 15 cm.

Microrellei: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 13, 1976.

Layer	Depth (cm)
Fibric	0- 18
Hemic	18-147
Clay	147+
loam	

Location: 1400 feet S and 740 feet E of the NW corner of Sec. 6, T.154N., R.27W. (Gov. Lot 5, Sec. 6, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of cotton grass with some leatherleaf and Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: R. Zarth and R. Fox on September 9, 1976.

Layer	Depth (cr
Fibric	0- 75
Hemic	75-338
Sand	338+

Location: 75 feet N and 80 feet E of the SW corner of Sec. 8, T.154N., R.27W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 8, T.154N., R.27W.)

Vegetation: Scattered black spruce, balsam fir, and paper birch; lush understory consists mostly of speckled alder with grasses.

Microrelief: 15 cm.

Depth To Water Table: Not visible.

Described By: M. Domeier on November 3, 1976.

Layer	Depth (cm
Hemic	0- 30
Coarse	30+
sand	
with	
pebbles	

Location: 50 feet N and 1600 feet E of the SW corner of Sec. 8, T.154N., R.27W. (SW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 8, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent with a tamarack crown cover of about 35 percent with scattered paper birch; understory consists of bog birch, speckled alder, and grasses with some sedges; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 3, 1976.

Layer	Depth (cm)
Hemic	0-101
Silt	101+
loam	

Location: 75 feet N and 2380 feet W of the SE corner of Sec. 8, T.154N., R.27W. (SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 8, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack and paper birch; lush understory consists mostly of speckled alder with Labrador tea and some leatherleaf and fern.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 3, 1976.

Depth (cm
0-122 122+

Location: 550 feet N and 1300 feet W of the SE corner of Sec. 8, T.154N., R.27W. (NW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 8, T.154N., R.27W.)

Vegetation: Consists mostly of sedges and speckled alder.

Microrellef: Negligible.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 3, 1976.

Laver Depth (cm) Hemic 0-76 Clay 76+ loam

Location: 2050 feet S and 25 feet E of the NW corner of Sec. 11, T.154N., R.27W. (NW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 11, T.154N., R.27W.)

Vegetation: Scattered black spruce; understory consists mostly of leatherleaf with some Labrador tea and sedges; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 1, 1976.

.ayer	Depth (cm
ibric	0- 61
łemic	61-257
Clay	257+
oam	

Location: 325 feet N and 40 feet E of the SW corner of Sec. 11, T.154N., R.27W.

Vegetation: Black spruce crown cover of about 50 percent; lush understory consists mostly of Labrador tea with leatherleaf and some sedges and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

(SW1/4 of SW1/4 of SW1/4, Sec. 11, T.154N., R.27W.)

Microrellef: 30 cm.,

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox on August 31, 1976.

Depth To Water Table: At surface.

Described By: R. Fox on September 1, 1976.

Layer Depth (cm) Fibric 0- 61 Hemic 61-254 Sapric 254-272 272+ Clay loam

Location: 1300 feet S and 25 feet E of the NW corner of Sec. 12, T.154N., R.27W.

Vegetation: Black spruce crown cover of about 60 percent; lush understory con-

sists mostly of leatherleaf with some Labrador tea, swamp laurel, sedges, cot-

ton grass, and bog rosemary; ground cover consists mostly of Sphagnum

(SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 12, T.154N., R.27W.)

mosses with some false Solomon's seal and cranberry.

Location: 250 feet N and 1175 feet E of the SW corner of Sec. 11, T.154N., R.27W. (SE1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 11, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of leatherleaf with cotton grass and some Labrador tea and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry and other mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 1, 1976.

Layer	Depth (cm)
Fibric	0- 76
Hemic	76-287
Coarse	287+
sand	

Location: 2400 feet S and 2025 feet E of the NW corner of Sec. 12, T.154N., R.27W. (SW1/4 of SW1/4 of SE1/4 of NW1/4. Sec. 12. T.154N.. R.27W.)

Vegetation: Black spruce crown cover of about 60 percent: understory consists of leatherleaf and Labrador tea with some swamp laurel: ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 3, 1976.

Layer Depth (cm) Fibric 0- 51 Hemic 51-244 Sapric 244-295 Sand 295+

Layer Depth (cm) 0-228 Hemic Sapric 228-246 Sand 246 -

Location: 2230 feet N and 25 feet E of the SW corner of Sec. 11, T.154N., R.27W. (SW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 11, T.154N., R.27W.)

Vegetation: Scattered tamarack; lush understory consists mostly of sedges and leatherleaf with some bog birch; ground cover consists mostly of Sphagnum mosses

Microrelief: Not recorded.

Depth To Water Table: At surface.

Described By: R. Fox on September 1, 1976.

Depth (cm
0- 46
46-203
203+
,

Location: 125 feet N and 2075 feet W of the SE corner of Sec. 11, T.154N., R.27W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 11, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of leatherleaf with some Labrador tea and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on August 31, 1976.

Layer	Depth (cm)
Fibric	0- 51
Hemic	51-279
Clay	279+
loam	

Location: 2425 feet N and 25 feet E of the SW corner of Sec. 12.T.154N., R.27W. (NW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 12, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of sedges with some Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Fox on August 31, 1976.

Depth (cm
0- 15
15- 51
51- 66
66-279
279+

Location: 25 feet N and 25 feet E of the SW corner of Sec. 12, T.154N, B.27W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 12, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of leatherleaf with some Labrador tea, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranherry

Microrellef: 45 cm.

Depth To Water Table: At surface. Described By: R Fox in August, 1976.

(Gov. Lot 2, Sec. 12, T.154N., R.27W.)

Described By: M. Domeier on September 3, 1976. Layer

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Sphagnum and other mosses with some cranberry.

Fibric

Hemic

Layer	Depth (cm)
Fibric	0- 15
Hemic	15-307
Sapric	307-333
Clay	333+
loam	

Location: 2375 feet N and 1575 feet W of the SE corner of Sec. 12, T.154N., R.27W. (NE1/4 of NE1/4 of NW1/4 of SE1/4, Sec. 12, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of Labrador tea with some leatherleaf and cotton grass; ground cover consists mostly of Sphagnum mosses with other mosses.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 3, 1976.

Layer	Depth (cm)
Hemic	0-185
Sapric	185-189
Sandy	189+
loam	

Location: 2480 feet S and 550 feet W of the NE corner of Sec. 12, T.154N., R.27W. (Gov. Lot 2, Sec. 12, T.154N., R.27W.)

Vegetation: Not recorded.

Depth To Water Table: Not recorded.

0- 7 7-172

Depth (cm)

178+

Location: 2575 feet S and 1100 feet W of the NE corner of Sec. 12, T.154N., R.27W.

Vegetation: Black spruce crown cover of about 85 percent; lush understory con-

sists mostly of Labrador tea with some leatherleaf; ground cover consists of

Sapric 172-178 Loam

Microrelief: Not recorded.

Described By: M. Domeier on September 3, 1976.

Layer	Depth (cm)
Hemic	0-127
Sapric	127-135
Loam	135+
with	
calcareous	
pebbles	

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 3, 1976.

Laver Fibric

Hemic

Sapric

(Gov. Lot 2, Sec. 12, T.154N., R.27W.)

Described By: M. Domeier on September 3, 1976.

Laver

Sapric

Hemic

Sapric

Sandy loam

of some Sphagnum mosses.

Depth To Water Table: Not recorded.

Microrelief: Not recorded.

Sampler would not penetrate

Location: 2590 feet S and 1825 feet W of the NE corner of Sec. 13, T.154N., R.27W. (SW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 13, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of Labrador tea with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Location: 975 feet S and 1800 feet E of the NW corner of Sec. 13, T.154N., R.27W. (SE1/4 of SW1/4 of NE1/4 of NW1/4, Sec. 13, T.154N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, swamp laurel, bog rosemary, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 20 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and L. Severson on March 27, 1980.

Layer Depth (cm) Fibric 0- 61 Hemic 61-122 122-213 Fibric Hemic 213-244 244-274 Fibric Hemic 274-366 366-381 Sapric Clay 381+

Location: 1700 feet N and 50 feet E of the SW corner of Sec. 13. T.154N., R.27W.

Location: 50 feet N and 1725 feet W of the SE corner of Sec. 12, T.154N, R 27W,

Vegetation: Black spruce crown cover of about 60 percent; understory consists of

leatherleaf with some Labrador tea and swamp laurel; ground cover consists

mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Depth (cm)

0- 51

51-325

325-376

Location: 2350 feet S and 250 feet W of the NE corner of Sec. 12, T.154N., R.27W.

Vegetation: Black spruce crown cover of about 60 percent; ground cover consists

Depth (cm)

0-25

25-40

40-46

46+

(SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 12, T.154N., R.27W.)

(NW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 13, T.154N., R.27W.) Vegetation: Black spruce crown cover of about 85 percent; lush understory consists mostly of leatherleaf with some cotton grass and Labrador tea; ground cover consists mostly of Sphagnum mosses with some pitcher plant and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Laver	Donath (non)
Layer	Depth (cm)
Hemic	0-434
Clay	434+
loam	

Microrellef: Not recorded. Depth To Water Table: Not recorded. Described By: M. Domeier on September 3, 1976.

> Laver Depth (cm) Hemic 0-427 Mineral 427+ soil

Location: 1300 feet N and 375 feet E of the SW corner of Sec. 13. T.154N., R.27W. (SW1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

Vegetation: Consists mostly of black spruce.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Depth (cm) Laver Peat 0-384 Mineral 384+ soil

Location: 1440 feet N and 475 feet E of the SW corner of Sec. 13, T.154N., R.27W. (NE1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

Vegetation: Consists mostly of tamarack.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm
Peat	0-381
Mineral soil	381+

Location: 625 feet N and 50 feet E of the SW corner of Sec. 13, T.154N., R.27W. (NW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., B.27W.)

Vegetation: Consists of tamarack bog birch, leatherleaf, and swamp laurel. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Laver

Peat

Sandy

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Peat	0-216
Fine	216+
sand	

Location: 50 feet N and 275 feet E of the SW corner of Sec. 13, T.154N., R.27W.

Vegetation: Consists of tamarack, bog birch, leatherleaf, and Sphagnum mosses.

Depth (cm)

0-127

127 +

(SW1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

Location: 300 feet N and 50 feet E of the SW corner of Sec. 13, T.154N., R.27W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

Vegetation: Consists of tamarack, leatherleaf, and Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Peat	0-216
Clay	216+
loam	

Location: 50 feet N and 525 feet E of the SW corner of Sec. 13, T.154N., R.27W. (SE1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

Vegetation: Crown cover consists of black spruce and tamarack; understory consists of leatherleaf, bog rosemary, and grasses; ground cover consists of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Peat	0-208
Fine	208+
sand	

Location: 50 feet N and 750 feet E of the SW corner of Sec. 13, T.154N., R.27W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

Location: 1125 feet N and 50 feet E of the SW corner of Sec. 13, T.154N., R.27W.

Depth (cm)

0-353

353 +

Location: 50 feet N and 50 feet E of the SW corner of Sec. 13, T.154N., R.27W.

Vegetation: Tamarack crown cover of about 60 percent with scattered black spruce; lush understory consists mostly of bog birch with Labrador tea and

Depth (cm)

0- 7

7-36

36+

(SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

(NW1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Laver

Peat Mineral

enil

some leatherleaf, swamp laurel and sedges.

Layer

Sapric

Hemic

Sand

Microrellef: Not recorded. Depth To Water Table: Not recorded. Described By: M. Domeier on September 3, 1976.

Vegetation: Consists of tamarack, bog birch, leatherleaf, and swamp laurel.

Vegetation: Crown cover consists of black spruce and tamarack; understory consists of leatherleaf, bog birch, bog rosemary, and grasses; ground cover consists of Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Peat	0-244
Fine	244+
sand	

Location: 50 feet N and 925 feet E of the SW corner of Sec. 13, T.154N., R.27W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.27W.)

Vegetation: Not recorded. Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 3, 1976.

Depth (cm) Layer Hemic 0-231 Mineral 231+

Location: 50 feet N and 1865 feet W of the SE corner of Sec. 13, T.154N., R.27W. (SW1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 13, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of leatherleaf with cotton grass and some Labrador tea and bog rosemary; ground cover consists mostly of Sphagnum mosses with some false

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 3, 1976,

Layer	Depth (cm)
Fibric Hemic	0- 25 25-432
Clay loam	432+

Location: 50 feet N and 1600 feet W of the SE corner of Sec. 14, T.154N., B 27W (SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 14, T.154N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 3, 1976

Layer	Depth (cm)
Hemic Sapric Mineral soil	0-335 335-360 360+

Location: 50 feet N and 800 feet W of the SE corner of Sec. 14, T.154N., R.27W. (SE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 14, T.154N., R.27W.)

Vegetation: Not recorded. Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 3, 1976.

Layer	Depth (cm)
Hemic	0-215
Sapric	215-231
Mineral soil	231+

Location: 50 feet N and 335 feet W of the SE corner of Sec. 14, T.154N., R.27W. (SW1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 14, T.154N., R.27W.)

Vegetation: Crown cover consists of tamarack; understory consists of bog birch, bog rosemary, leatherleaf, and grasses; ground cover consists of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 3, 1976.

Layer	Depth (cm)
Peat Fine	0-132
sand	132+

Location: 2150 feet S and 175 feet E of the NW corner of Sec. 15, T.154N., R.27W. (NW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 15, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; lush understory consists mostly of Labrador tea with speckled alder. ferns, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 30, 1976.

Layer	Depth (cm)
Hemic	0-300
Sapric	300-312
Clay	312+

Location: 2270 feet N and 1290 feet W of the SE corner of Sec. 16, T.154N., R.27W. (SW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 16, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; lush understory consists mostly of Labrador tea with some cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 30, 1976.

Layer	Depth (cm)
Hemic	0-274
Clay	274+

Location: 1050 feet S and 1880 feet W of the NE corner of Sec. 17, T.154N., R.27W. (SW1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 17, T.154N., R.27W.)

Vegetation: Balsam fir crown cover of about 60 percent; lush understory consists mostly of grasses with speckled alder and some ferns; ground cover consists of some feather mosses.

Microrelief: Negligible.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 3, 1976.

Layer	Depth (cm)
Sapric	0-120
Sandy	120+
loam	

Location: 2075 feet S and 1880 feet W of the NE corner of Sec. 17, T.154N., R.27W. (NW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 17, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent with a northern white cedar crown cover of about 35 percent; understory consists of Labrador tea. sedges, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded

Described By: M. Domeier on November 3, 1976.

Depth (cn
0-137 137+

Location: 2260 feet N and 1880 feet W of the SE corner of Sec. 17, 7,154N., R.27W. (SW1/4 of NE1/4 of NW1/4 of SE1/4, Sec. 17, T.154N., R.27W.)

Vegetation: Scattered black spruce, tamarack, and northern white cedar; understory consists of speckled alder with some Labrador tea and leatherleaf; ground cover consists of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 3, 1976.

Depth (cm) Laver Hemic 0-137 Clav 137+ loam

Location: 1065 feet N and 1880 feet W of the SE corner of Sec. 17, T.154N., R.27W. (NW1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 17, T.154N., R.27W.)

Vegetation: Northern white cedar crown cover of about 70 percent with a black spruce crown cover of about 25 percent; understory consists of some grasses. leatherleaf, sedges, and Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 3, 1976.

Layer	Depth (cm)
Hemic	0-97
Clay	97+
nam	

Location: 1200 feet S and 800 feet W of the NE corner of Sec. 22, T.154N., R.27W.

Vegetation: Northern white cedar crown cover of about 60 percent with scattered paper birch; understory consists of grasses with some speckled aider; ground cover consists mostly of Sphagnum mosses.

(SE1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 22, T.154N., R.27W.)

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 2, 1976.

Layer	Depth (cm
Hemic	0-127
Sand	127⊥

Location: 430 feet S and 1500 feet W of the NE corner of Sec. 22, T.154N., R.27W. (SE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 22, T.154N., R.27W.)

Vegetation: Consists of sedges with some bog birch, willow, and speckled alder.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 30, 1976.

Layer	Depth (cn
Peat Mineral soil	0-58 58+

Microrelief: Not recorded.

Location: 2275 feet S and 425 feet E of the NW corner of Sec. 23. T.154N.. R.27W. (NE1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 23, T.154N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on September 2, 1976.

Depth (cm) Layer 0-259 Hemic 259-289 Sapric Sandy 289+ loam

Location: 1990 feet N and 1610 feet E of the SW corner of Sec. 23, T.154N., R.27W.

Location: 900 feet S and 1275 feet W of the NE corner of Sec. 22, T.154N., R.27W.

Vegetation: Tamarack crown cover of about 50 percent; lush understory consists

Depth (cm)

0-127

127-134

134+

Location: 2075 feet N and 2090 feet W of the SE corner of Sec. 22, T.154N., R.27W.

Vegetation: Tamarack crown cover of about 35 percent; lush understory consists

mostly of bog birch and grasses with some fern; ground cover consists of

Depth (cm)

0-290

290+

(SE1/4 of NW1/4 of NW1/4 of SE1/4, Sec. 22, T.154N., R.27W.)

(NW1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 22, T.154N., R.27W.)

mostly of bog birch with some speckled alder.

Layer

Sapric

sand

Clay

sand

Medium

Described By: M. Domeier on September 2, 1976.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

(NW1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 23, T.154N., R.27W.) Vegetation: Balsam fir crown cover of about 60 percent with scattered northern white cedar; lush understory consists mostly of grasses with speckled alder and ferns; ground cover consists of some feather mosses.

Microrelief: Not recorded.

Sphagnum mosses.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 30, 1976.

Layer Hemic

Fine

sandy

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth on August 26, 1976.

Layer	Depth (cm
Hemic	0-391
Sapric	391-427
Sandy	427+
loam	

cover consists mostly of Sphagnum mosses. Microrellef: 30 cm. Depth To Water Table: Not recorded. Described By: R. Fox on September 2, 1976.

> Layer Depth (cm) Hemic 0-188 188-216 Sapric 216+

Location: 1700 feet S and 50 feet E of the NW corner of Sec. 23, T.154N., R.27W.

Vegetation: Northern white cedar crown cover of about 60 percent with scattered

paper birch; understory consists of sedges with some speckled alder; ground

(SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 23, T.154N., R.27W.)

Location: 1125 feet N and 125 feet W of the SE corner of Sec. 23, T.154N., R.27W. (SE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 23, T.154N., R.27W.)

Vegetation: Tamarack crown cover of about 35 percent; lush understory consists mostly of grasses with bog birch and some bog rosemary and fern; ground cover consists of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Hemic Fine sand	0-368 368+

Location: 575 feet N and 775 feet W of the SE corner of Sec. 23, T.154N., R.27W. (NW1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 23, T.154N., R.27W.)

Vegetation: Tamarack crown cover of about 35 percent with scattered black spruce; lush understory consists mostly of grasses with some bog birch; ground cover consists of Sphagnum and other mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Hemic Fine	0-353 353+
sand	333 T

Location: 965 feet N and 390 feet W of the SE corner of Sec. 23, T.154N., R.27W. (SW1/4 of NW1/4 of SE1/4 of SE1/4, Sec. 23, T.154N., R.27W.)

Vegetation: Scattered black spruce and northern white cedar; understory consists of bog birch, cattails, Labrador tea, and leatherleaf; ground cover consists of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm
Peat	0-389
Fine	389+
sandy	
loam	

Location: 250 feet S and 25 feet E of the NW corner of Sec. 24, T.154N., R.27W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 24, T.154N., R.27W.)

Depth (cm)

0-91

91+

Vegetation: Consists of tamarack and bog birch.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer Peat

Mineral

soil

Location: 475 feet S and 25 feet E of the NW corner of Sec. 24, T.154N., R.27W. (SW1/4 of NW1/4 of NW1/4, Sec. 24, T.154N., R.27W.)

Vegetation: Not recorded.
Microrelief: Not recorded.
Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Hemic	0-127
Sand	127+

Location: 725 feet S and 25 feet E of the NW corner of Sec. 24, T.154N., R.27W. (NW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 24, T.154N., R.27W.)

Vegetation: Consists of tamarack, bog birch, leatherleaf, and bog rosemary.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Hemic	0-191
Clay	191+

Location: 1000 feet S and 25 feet E of the NW corner of Sec. 24, T.154N., R.27W. (SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 24, T.154N., R.27W.)

Vegetation: Crown cover consists of tamarack; understory consists of leatherleaf, bog birch, and bog rosemary; ground cover consists of Sphagnum mosses.

Depth (cm)

0-282

282+

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer

Peat

sand

Coarse

Location: 1910 feet N and 660 feet E of the SW corner of Sec. 24, T.154N., R.27W. (SE1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 24, T.154N., R.27W.)

Vegetation: Tamarack crown cover of about 60 percent; lush understory consists mostly of grasses and bog rosemary with bog birch; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)
Peat	0-356
Fine	356+
sand	

Location: 400 feet S and 575 feet W of the NE corner of Sec. 24, T.154N., R.27W. (Gov. Lot 1, Sec. 24, T.154N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists of grasses with some swamp laurel and Labrador tea; ground cover consists mostly of Sphagnum mosses with some snowberry.

Microrelief: 15 cm.

Depth To Water Table: 13 cm.

Described By: H. Mooers and L. Severson on March 26, 1980.

Layer	Depth (cm)	
Fibric	0- 91	
Hemic	91-122	
Fibric	122-183	
Hemic	183-305	
Fibric	305-335	
Hemic	335-457	
Sapric	457-488	
Fine	488+	
silt		

Location: 2480 feet N and 1175 feet E of the SW corner of Sec. 25, T.154N., R.27W. (NE1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 25, T.154N., R.27W.)

Vegetation: Tamarack crown cover of about 35 percent with scattered black spruce; lush understory consists mostly of sedges and cotton grass with leatherleaf and bog birch with some bog rosemary, Labrador tea, horsetail, cattail, and pitcher plant; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Zarth on August 26, 1976.

some other mosses and cranberry.

Laver

Sandy

Depth To Water Table: Not recorded. Described By: R. Zarth on August 26, 1976.

Microrelief: Not recorded

Layer Depth (cm) Hemic 0-279 Sandy 279+ loam

Location: 125 feet S and 1500 feet W of the NE corner of Sec. 26, T.154N., R.27W.

spruce: lush understory consists mostly of sedges and cotton grass with

leatherleaf and bog birch with some bog rosemary, Labrador tea, horsetail, cat-

tail, and pitcher plant; ground cover consists mostly of Sphagnum mosses with

Depth (cm)

0-325

325+

(NE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 26, T.154N., R.27W.) Vegetation: Tamarack crown cover of about 35 percent with scattered black Location: 1050 feet N and 2630 feet W of the SE corner of Sec. 25, T.154N., R.27W. (SE1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 25, T.154N., R.27W.)

Vegetation: Tamarack crown cover of about 35 percent; lush understory consists mostly of leatherleaf with bog birch, bog rosemary, and cattails; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Zarth on August 27, 1976.

Layer	Depth (cm)
Hemic Sandy Ioam	0-262 262+

Location: 1550 feet S and 25 feet W of the NE corner of Sec. 26, T.154N., R.27W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 26, T.154N., R.27W.)

Vegetation: Tamarack crown cover of about 35 percent with scattered black spruce; lush understory consists mostly of sedges and cotton grass with leatherleaf and bog birch with some bog rosemary, Labrador tea, horsetail, cattail, and pitcher plant; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 30 cm.

Denth To Water Table: At surface.

Described By: R. Zarth on August 26, 1976.

Depth (cm) Layer 0-287 Sandy

Location: 2150 feet S and 1550 feet E of the NW corner of Sec. 26, T.154N., R.27W. (NW1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 26, T.154N., R.27W.)

Location: 60 feet N and 1550 feet W of the SE corner of Sec. 25, T.154N., R.27W.

Vegetation: Scattered tamarack and black spruce: Jush understory consists mostly

of leatherleaf with Labrador tea, bog birch, and bog rosemary; ground cover

Depth (cm)

0-196

196+

(SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 25, T.154N., R.27W.)

consists mostly of Sphagnum mosses.

Described By: R. Zarth on August 27, 1976.

Laver Hemic

Sandy

Depth To Water Table: At surface.

Microrelief: 45 cm.

Vegetation: Tamarack crown cover of about 35 percent; lush understory consists mostly of grasses with bog birch and cattails with some swamp laurel; ground cover consists of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 27, 1976.

Layer	Depth (cm)	
Hemic	0-264	
Sandy	264+	
loam		

Location: 1110 feet N and 150 feet E of the SW corner of Sec. 26, T.154N., R.27W. (NW1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 26, T.154N., R.27W.)

Vegetation: Tamarack crown cover of about 35 percent; lush understory consists mostly of grasses with some cattail, reed sedge, leatherleaf, bog rosemary, and bog birch; ground cover consists of Sphagnum mosses with some feather mosses, snowberry, and false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: B. Sether and S. Nelson on April 3, 1980.

Fibric

Hemic

Loamy

sand

Layer Depth (cm)

0-213

213-274

274+

Location: 1610 feet S and 875 feet E of the NW corner of Sec. 27. T.154N., R.27W. (NW1/4 of NE1/4 of SW1/4 of NW1/4, Sec. 27, T.154N., R.27W.)

Vegetation: Scattered tamarack and black spruce; lush understory consists mostly of bog birch with northern white cedar, sedges, and grasses with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded. Depth To Water Table: Not recorded.

Described By: R. Fox on October 1, 1976.

Layer	Depth (cm)
Hemic	0-217
Fine	217+
and	
coarse	
aaad	

Location: 1825 feet S and 700 feet E of the NW corner of Sec. 32, T.154N., R.27W. (SE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 32, T.154N., R.27W.)

Vegetation: Scattered paper birch and tamarack; lush understory consists mostly of speckled alder and sedges; ground cover consists of some Sphagnum mosses.

Microrelief: Not recorded. Depth To Water Table: Not recorded. Described By: R. Fox on October 1, 1976.

Layer	Depth (cm)	
Sapric	0-56	
Loam	56+	

Location: 1250 feet S and 100 feet E of the NW corner of Sec. 3, T.155N., R.27W. (Gov. Lot 4, Sec. 3, T.155N., R.27W.)

Depth (cm)

Vegetation: Consists of some ferns.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: T. Malterer on October 27, 1976.

Layer

Sand

Location: 2405 feet S and 1455 feet E of the NW corner of Sec. 13, T.155N., R.27W. (SW1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 13, T.155N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 12, 1976.

Layer	Depth (cm)
Hemic	0-76
Sandy clay loam	76+

Location: 1285 feet N and 1320 feet E of the SW corner of Sec. 13, T.155N., R.27W. (SE1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 13, T.155N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 12, 1976.

Layer	Depth (cm)	
Fibric	0- 30	
Hemic	30-101	
Sand	101+	

Location: 20 feet N and 1140 feet E of the SW corner of Sec. 13, T.155N., R.27W. (SE1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 13, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 95 percent; lush understory consists mostly of leatherleaf with Labrador tea and some swamp laurel, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 45 cm.

Depth To Water Table: 10 cm.

Described By: M. Domeier on August 12, 1976.

Layer	Depth (cm	
Fibric	0- 20	
Hemic	20-225	
Medium	225+	
eand		

Location: 1490 feet N and 2150 feet W of the SE corner of Sec. 13, T.155N., R.27W. (SE1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 13, T.155N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 12, 1976.

Layer	Depth (cm)
Fibric	0- 25
Hemic	25-127
Mineral	127+

Location: 100 feet S and 2125 feet E of the NW corner of Sec. 16, T.155N., R.27W.

(NW1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 16, T.155N., R.27W.)

Location: 1305 feet S and 135 feet E of the NW corner of Sec. 14, T.155N., R.27W. (SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 14, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; lush understory consists mostly of speckled alder with some Labrador tea; bog birch, shrubs, and ferns; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 1, 1976.

.ayer	Depth (cm)
eat	0-130
/ineral	130+
oil	

Location: 1930 feet N and 145 feet E of the SW corner of Sec. 14, T.155N., R.27W. (SW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 14, T.155N., R.27W.)

Vegetation: Scattered black spruce and tamarack; jush understory consists mostly of speckled alder with some Labrador tea, leatherleaf, shrubs, and ferns; ground cover consists mostly of Sphagnum mosses.

> Depth (cm) 0-193

193+

Microrellef: 60 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 1, 1976.

'	Vegetation: Not recorded.
	Microrelief: Not recorded.
	Depth To Water Table: Not recorded.
	Described By: R. Zarth on August 17, 1976.

Layer	Depth (cm)	
Hemic	0-140	
Sandy Ioam	140+	

Location: 100 feet S and 575 feet E of the NW corner of Sec. 16, T.155N., R.27W. (NE1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 16, T.155N., R.27W.) Vegetation: Cut over area consisting mostly of speckled alder.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth on August 17, 1976.

Depth (cm	
0-185 185+	

Hemic Sandy

Layer

Location: 1900 feet S and 25 feet E of the NW corner of Sec. 17, T.155N., R.27W. (SW1/4 of NW1/4 of SW1/4 of NW1/4. Sec. 17, T.155N., R.27W.)

Vegetation: Black spruce and tamarack crown cover of about 35 percent each; understory consists of Labrador tea and grasses with some paper birch; ground cover consists mostly of Sobagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on November 3, 1976.

Layer Depth (cm)
Hemic 0-127
Mineral 127+
soil

Location: 450 feet N and 75 feet E of the SW corner of Sec. 17, T.155N., R.27W. (NW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 17, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 95 percent; sparse understory consists of some Labrador tea and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on November 3, 1976.

Layer	Depth (cm)
Hemic	0-135
Sapric	135-175
Clay with calcareous pebbles	175+

Location: 2020 feet S and 100 feet E of the NW corner of Sec. 20, T.155N., R.27W. (SW1/4 of NW1/4 of NW1/4, Sec. 20, T.155N., R.27W.)

Vegetation: Scattered black spruce and paper birch; lush understory consists mostly of grasses and dogwood.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth on November 3, 1976.

Layer	Depth (cm)
Sapric	0-74
Clay Ioam	74+

Location: 700 feet N and 150 feet E of the SW corner of Sec. 20, T.155N., R.27W. (SW1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 20, T.155N., R.27W.)

Vegetation: Logged area, consists of scattered black spruce, tamarack, and paper birch; lush understc/y consists mostly of speckled alder with grasses and dogwood.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth on November 3, 1976.

Layer	Depth (cm)
Sapric	0-51
Clay	51+

Location: 2575 feet N and 115 feet E of the SW corner of Sec. 23, T.155N., R.27W. (NW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 23, T.155N., R.27W.)

Vegetation: Scattered tamarack and paper birch; lush understory consists mostly of speckled alder and grasses; ground cover consists mostly of Sphagnum masses

Microrelief: 40 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 1, 1976.

Layer	Depth (cm)	
Peat Mineral	0-179 179+	
soil		

Location: 585 feet N and 110 feet E of the SW corner of Sec. 23, T.155N., R.27W. (NW1/4 of SW1/4 of SW1/4, Sec. 23, T.155N., R.27W.)

Vegetation: Scattered tamarack and paper birch; lush understory consists mostly of speckled alder and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 40 cm.

Depth To Water Table: Not recorded.

Described By: T. Malterer on October 1, 1976.

Layer	Depth (cr	
Hemic Clay	0-45 45+	
loam		

Location: 1445 feet S and 2010 feet W of the NE corner of Sec. 24, T.155N., R.27W. (NW1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 24, T.155N., R.27W.)

Vegetation: Not recorded.
Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 12, 1976.

Layer Depth (cm)

Fibric 0- 8

Hemic 8-254

Clay 254+

loam

Location: 2520 feet N and 1480 feet E of the SW corner of Sec. 24, T.155N., R.27W. (NW1/4 of NW1/4 of NE1/4 of SW1/4, Sec. 24, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 65 percent; understory consists of Labrador tea and cotton grass with some bog birch and leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described by: M. Domeier on August 12, 1976.

Layer	Depth (cm)
Hemic	0-200
Sapric	200-205
Clay	205+
loam	

Location: 900 feet N and 1010 feet E of the SW corner of Sec. 24, T.155N., R.27W. (SW1/4 of NE1/4 of SW1/4 of SW1/4, Sec. 24, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of Labrador tea with some swamp laurel; ground cover consists mostly of Sphagnum mosses with some blueberry.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 12, 1976.

Layer	Depth (cm	
Hemic	0-54	
Sapric	54-97	
Clay	97+	
loam		

Location: 2550 feet N and 2030 feet W of the SE corner of Sec. 24, T.155N., R.27W. (NE1/4 of NW1/4 of NW1/4 of SE1/4, Sec. 24, T.155N., R.27W.)

Vegetation: Not recorded. Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M Domeier on August 12, 1976.

Laver Depth (cm) Fibric 0- 20 Hemic 20-183 Sand 183 +

Location: 175 feet S and 285 feet E of the NW corner of Sec. 25, T.155N., R.27W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 25, T.155N., R.27W.)

Vegetation: Scattered black spruce; understory consists of Labrador tea, bog birch, and speckled alder; ground cover consists of Sphagnum mosses. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 12, 1976.

Layer	Depth (cm)
Sapric Clay loam	0-33 33+

Peat 0-33 Mineral 33 +soil

Vegetation: Not recorded. Microrelief: Not recorded

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 19, 1976.

Layer

Location: 1025 feet N and 2050 feet E of the SW corner of Sec. 27, T.155N., R.27W. (NW1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 27, T.155N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 19, 1976.

Layer	Depth (cm)
Peat Mineral soil	0-61 61+

Location: 525 feet N and 2075 feet E of the SW corner of Sec. 27, T.155N., R.27W. (NW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 27, T.155N., R.27W.)

Vegetation: Not recorded. Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 19, 1976.

Layer	Depth (cm)
Peat Mineral soil	0-99 99+

Location: 50 feet N. and 2025 feet E of the SW corner of Sec. 27, T.155N., R.27W. (SW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 27, T.155N., R.27W.) Vegetation: Black spruce crown cover of about 60 percent; understory consists of

Location: 1415 feet N and 2000 feet E of the SW corner of Sec. 27, T.155N., R.27W.

Depth (cm)

(SW1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 27, T.155N., R.27W.)

Labrador tea; ground cover consists of Sphagnum and other mosses. Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 19, 1976.

Layer	Depth (cm)
Hemic	0-132
Sandy	132+
clay loam	

Location: 335 feet S and 120 feet W of the NE corner of Sec. 31, T.155N., B 27W. (NE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 31, T.155N., R.27W.)

Vegetation: Scattered black spruce, paper birch, and tamarack; lush understory consists mostly of Labrador tea with cotton grass; ground cover consists of Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 16, 1976.

Location: 575 feet S and 100 feet W of the NE corner of Sec. 31, T.155N., R.27W. (SE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 31, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent; understory consists of Labrador tea and grasses; ground cover consists of Sphagnum and feather mosses with some false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 16, 1976.

Layer	Depth (cm)	Layer	Depth (cm)
Hemic	0-30	Hemic	0-61
Coarse . sand	30+	Mineral soil	61+

Location: 860 feet S and 165 feet W of the NE corner of Sec. 31, T.155N., R.27W. (NE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 31, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 75 percent; lush understory consists mostly of Labrador tea with some leatherleaf and grasses; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 16, 1976.

Layer	Depth (cm)
Hemic	0-95
Sapric	95-99
Clay	99+
loam	

Vegetation: Black spruce crown cover of about 35 percent; understory consists of leatherleaf with some Labrador tea, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 9, 1976.

Layer	Depth (cm)
Fibric	0- 36
Sapric	36-200
Hemic	200-211
Sandy	211+
loam	

Location: 2350 feet S and 2140 feet E of the NW corner of Sec. 31, T.155N., R.27W. (SE1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 31, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of leatherleaf with some grasses, Labrador tea, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 9, 1976.

Depth (cm
0- 41
41-115
115-183
183+

41 Fibric 0-23
15 Hemic 23-229
83 Loam 229+
with

Microrelief: 15 cm

Depth To Water Table: At surface.

Location: 2620 feet S and 1930 feet W of the NE corner of Sec. 32, T.155N., R.27W. (SW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 32, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of Labrador tea and leatherleaf with some grasses; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 9, 1976.

Layer	Depth (cm)
Hemic Medium	0-71 71+
cand	

Location: 500 feet N and 2030 feet W of the SE corner of Sec. 32, T.155N., R.27W. (NE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 32, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists of Labrador tea and leatherleaf with some grasses and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 13, 1976.

Layer	Depth (cm)
Fibric	0- 15
Hemic	15-147
Loam	147+

Location: 500 feet N and 1400 feet W of the SE corner of Sec. 32, T.155N., R.27W. (NE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 32, T.155N., R.27W.)

Location: 1200 feet N and 875 feet E of the SW corner of Sec. 31, T.155N., R.27W.

Vegetation: Black spruce crown cover of about 35 percent; lush understory con-

sists of leatherleaf and grasses with some Labrador tea and swamp laurel;

(Gov. Lot 4, Sec. 31, T.155N., R.27W.)

Described By: R. Zarth on September 9, 1976.

ground cover consists mostly of Sphagnum mosses.

calcareous pebbles

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; lush understory consists of cotton grass and leatherleaf with some swamp laurel and Labrador tea; ground cover consists of Sphagnum mosses with some other mosses.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 16, 1976.

Layer	Depth (cm)
Hemic	0-117
Loam	117+

Location: 390 feet N and 920 feet W of the SE corner of Sec. 32, T.155N., R.27W. (NE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 32, T.155N., R.27W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of Labrador tea with some horsetail; ground cover consists of Sphagnum mosses with some feather mosses, cranberry, and false Solomon's seal.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 16, 1976.

Layer	Depth (cm)
Hemic	0-66
Loam	66+

Location: 500 feet N and 1150 feet W of the SE corner of Sec. 32, T.155N., R.27W. (NW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 32, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 90 percent; lush understory consists mostly of leatherleaf with grasses and some Labrador tea and swamp laurel; ground cover consists of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 16, 1976.

Layer	Depth (cm)
Peat	0-94
Loam	94+

Location: 600 feet S and 585 feet W of the NE corner of Sec. 34, T.155N., R.27W. (SW1/4 of NE1/4 of NE1/4 of NE1/4, Sec, 34, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of leatherleaf with some cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on August 19, 1976.

Layer	Depth (cm)
Fibric	0- 8
Hemic	8-102
Sandy	102+
clay	
loam	

Location: 605 feet S and 2010 feet E of the NW corner of Sec. 34. T.155N.. R.27W. (SW1/4 of NE1/4 of NE1/4 of NW1/4. Sec. 34. T.155N.. R.27W.)

Vegetation: Black spruce grown cover of about 60 percent, understory consists of Labrador tea and leatherleaf with some swamp laurel, ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on August 19, 1976.

Laver Depth (cm) Hemic 0-117 Sandy 117+ clay loam

Location: 1890 feet N and 950 feet E of the SW corner of Sec. 34, T.155N., R.27W. (NE1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 34, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of Labrador tea, leatherleaf, cotton grass, and sedges; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth on August 19, 1976.

Layer	Depth (cm)
Fibric Hemic Sandy	0- 15 15-137 137+
clay	107 1

Location: 1800 feet N and 710 feet W of the SE corner of Sec. 34, T.155N., R.27W. (NE1/4 of SW1/4 of NE1/4 of SE1/4, Sec. 34, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of cotton grass with leatherleaf and some Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: M. Domeier and R. Zarth on August 19, 1976.

Layer	Depth (cm)
Fibric	0- 51
Hemic	51-196
Clay	196+
loam	

Location: 1825 feet N and 1485 feet W of the SE corner of Sec. 34, T.155N., R.27W. (NE1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 34, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 40 percent; understory consists of some black spruce, sedges, Labrador tea, leatherleaf, swamp laurel, and bod rosemary; ground cover consists mostly of Sphagnum mosses with some feather mosses and snowberry.

Microrellef: 20 cm.

Depth To Water Table: 45 cm.

Described By: B. Sether and D. Haverkost on April 21, 1980.

Depth (cm)
0- 91
91-183
183-213
213+

Location: 555 feet S and 1730 feet W of the NE corner of Sec. 35, T.155N., R.27W. (SW1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 35, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists of Labrador tea and cotton grass with some leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: 5 cm.

Described By: M. Domeier and R. Zarth on August 18, 1976.

Layer	Depth (cm)
Hemic	0-130
Sandy	130+
clay	
loam	

Location: 2050 feet S and 1775 feet W of the NE corner of Sec. 35, T.155N., R.27W. (NW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 35, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 95 percent; lush understory consists mostly of Labrador tea with some cattail; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier and R. Zarth on August 18, 1976.

Layer	Depth (cm)
Sapric	0-46
Sandy	46+
clay	
loam	

Location: 575 feet S and 2225 feet F of the NW corner of Sec. 35, T.155N., R.27W. (SW1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 35, T.155N., R.27W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth and M. Domeier on August 18, 1976.

Depth (cm)
0-135
135+

Location: 1000 feet S and 2175 feet E of the NW corner of Sec. 35, T 155N, R 27W (SW1/4 of SE1/4 of NE1/4 of NW1/4, Sec. 35, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists of Labrador tea and leatherleaf with some cotton grass and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: 5 cm.

Described By: M. Domeier and R. Zarth on August 18, 1976.

Layer	Depth (cm)
Fibric	0-30
Hemic	30-86
Sandy	86+
clay	
loam	

Location: 2075 feet S and 2140 feet E of the NW corner of Sec. 35, T.155N., R.27W. (NW1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 35, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 65 percent; lush understory consists of Labrador tea and leatherleaf with some sedges and cotton grass; ground cover consists of some cranberry and false Solomon's seal.

Microrelief: 45 cm.

Depth To Water Table: 5 cm.

Described By: R. Zarth and M. Domeier on August 18, 1976.

Layer	Depth (cm)
Fibric	0- 51
Hemic	51-191
Sapric	191-203
Sandy	203+
clay	
loam	

Location: 1665 feet N and 2075 feet E of the SW corner of Sec. 35, T.155N., R.27W. (NW1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 35, T.155N., R.27W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of Labrador tea with some leatherleaf and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier and R. Zarth on August 16, 1976.

Layer	Depth (cm)
Hemic	0- 20
Fibric	20- 75
Hemic	75-196
Sandy	196+
clay	
loam	

Location: 2425 feet N and 75 feet E of the SW corner of Sec. 12, T.156N., R.27W.

Vegetation: Black spruce crown cover of about 40 percent with scattered northern

white cedar and tamarack; understory consists of Labrador tea with some

balsam fir, swamp laurel, grasses, and horsetail; ground cover consists mostly

of Sphagnum mosses with some other mosses, snowberry, bunchberry, lingen

Depth (cm)

0- 91

91-106

106 +

(NW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 12, T.156N., R.27W.)

berry, cranberry, false Solomon's seal, and twin flower.

Described By: R. Graber and N. Aaseng on November 13, 1979.

Laver

Hemic

Sapric

Clay

loam

Location: 1590 feet N and 1850 feet W of the SE corner of Sec. 35. T 155N . R 27W (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 35, T 155N, R.27W)

Vegetation: Black spruce crown cover of about 60 percent, lush understory consists mostly of Labrador tea with bog rosemary and cotton grass, ground cover consists mostly of Sphagnum mosses with some other mosses

Microrelief: 30 cm

Depth To Water Table: Not recorded.

Described By: M. Domeier and R. Zarth on August 18, 1976.

Layer	Depth (cm
Hemic	0-125
Sapric	125-140
Sandy	140+
clay	
loam	

Location: 1840 feet N and 125 feet E of the SW corner of Sec. 15, T.156N., R.27W.

Depth To Water Table: Not recorded.

Described By: D. Olson and D. Haverkost on November 13, 1979.

(NW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 15, T.156N., R.27W.)

Vegetation: Tamarack crown cover of about 90 percent.

Microrelief: Not recorded.

Layer	Depth (cm)
Sapric	0-335
Very	335+
fine	
sand	

Location: 100 feet S and 160 feet E of the NW corner of Sec. 17, T.156N., R.27W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 17, T.156N., R.27W.)

Vegetation: Black spruce crown cover of about 95 percent; understory consists of Labrador tea and leatherleaf with some grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 25 cm.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on November 13, 1979.

Layer	Depth (cm)
Hemic	0-208
Loam	208+

Location: 250 feet S and 100 feet E of the NW corner of Sec. 22, T.156N., R.27W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 22, T.156N., R.27W.)

Vegetation: Northern white cedar crown cover of about 40 percent with a paper birch crown cover of about 30 percent with scattered ash; understory consists of grasses with some balsam fir; ground cover consists of mosses.

Depth To Water Table: Not recorded.

Described By: R. Wakanabo on August 3, 1977.

Layer	Depth (cm)
Hemic	0-25
Sand	25+

Location: 350 feet N and 100 feet W of the SE corner of Sec. 9, T.156N., R.27W. (NE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 9, T.156N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered northern white cedar and tamarack; understory consists of some sedges, ferns, swamp laurel, and grasses; ground cover consists mostly of mosses with some Sphagnum mosses, snowberry, and pitcher plant.

Microrelief: 30 cm.

Denth To Water Table: At surface

Described By: R. Graber and N. Aaseng on November 13, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-213
Fine	213+
eand	

Location: 2025 feet N and 1190 feet W of the SE corner of Sec. 15, T.156N., R.27W. (SW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 15, T.156N., R.27W.)

Vegetation: Paper birch, black spruce, and tamarack crown cover of about 20 percent each with scattered northern white cedar; lush understory consists mostly of shrubs with grasses and some ferns; ground cover consists of some mosses. Microrelief: 5 cm.

Depth To Water Table: Below 60 cm.

Described By: D. Olson on August 3, 1977.

Layer	Depth (cm)
Hemic	0-40
Sand	40+

Location: 1250 feet S and 90 feet E of the NW corner of Sec. 22, T.156N., R.27W. (SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 22, T.156N., R.27W.)

Vegetation: Northern white cedar crown cover of about 40 percent with scattered birch; lush understory consists of grasses with some balsam fir, Labrador tea, and ferns; ground cover consists of some Sphagnum and other mosses.

Microrelief: 5 cm.

Depth To Water Table: Not recorded.

Described By: B. Wakanabo on August 3, 1977.

Layer	Depth (cm)
Hemic	0-34
Sand	34+
with	
pebbles	

Location: 2260 feet S and 100 feet E of the NW corner of Sec. 22. T.156N.. R.27W. (SW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 22. T.156N.. R.27W.)

Vegetation: Black spruce crown cover of about 30 percent with scattered birch; understory and ground cover not recorded.

Microrellet: 5 cm.

Depth To Water Table: Not recorded.

Described By: B. Wakanabo on August 2, 1977.

Layer Depth (cm)
Hemic 0-14
Coarse 14+
sand

Location: 1310 feet N and 100 feet E of the SW corner of Sec. 22, T.156N., R.27W. (SW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 22, T.156N., R.27W.)

Vegetation: Hazelnut crown cover of about 40 percent with scattered paper birch, black spruce, tamarack, and northern white cedar; understory consists of grasses with some ferns; ground cover consists of mosses with some false Solomon's seal.

Microrelief: 15 cm.

Depth To Water Table: 48 cm.

Described By: R. Wakanabo on August 2, 1977.

Layer	Depth (cm)
Hemic	0-36
Sand	36+

Location: 2375 feet S and 100 feet E of the NW corner of Sec. 27, T.156N., R.27W. (SW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 27, T.156N., R.27W.)

Vegetation: Tamarack crown cover of about 25 percent with a paper birch and cedar crown cover of about 20 percent each; lush understory consists mostly of shrubs with raspberry; ground cover not recorded.

Microrelief: 5 cm.

Depth To Water Table: Not visible.

Described By: D. Olson on August 2, 1977.

Layer	Depth (cm)
Hemic	0-41
Sandy clav	41-48
loam	
Sand	48+

Location: 90 feet N and 1030 feet W of the SE corner of Sec. 31, T.156N., R.27W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 31, T.156N., R.27W.)

Vegetation: Consists of stunted tamarack, leatherleaf, sedges, and bog birch.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier and C. Cameron on July 24, 1976.

Layer	Depth (cm)
Hemic Coarse sand	0-122 122+

Location: 75 feet N and 2055 feet W of the SE corner of Sec. 32, T.156N., R.27W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 32, T.156N., R.27W.)

Vegetation: Consists of willow and grasses.

Microrellef: Not recorded.

Deoth To Water Table: At surface.

Depth to water table: At surface

Described By: M. Domeier and C. Cameron on July 24, 1976.

Layer	Depth (cm)
Muck Sand	0-30 30+
Sanu	30 F

Location: 2400 feet S and 1835 feet E of the NW corner of Sec. 5, T.157N., R.27W. (SE1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 5, T.157N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, and cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 11, 1979.

.ayer	Depth (cm
ibric	0- 61
lemic	61-106
Silt	106-122
Clay	122+
oam	

Location: 730 feet N and 1825 feet E of the SW corner of Sec. 5, T.157N., R.27W. (SE1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 5, T.157N., R.27W.)

Vegetation: Black spruce crown cover of about 75 percent with scattered tamarack; understory consists of leatherleaf with some sedges, Labrador tea, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 11, 1979.

Location: 685 feet N and 2180 feet W of the SE corner of Sec. 5, T.157N., R.27W. (SE1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 5, T.157N., R.27W.)

Vegetation: Scattered black spruce and tamarack; understory consists of some Labrador tea, leatherleaf, sedges, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 11, 1979.

Location: 825 feet N and 590 feet W of the SE corner of Sec. 5, T.157N., R.27W. (SE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 5, T.157N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of Labrador tea, leatherleaf, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrelief: 15 cm.

Depth To Water Table: At surface,

Described By: D. Mellem and D. Haverkost on October 11, 1979.

Layer	Depth (cm)	
Fibric	0- 61	
Hemic	61-335	
Sapric	335-350	
Clay	350+	
loam		

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-305
Clay	305+
loam	

Layer	Depth (cm)
Peat	0-166
Clay	166+
loam	

berry.

Microrellef: 20 cm.

Solomon's seal.

Depth To Water Table: At surface.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Location: 390 feet S and 1975 feet E of the NW corner of Sec. 5, T.157N., R.27W. (Gov. Lot 3, Sec. 5, T.157N., R.27W.)

Vegetation: Black spruce crown cover of about 70 percent; understory consists of Labrador tea with some bog rosemary, leatherleaf, swamp laurel, and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 11, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-120
Silty	120+
clay	

Location: 2450 feet S and 125 feet W of the NE corner of Sec. 6, T.157N., R.27W. (SE1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 6, T.157N., R.27W.)

Vegetation: Black spruce crown cover of about 85 percent; understory consists of Labrador tea with some swamp laurel and sedges; ground cover consists mostly of Sphagnum mosses.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 10, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-165
Sandy	165+
clay	
loam	

Location: 1750 feet N and 2030 feet W of the SE corner of Sec. 8, T.157N., R.27W. (NE1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 8, T.157N., R.27W.)

Vegetation: Black spruce crown cover of about 65 percent; lush understory consists mostly of Labrador tea with some bog rosemary and sedges; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 11, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-342
Sapric	342-362
Sandy	362+
clay	
loam	

Location: 1700 feet S and 2620 feet W of the NE corner of Sec. 14, T.157N., R.27W.

Vegetation: Black spruce crown cover of about 50 percent; understory consists of

Labrador tea with some swamp laurel, bog rosemary, leatherleaf, and cotton grass; ground cover consists mostly of Sphagnum mosses with some false

(SW1/4 of NW1/4 of SW1/4 of NE1/4, Sec. 14, T.157N., R.27W.)

Described By: D. Meliem and D. Haverkost on October 8, 1979

Location: 1750 feet S and 375 feet E of the NW corner of Sec. 8, T.157N., R.27W.

Vegetation: Scattered black spruce; understory consists of sedges with some

Labrador tea, leatherleaf, swamp laurel, and bog rosemary; ground cover con-

sists mostly of Sphagnum mosses with some false Solomon's seal and cran-

(SE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 8, T.157N., R.27W.)

Described By: D. Mellem and D. Haverkost on October 10, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-286
Sapric	286-302
Sandy	302+
clay	
loam	

Location: 2565 feet N and 1100 feet E of the SW corner of Sec. 14, T.157N., R.27W. (NE1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 14, T.157N., R.27W.)

Vegetation: Black spruce crown cover of about 50 percent; understory consists of leatherleaf with some Labrador tea, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 8, 1979.

Layer Depth (cm)
Hemic 0-196
Silty 196+
clay

Layer Depth (cm)

Fibric 0- 60
Hemic 60-199
Silty 199+
clay with
calcareous pebbles

Location: 75 feet N and 75 feet W of the SE corner of Sec. 6, T.157N., R.27W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 6, T.157N., R.27W.)

Vegetation: Scattered black spruce; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, bog rosemary, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 10, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-420
Sapric	420-438
Sandy	438+
clay	
loam	

Location: 80 feet N and 2090 feet E of the SW corner of Sec. 13, T.157N., R.27W. (SW1/4 of SE1/4 of SE1/4 of SW1/4. Sec. 13, T.157N., R.27W.)

Vegetation: Black spruce crown cover of about 75 percent; understory consists mostly of Labrador tea with some sedges, leatherleaf, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, and cranberry.

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 8, 1979.

Layer	Depth (cm)
Fibric	0-120
Hemic	120-302
Silty clay	302+

Location: 2225 feet N and 660 feet W of the SE corner of Sec. 14, T.157N., R.27W. (SE1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 14, T.157N., R.27W.)

Vegetation: Scattered black spruce; sparse understory consists of some sedges, Labrador tea, bog rosemary, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and mushrooms.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 8, 1979.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-211
Silty clay	211+

Location: 500 feet S and 450 feet E of the NW corner of Sec. 17, T.157N., R.27W. (SE1/4 of NW1/4 of NW1/4 of NW1/4. Sec. 17. T.157N.. R.27W.)

Vegetation: Scattered black spruce and tamarack: understory consists of grasses and blue flag with some leatherleaf, speckled alder. Labrador tea, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 11, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-211
Silty	211+
clay	
loam	

Location: 1950 feet N and 1970 feet W of the SE corner of Sec. 17, T.157N., R.27W. (NE1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 17, T.157N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists mostly of Labrador tea with some leatherleaf and grasses: ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 11, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-181
Silty	181+
clay	

Location: 1600 feet N and 1970 feet W of the SE corner of Sec. 17, T.157N., R.27W. (SE1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 17, T.157N., R.27W.) Vegetation: Black spruce crown cover of about 60 percent; understory consists

mostly of Labrador tea with some grasses and leatherleaf; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 11, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-135
Silty	135+
clay	

Location: 1035 feet S and 75 feet E of the NW corner of Sec. 19, T.157N., R.27W. (Gov. Lot 1, Sec. 19, T.157N., R.27W.)

Vegetation: Northern white cedar crown cover of about 45 percent; lush understory consists mostly of grasses with speckled alder.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and R. Graber on November 13, 1979.

Layer	Depth (cm)
Hemic	0-167
Fine	167+
sand	

Location: 1900 feet S and 500 feet E of the NW corner of Sec. 23, T.157N., R.27W. (SE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 23, T.157N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 8, 1979.

Layer	Depth (cm)
Fibric Hemic Coarse sandy clay	0- 60 60-180 180+

Location: 575 feet N and 455 feet E of the SW corner of Sec. 23, T.157N., R.27W. (NW1/4 of SW1/4 of SW1/4, Sec. 23, T.157N., R.27W.)

Vegetation: Black spruce crown cover of about 75 percent; understory consists mostly of Labrador tea with some sedges and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cran-

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 8, 1979.

Layer	Depth (cm
Fibric	0-151
Hemic	151-347
Silty	347+
clay	

Location: 1880 feet S and 2315 feet E of the NW corner of Sec. 24, T.157N., R.27W. (SW1/4 of NE1/4 of SE1/4 of NW1/4, Sec. 24, T.157N., R.27W.)

Vegetation: Scattered black spruce; understory consists of some Labrador tea, bog rosemary, sedges, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry. Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 8, 1979.

Location: 725 feet N and 1975 feet E of the SW corner of Sec. 24, T.157N., B.27W. (SE1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 24, T.157N., R.27W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges with some leatherleaf, swamp laurel, bog rosemary, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 4, 1979.

Layer	Depth (cm)	Layer	Depth (cm)
Fibric	0-120	Fibric	0-180
Hemic	120-286	Hemic	180-453
Silty	286+	Sapric	453-484
clay		iity	484+
		clay	

Location: 625 feet S and 1055 feet W of the NE corner of Sec. 25, T.157N., R 27W. (SW1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 25, T.157N., R.27W.)

Vegetation: Scattered black spruce; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 4, 1979.

Layer	Depth (cm)
Fibric	0-210
Hemic	210-363
Sapric	363-393
Silty	393+
clay	

Location: 525 feet S and 2335 feet E of the NW corner of Sec. 25, T.157N., R.27W. (SE1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 25, T.157N., R.27W.)

Vegetation: Scattered black spruce and tamarack; understory consists of sedges with some Labrador tea, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 4, 1979.

Layer	Depth (cm)	
Fibric	0-181	
Hemic	181-453	
Sapric	453-483	
Silty	483+	
clay		

Location: 15 feet S and 410 feet E of the NW corner of Sec. 26. T.157N., R.27W.

Vegetation: Black spruce crown cover of about 70 percent with scattered

tamarack; lush understory consists mostly of sedges with some bog birch and

bog rosemary; ground cover consists mostly of Sphagnum mosses with some

(NE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 26, T.157N., R.27W.)

Described By: D. Mellem and D. Haverkost on October 8, 1979.

pitcher plant and faise Solomon's seal.

Depth To Water Table: At surface.

Location: 350 feet N and 2175 feet E of the SW corner of Sec. 25, T.157N., R.27W. (NW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 25, T.157N., R.27W.)

Vegetation: Scattered black spruce and tamarack: understory consists mostly of sedges and Phragmites with some leatherleaf and bog rosemary; ground cover consists of some Sphagnum mosses and pitcher plant.

Microrellef: 10 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on October 13, 1979.

Layer	Depth (cm)	
Fibric	0-152	
Hemic	152-289	
Loam	289+	

Location: 700 feet N and 1655 feet W of the SE corner of Sec. 26, T.157N., R.27W. (NW1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 26, T.157N., R.27W.)

Vegetation: Scattered black spruce and tamarack: lush understory consists mostly of sedges with some bog rosemary and Labrador tea; ground cover consists mostly of Sphagnum mosses.

Depth (cm)

0- 91

91-320

320+

Microrellef: Not recorded.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on October 13, 1979.

Laver

Fibric

Hemic

Clay

loam

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-151
Fibric	151-211
Hemic	211-317
Sapric	317-332
Medium	332+
and	
coarse	
sandy	
clay	

Location: 740 feet S and 1160 feet E of the NW corner of Sec. 1, T,158N., R,27W.

Vegetation: Scattered black spruce; sparse understory consists of Labrador tea, leatherleaf, swamp laurel, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrellef: 45 cm.

Microrellef: 15 cm.

Depth To Water Table: At surface.

(Gov. Lot 4, Sec. 1, T.158N., R.27W.)

Described By: D. Mellem and D. Haverkost on September 13, 1979.

Layer	Depth (cm)	
Fibric	0- 30	
Hemic	30-250	
Sapric	250-263	
Silty	263+	
clav		

Location: 1875 feet N and 2315 feet E of the SW corner of Sec. 5, T.158N., R.27W.

(NE1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 5, T.158N., R.27W.) Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf with some swamp laurel, cotton grass, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, lingen berry, false Solomon's seal and lichens.

Microrellef: 40 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Haverkost on November 2, 1979.

Layer	Depth (cm)	
Fibric	0-240	
Hemic	240-360	
Sapric	360-450	
Silty	450+	
clav		

Location: 865 feet S and 2000 feet E of the NW corner of Sec. 26, T.157N., R.27W. (NW1/4 of SE1/4 of NE1/4 of NW1/4, Sec. 26, T.157N., B.27W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some leatherleaf, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses with some snowberry.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 8, 1979.

Layer	Depth (cm)	
Fibric	0-120	
Hemic	120-210	
Fibric	210-270	
Hemic	270-484	
Silty	484+	
clay		

Location: 700 feet S and 2600 feet W of the NE corner of Sec. 27, T.157N., R.27W. (NW1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 27, T.157N., R.27W.)

Vegetation: Scattered black spruce and northern white cedar; understory consists of grasses; ground cover consists of some Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and R. Graber on November 13, 1979.

Layer	Depth (cm)	
Fibric	0-106	
Hemic	106-396	
Sandy	396+	
loam		

Location: 800 feet N and 750 feet E of the SW corner of Sec. 6, T.158N., R.27W. (Gov. Lot 7, Sec. 6, T.158N., R.27W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of leatherleaf with some swamp laurel, sedges, bog rosemary, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Haverkost on November 2, 1979.

Layer	Depth (cm)	
Fibric	0- 30	
Hemic	30-120	
Sapric	120-185	
Silty	185+	
clay		

Location: 550 feet N and 350 feet E of the SW corner of Sec. 10, T.158N., R.27W. (NE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 10, T.158N., R.27W.)

Vegetation: Scattered tamarack; understory consists of some leatherleaf, swamp laurel, sedges, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrellef: 30 cm.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: N. Aaseng and R. Graber on November 13, 1979.

Layer Depth (cm) Fibric 0-137 137-335 Hemic Clay 335+

Location: 360 feet N and 250 feet W of the SE corner of Sec. 14, T.158N., R.27W.

Vegetation: Scattered tamarack and black spruce; understory consists of

leatherleaf with some sedges and swamp laurel; ground cover consists mostly

Depth (cm)

0-183

183-213

213+

(NE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 14, T.158N., R.27W.)

Described By: D. Olson and D. Haverkost on November 13, 1979.

of Sphagnum mosses with some cranberry.

Laver

Fibric

Hemic

Sand

Location: 1770 feet N and 690 feet E of the SW corner of Sec. 12, T.158N., R.27W. (NW1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 12, T.158N., R.27W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with leatherleaf and some swamp laurel.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on November 13, 1979.

Layer	Depth (cm)	
Fibric	0- 91	
Hemic	91-259	
Silty	259+	
clay		

Location: 175 feet N and 175 feet E of the SW corner of Sec. 16, T.158N., R.27W.

Vegetation: Lush understory consists mostly of sedges with leatherleaf and some

Microrelief: 20 cm.

Depth To Water Table: At surface.

Layer	Depth (cm)	
Hemic	0-193	
Clay	193+	

Location: 660 feet S and 130 feet W of the NE corner of Sec. 13, T.158N., R.27W. (SE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 13, T.158N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some sedges, swamp laurel, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)	
Fibric	0-238	
Hemic	238-335	
Silty	335+	
ala.		

Location: 1525 feet S and 2060 feet E of the NW corner of Sec. 17, T.158N., R.27W.

Vegetation: Scattered black spruce; lush understory consists mostly of sedges and

cotton grass; ground cover consists of some Sphagnum and feather mosses.

(NW1/4 of NE1/4 of SE1/4 of NW1/4, Sec. 17, T.158N., R.27W.)

Described By: D. Olson and R. Graber on November 2, 1979.

clav

(SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 16, T.158N., R.27W.)

grasses.

Described By: D. Olson and D. Haverkost on November 13, 1979.

Depth (cm)	Layer	Depth (cm)
0-193	Fibric	0- 91
93+	Hemic	91-122
	Fibric	122-183
	Hemic	183-442
	Silty	442+

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Location: 640 feet S and 1125 feet W of the NE corner of Sec. 18, T.158N., R.27W. (SW1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 18, T.158N., R.27W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges and grasses; ground cover consists mostly of cranberry with some Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: At surface.

Described By: D. Olson and R. Graber on November 2, 1979.

Laver

Fibric Hemic Silty clay

Location: 1130 feet S and 125 feet E of the NW corner of Sec. 29, T.158N., R.27W. (SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 29, T.158N., R.27W.)

Vegetation: Tamarack crown cover of about 60 percent with scattered black spruce; lush understory consists mostly of Labrador tea, sedges, and cotton grass with some leatherleaf; ground cover consists of Pulesterum mosses. Microrellef: 20 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and R. Graber on November 2, 1979.

Depth (cm)	Layer	Depth (cm)
0-152	Hemic	0-111
152-442 442+	Silty clay	111+

Location: 925 feet N and 25 feet W of the SE corner of Sec. 36, T. 158N., R.27W. (SE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 36, T.158N., R.27W.)

Vegetation: Aspen crown cover of about 30 percent with scattered black spruce, balsam fir, and paper birch; understory consists of grasses and speckled alder: ground cover consists of mosses including some Sphagnum mosses. Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: R. Graber and N. Aaseng on November 13, 1979.

Layer	Depth (cm)	
Silty clay	0+	

Microrellef: 40 cm.

Dooth To Water Table: At surface

Location: 100 feet S and 2350 feet E of the NW corner of Sec. 6, T.159N., R.27W. (Gov. Lot 3, Sec. 6, T.159N., R.27W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered aspen and balsam fir; understory consists of grasses; ground cover consists of some Sphagnum mosess.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem on November 13, 1979.

Layer Depth (cm)
Hemic 0-183
Silty 183+
clay

Location: 1890 feet N and 1940 feet E of the SW corner of Sec. 20, T.159N., R.27W. (NE1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 20, T.159N., R.27W.)

Vegetation: Black spruce crown cover of about 70 percent with about 25 percent

and swamp laurel; ground cover consists of feather mosses.

Described By: D. Olson and R. Graber on November 2, 1979.

Layer

Hemic

crown cover of tamarack; understory consists of grasses with some leatherleaf

Depth (cm)

0-350 350+ Location: 1075 feet S and 60 feet W of the NE corner of Sec. 11, T.159N., R.27W. (SE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 11, T.159N., R.27W.)

Vegetation: Aspen crown cover of about 50 percent with scattered paper birch and balsam fir; understory consists of speckled alder and grasses with some labrador tea.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem on November 16, 1979.

Layer	Depth (cm	
Fibric	0- 61	
Hemic	61-122	
Clay	122+	

Location: 1550 feet N and 150 feet E of the SW corner of Sec. 21, T.159N., R.27W. (SW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 21, T.159N., R.27W.)

Vegetation: Lush understory consists mostly of sedges with some bog birch and willow; ground cover consists of some mosses.

Microrelief: 5 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on November 14, 1979.

 Layer
 Depth (cm)

 Fibric
 0-122

 Hemic
 122-310

 Clay
 310+

Location: 1615 feet N and 205 feet E of the SW corner of Sec. 22, T.159N., R.27W. (SW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 22, T.159N., R.27W.)

Location: 2510 feet N and 1115 feet W of the SE corner of Sec. 12, T.159N., R.27W.

Vegetation: Scattered black spruce and tamarack; understory consists of some

bog birch, sedges, swamp laurel, leatherleaf, and Labrador tea; ground cover

consists mostly of Sphagnum mosses with some other mosses and cranberry.

Depth (cm)

0-335

335+

(NW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 12, T.159N., R.27W.)

Described By: D. Olson and N. Aaseng on November 14, 1979.

Laver

Hemic

Sandy

clav

Vegetation: Scattered tamarack, black spruce, and paper birch; sparse understory consists of some balsam fir, willow, Labrador tea, sedges, grasses, and ferns; ground cover consists of Sphagnum and other mosses.

Microrelief: 15 cm.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: D. Oison and N. Aaseng on November 14, 1979.

Layer	Depth (cm	
Hemic	0-177	
Silty	177+	
clay		
loam		

Location: 2510 feet S and 2250 feet W of the NE corner of Sec. 24, T.159N., R.27W. (SE1/4 of SW1/4 of SW1/4 of NE1/4, Sec. 24, T.159N., R.27W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some speckled alder, Labrador tea, leatherleaf, swamp laurel, bog rosemary, grasses, and cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Location: 2450 feet S and 800 feet E of the NW corner of Sec. 32, T.159N., R.27W. (SW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 32, T.159N., R.27W.)

Vegetation: Scattered tamarack and black spruce; understory consists mostly of leatherleaf with sedges and some bog birch, bog rosemary, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 10 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Haverkost on November 2, 1979.

Layer	Depth (cm)	
Fibric	0- 90	
Hemic	90-105	
Silty	105+	
clav		

Location: 1675 feet S and 1100 feet E of the NW corner of Sec. 33, T.159N., R.27W. (SE1/4 of NE1/4 of SW 1/4 of NW1/4, Sec. 33, T.159N., R.27W.)

Vegetation: Lush understory consists mostly of bog birch with sedges and some bog rosemary, leatherleaf, tamarack, and aster; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 5 cm.

Depth To Water Table: At surface.

Described By: D. Oison and N. Aaseng on November 14, 1979.

Layer	Depth (cm)	
Hemic Clay	0-228 228+	
loam		

Location: 650 feet N and 2000 feet E of the SW corner of Sec. 35, T.159N., R.27W. (NW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 35, T.159N., R.27W.)

Vegetation: Black spruce crown cover of about 40 percent; understory consists of Labrador tea, leatherleaf, swamp laurel, and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 13, 1979.

Layer Depth (cm)
Hemic 0-120
Sandy 120+
clay

Location: 590 feet N and 715 feet W of the SE corner of Sec. 35, T.159N., R.27W. (NE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 35, T.159N., R:27W.)

Vegetation: Scattered black spruce; sparse understory consists of some Labrador tea, swamp laurel, and sedges; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface

Described By: D. Mellem and D. Haverkost on September 13, 1979.

Layer	Depth (cm)
Fibric Hemic Sapric Silty clay	0- 90 90-212 212-215 215+

Location: 1960 feet S and 1400 feet W of the NE corner of Sec. 36, T.159N., R.27W. (SE1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 36, T.159N., R.27W.)

Vegetation: Scattered tamarack and black spruce; sparse understory consists of some leatherleaf, swamp laurel, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses.

Microrellef: 15 cm.

Depth To Water Table: Near surface.

Described By: R. Graber and D. Haverkost on November 14, 1979.

Layer	Depth (cm)	
Fibric	0-152	
Hemic	152-274	
Clay	274+	
loam		

Location: 180 feet N and 1390 feet E of the SW corner of Sec. 36, T.159N., R.27W. (SW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 36, T.159N., R.27W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some leatherleaf, bog rosemary, and horsetail; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 13, 1979.

Layer	Depth (cm)	
Fibric	0- 30	
Hemic	30-150	
Silty	150+	
clay		

Location: 425 feet S and 690 feet E of the NW corner of Sec. 6, T.151N., R.28W. (Gov. Lot 3, Sec. 6, T.151N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some leatherleaf, swamp laurel, bog rosemary, and iris; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and N. Aaseng on December 10, 1979.

Layer	Depth (cm)
Hemic Coarse and medium	0-183 183+

Location: 75 feet S and 410 feet E of the NW corner of Sec. 7, T.151N., R.28W. (Gov. Lot 1, Sec. 7, T.151N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of leatherleaf with some grasses, swamp laurel, and Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrellef: Negligible.

Depth To Water Table: Not visible.

Described By: D. Mellem and D. Haverkost on December 10, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-152 152+

Location: 1505 feet S and 600 feet E of the NW corner of Sec. 27, T.151N., R.28W. (NE1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 27, T.151N., R.28W.)

Vegetation: Scattered black spruce and northern white cedar; lush understory consists of grasses and raspberry with some alder, dogwood, and bog birch; ground cover consists of some Sphagnum mosses.

Depth (cm)

0-152

152 +

Microrellef: 10 cm.

Depth To Water Table: 15 cm.

Described By: H. Mooers and D. Haverkost on May 20, 1980.

Layer

Hemic

Silt

loam

Location: 70 feet N and 2325 feet E of the SW corner of Sec. 13, T.152N., R.28W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 13, T.152N., R.28W.)

Vegetation: Northern white cedar crown cover of about 90 percent with scattered black spruce; lush understory consists mostly of speckled alder with some grasses.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on December 10, 1979.

Layer	Depth (cm)
Hemic	0-183
Fibric	183-198
Limnic	198-213
Silty	213+
clay	
with	
nebbles	

Location: 1100 feet S and 2000 feet W of the NE corner of Sec. 1, T.153N., R.28W. (Gov. Lot 2, Sec. 1, T.153N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; understory consists of leatherleaf with some Labrador tea, bog rosemary, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some snowberry and false Solomon's seal.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: H. Mooers and D. Haverkost on April 3, 1980.

Layer	Depth (cm)
Hemic	0-138
Loamy	138+
sand	

Location: 625 feet S and 530 feet W of the NE corner of Sec. 25, T.153N., R.28W (SW1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 25, T.153N., R.28W.)

Vegetation: Tamarack crown cover of about 35 percent; understory consists mostly of speckled alder with some Labracur tea and fern.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox on October 27, 1976.

 Layer
 Depth (cm)

 Hemic Sand
 0-157

 157+
 157+

Location: 1650 feet S and 625 feet W of the NE corner of Sec. 25, T.153N., R.28W. (NW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 25, T.153N., R.28W.)

Vegetation: Not recorded.
Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on October 27, 1976.

Layer	Depth (cm)
Peat	0-122
Mineral soil	122+
SOII	

Location: 1600 feet S and 500 feet W of the NE corner of Sec. 1, T.154N., R.28W. (NW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 1, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of leatherleaf with some swamp laurel, Labrador tea, and cotton grass; ground cover consists most; of Sphagnum mosses with some false Solomon's seal.

Micrografie: 15 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 9, 1976.

Layer	Depth (cm)
Fibric	0- 13
Hemic	13-198
Loam	198+

Location: 100 feet N and 675 feet W of the SE corner of Sec. 1, T.154N., R.28W. (SW1/4 of SE1/4 of SE1/4 of SE1/4. Sec. 1, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; understory consists of leatherleaf with some Labrador tea and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 9, 1976.

Layer	Depth (cm)
Hemic	0-183
Clay	183+
loam	

Location: 125 feet N and 1600 feet E of the SW corner of Sec. 2, T.154N., R.28W. (SW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 2, T.154N., R.28W.)

Vegetation: Scattered black spruce; lush understory consists mostly of leatherleaf with sedges and some Labrador tea and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth and R. Fox on September 16, 1976.

 Layer
 Depth (cm)

 Fibric
 0- 51

 Hemic
 51-122

 Loam
 122+

Location: 2375 feet S and 1660 feet E of the NW corner of Sec. 2, T.154N., R.28W.

Vegetation: Black spruce crown cover of about 40 percent; lush understory consists of leatherleaf, sedges, and cotton grass with some Labrador tea; ground

(SE1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 2, T.154N., R.28W.)

cover consists mostly of Sphagnum mosses.

Described By: R. Fox and R. Zarth on September 16, 1976.

Depth To Water Table: Not recorded.

Microrellef: 30 cm.

 Layer
 Depth (cm)

 Fibric
 0- 20

 Hemic
 20-140

 Loam
 146+

Location: 1060 feet N and 2425 feet W of the SE corner of Sec. 34, T.153N., R.28W. (NW1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 34, T.153N., R.28W.)

Vegetation: Tamarack crown cover of about 30 percent; understory consists mostly of grasses with some black spruce, northern white cedar, alder, sedges, cattail, and bog rosemary; ground cover consists of some Sphagnum mosses and pitcher plant.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and S. Nelson on April 3, 1980.

Layer	Depth (cm
Fibric .	0-122
Hemic	122-223
Fine	223+
sand	

Location: 230 feet S and 625 feet E of the NW corner of Sec. 1, T.154N., R.28W.

(Gov. Lot 4, Sec. 1, T.154N., R.28W.)

Vegetation: Not recorded.
Microrelief: Not recorded.

Depth To Water Table: Not recorded

Described By: R. Fox on September 16, 1976.

Layer	th (cm) عرب
Fibric	0- 69
Hemic	69-203
Clay	203+
loam	

Location: 2410 feet N and 175 feet W of the SE corner of Sec. 2, T.154N., R.28W. (NE1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 2, T.154N., R.28W.)

Vegetation: Scattered tamarack; understory consists mostly of sedges and cotton grass with some black spruce; ground cover consists of some Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on September 15, 1976.

Layer	Depth (cm)
Fibric	0- 51
Hemic	51-163
Loam	163+

Location: 175 feet N and 250 feet W of the SE corner of Sec. 2, T.154N., R.28W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 2, T.154N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of leatherleaf with Labrador tea and some bog rosemary, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded. Described On: September 13, 1976.

Layer	Depth (cr
Fibric	0- 23
Hemic	23-155
Loam	155+

Location: 900 feet S and 1700 feet E of the NW corner of Sec. 2, T.154N., R.28W. (Gov. Lot 3, Sec. 2, T.154N., R.28W.)

Vegetation: Not recorded.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on September 16, 1976.

Layer	Depth (cm)
Fibric	0- 33
Hemic	33-142
Sandy	142+
loam	

Location: 2370 feet S and 1075 feet W of the NE corner of Sec. 3, T.154N., R.28W. (NW1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 3, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory con-

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of leatherleaf with Labrador tea and some sedges; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth and R. Fox on September 16, 1976.

Layer	Depth (cm
Fibric	0- 50
Hemic	50-173
Clay	173+
loam	

Location: 85 feet N and 25 feet E of the SW corner of Sec. 3, T.154N., R.28W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 3, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of Labrador tea with some leatherleaf and sedges.

Microrellet: 45 cm

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on September 16, 1976.

Layer	Depth (cm)
Fibric	0- 25
Hemic	25-180
Sandy	180+
loam	

Location: 1125 feet N and 1390 feet E of the SW corner of Sec. 3, T.154N., R.28W. (NW1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 3, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; lush understory consists mostly of sedges with Labrador tea and leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: R. Zarth and R. Fox on September 16, 1976.

Layer	Depth (cm)
Fibric	0-15
Hemic	15-86
Fine	86+
sand	

Location: 75 feet N and 1440 feet E of the SW corner of Sec. 3, T.154N., R.28W. (SW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 3, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists of sedges with some Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on September 16, 1976.

Layer	Depth (cm)
Fibric	0- 18
Hemic	18-231
Sandy	231+
loam	

Location: 75 feet N and 1150 feet W of the SE corner of Sec. 3, T.154N., R.28W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 3, T.154N., R.28W.)

Vegetation: Scattered black spruce; lush understory consists of sedges and leatherleaf with some bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 3 cm.

Described By: R. Zarth and R. Fox on September 16, 1976.

Layer

Fibric Hemic Sandy Ioam Location: 860 feet S and 1050 feet W of the NE corner of Sec. 3, T.154N., R.28W. (Gov. Lot 1, Sec. 3, T.154N., R.28W.)

Vegetation: Scattered black spruce; lush understory consists of Labrador tea, leatherleaf, and cotton grass with some swamp laurel and bog birch: ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on September 16, 1976.

Depth (cm)	Layer	Depth (cm)
0- 60	Fibric	0- 20
60-208	Hemic	20-102
208+	Sand	102+

Location: 75 feet N and 1105 feet E of the SW corner of Sec. 4, T.154N., R.28W. (SE1/4 of SE1/4 of SW1/4, Sec. 4, T.154N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of leatherleaf with bog birch and Labrador tea with some swamp laurel, willow, sedges, cotton grass, and bog rosemary; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on September 27, 1976.

Layer	Depth (cm)
Hemic	0-155
Sandy	155+
loam	

Location: 1435 feet N and 1790 feet E of the SW corner of Sec. 9. T 154N R 28W (SE1/4 of SW1/4 of NE1/4 of SW1/4. Sec. 9. T 154N, R 28W)

Vegetation: Black spruce crown cover of about 35 percent: understory consists of sedges with some leatherleaf, Labrador tea, and pitcher plant: ground cover consists mostly of Schagnum mosses with some other mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on September 24, 1976.

Layer Depth (cm)
Fibric 0- 84
Hemic 84-318
Clay 318+
loam

Location: 1410 feet N and 80 feet W of the SE corner of Sec. 9, T.154N., R.28W. (SE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 9, T.154N., R.28W.)

Vegetation: Lush understory consists mostly of bog rosemary and leatherleaf with stunted black spruce, tamarack, willow, and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrellef: 75 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on September 24, 1976.

Layer	Depth (cm)
Fibric	0- 51
Sapric	51-124
Hemic	124-318
Mineral	318+
soil	

Location: 2390 feet S and 1130 feet W of the NE corner of Sec. 10, T.154N., R.28W. (SW1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 10, T.154N., R.28W.)

Vegetation: Scattered black spruce; lush understory consists of Labrador tea, leatherleaf, and cotton grass with some swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox on September 20, 1976.

Layer	Depth (cm)
Fibric	0- 76
Hemic	76-411
Mineral	411+
soil	

Location: 2350 feet S and 225 feet E of the NW corner of Sec. 11, T.154N., R 28W (SW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 11, T.154N., R.28W.)

Vegetation: Scattered black spruce: lush understory consists mostly of cotton grass with some leatherleaf and swamp laurel: ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 21, 1976.

Layer	Depth (cm
Fibric	0- 61
Sapric	61-140
Hemic	140-262
Sandy	262+
loam	

Location: 2300 feet S and 1625 feet E of the NW corner of Sec. 11, T.154N., R.28W. (SW1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 11, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 50 percent; lush understory consists of cotton grass and grasses with some Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 5 cm.

Described By: M. Domeier on September 21, 1976.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-305
Sand	305+
over	
clay	

Location: 550 feet N and 1500 feet E of the SW corner of Sec. 11, T.154N., R.28W. (NW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 11, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 70 percent; lush understory consists mostly of Labrador tea with cotton grass and grasses with some leatherleaf and swamp laurel; ground cover consists of Sphagnum mosses and cranberry with some other mosses and lichens.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 21, 1976.

Layer	Depth (cm)
Fibric	0-228
Hemic	228-445
Clay Ioam	445+
Hemic Clay	228-445

Location: 1540 feet N and 2465 feet W of the SE corner of Sec. 11, T.154N., R.28W. (NW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 11, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of grasses and leatherleaf with swamp laurel and some Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 21, 1976.

Layer	Depth (cm)
Fibric	0- 51
Sapric	51- 61
Hemic	61-432
Clay	432+
loam	

Location: 875 feet S and 280 feet W of the NE corner of Sec. 12, T.154N., R.28W. (NE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 12, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 50 percent with scattered tamarack; lush understory consists mostly of leatherleaf with grasses and some Labrador tea and swamp laurel; ground cover consists mostly of Sphagnum mosses with some feather mosses.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 28, 1976.

Layer	Depth (cm)
Fibric	0- 23
Hemic	23-155
Clay	155+
loam	

Location: 2340 feet S and 90 feet E of the NW corner of Sec. 12, T.154N., R.28W. (SW1/4 of SW1/4 of SW1/4, Sec. 12, T.154N., R.28W.)

Vegetation: Lush understory consists mostly of sedges and leatherleaf with some black spruce and tamarack; ground cover consists mostly of Sphagnum mosses.

Microrellef: 60 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox on September 30, 1976.

Layer	Depth (cm)
Fibric	0- 58
-lemic	58-236
Mineral	236+
soil	

Location: 2385 feet S and 2410 feet E of the NW corner of Sec. 12, T.154N R 28W (SE1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 12. T.154N., R.28W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges and leatherleaf with some tamarack; ground cover consists mostly of Sphagnum mosses

Depth (cm)

0- 13

13-137

137+

Microrellef: Not recorded.

Depth To Water Table: Not recorded. Described On: September 30, 1976.

Laver

Fibric

Hemic

Loam

Microrelief: Not recorded. Depth To Water Table: Not recorded. Described By: R. Fox on September 30, 1976.

Layer	Depth (cm)
Fibric	0- 66
Hemic	66-408
Loam	408+

Location: 925 feet S and 2615 feet W of the NE corner of Sec. 13, T 154N, B 28W. (NW1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 13, T.154N., R.28W.)

Vegetation: Scattered tamarack; lush understory consists mostly of sedges with

leatherleaf; ground cover consists mostly of Sphagnum mosses.

Location: 700 feet S and 1430 feet W of the NE corner of Sec. 13, T.154N., R.28W. (NE1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 13, T.154N., R.28W.)

Vegetation: Scattered black spruce and tamarack; understory consists mostly of leatherleaf with bog rosemary.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on September 30, 1976.

Layer	Depth (cm)
Peat	0-107
Mineral	107+
eoil	

Location: 890 feet S and 1575 feet W of the NE corner of Sec. 13, T.154N., R.28W. (NW1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 13, T.154N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges and leatherleaf with some bog rosemary; ground cover consists mostly of Sphagnum mosses.

Depth (cm)

0- 60 60-240

240-244

244 +

Microrelief: 45 cm

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 30, 1976.

Laver

Fibric

Hemic

Sapric

Sand

Location: 815 feet N and 875 feet E of the SW corner of Sec. 13, T.154N., R.28W. (SW1/4 of NE1/4 of SW1/4 of SW1/4, Sec. 13, T.154N., R.28W.)

Vegetation: Scattered tamarack and black spruce; lush understory consists mostly of bog rosemary with some sedges and bog birch; ground cover consists mostly of Sphagnum mosses.

Microrelief: 60 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox in September . 1976.

Layer	Depth (cm)	
Fibric	0- 20	
Hemic	20-381	
Silt	381+	

Location: 825 feet S and 1050 feet W of the NE corner of Sec. 14, T.154N., R.28W. (NW1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 14, T.154N., R.28W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges and leatherleaf with some tamarack; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: At surface.

Described By: R. Zarth on September 30, 1976.

Layer	Depth (cm)
Fibric	0- 64
Hemic	64-500
Sapric	500-511
Loam	511+

Location: 1875 feet S and 1475 feet E of the NW corner of Sec. 14, T 154N . R 28W (SW1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 14, T.154N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent, lush understory consists of Labrador tea, swamp laurel, and cotton grass; ground cover consists of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 5 cm.

Described By: M. Domeier on September 21, 1976.

Location: 125 feet S and 1200 feet W of the NE corner of Sec. 15, T.154N., R.28W. (NW1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 15, T.154N., R.28W.)

Vegetation: Scattered black spruce: Jush understory consists mostly of leatherleaf with Labrador tea and swamp laurel with some cotton grass: ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 20, 1976.

	A Section of the sect		
Layer	Depth (cm)	Layer	Depth (cm)
Fibric	0-290	Fibric	0-127
Hemic	290-549	Hemic	127 ·
Clay	549+	Bottom	_
loam		unknown	

Location: 400 feet S and 1410 feet E of the NW corner of Sec. 15, T.154N., R.28W. (SW1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 15, T.154N., R.28W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges with leatherleaf and some tamarack; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 21, 1976.

Layer	Depth (cm)	
Fibric	0-117	
Hemic	117-366	
Loarn	366+	

Location: 1850 feet S and 1425 feet E of the NW corner of Sec. 15, T.154N., R.28W. (SW1/4 of NW1/4 of SE1/4 of NW1/4, Sec. 15, T.154N., R.28W.)

Vegetation: Not recorded. Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: R. Fox and R. Zarth on September 21, 1976.

 Layer
 Depth (cm)

 Fibric
 0- 89

 Hemic
 89-564

 Silt
 564+

 loam

Location: 75 feet N and 1510 feet W of the SE corner of Sec. 15. T.154N., R.28W/.

Vegetation: Scattered black spruce and tamarack; understory consists of bogs

rosemary and cotton grass with some swamp laured ground cover consists

Depth (cm)

0-127

127-162

162-503

503+

(SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 15, T. 154Nk., RIZ8W.)

Described By: R. Fox and R. Zarth on September 22. 19376.

Layer

Fibric

Sapric

Hemic

Sand

Location: 575 feet N and 1425 feet E of the SW corner of Sec. 15, T.154N., R.28W. (NW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 15, T.154N., R.28W.)

Vegetation: Scattered tamarack; lush understory consists mostly of sedges and bog birch with bog rosemary and some leatherleaf and willow; ground cover consists mostly of sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: R. Fox and R. Zarth on September 21, 1976.

Layer	Depth (cm)
Fibric	0- 76
Hemic	76-417
Verv	417+
fine	
sand	

Location: 1830 freet S and 1140 feet W'off the NE corner of Sec. 16, T.154N., R.28W. (SW1/4 of NW#1/4 of SE1/4 of NE1//4, Sec. 16, T.154N., R.28W.)

Vegetation: Lusti winderstory consists mostly of sedges with some tamarack and northern white cedar; ground cover consists of some Sphagnum mosses and pitcher plant:

Microrellef: Not recorded.

Depth To Water Table: At surface.

Described By: R: Zarth and R. Fox on September 27, 1976.

Hemic 0-396 Sandy 396+ loam Location: 630 feet S and 1615 feet W of the NE corner of Sec. 22, T.154N., R.28W. (SE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 22, T.154N., R.28W.)

Location: 2200 feet N and 1050 feet W of the SE corner of Sec. 15, T.154N., R.28W.

Vegetation: Scattered black spruce; lush understory consists mostly of leatherleaf

Depth (cm)

0-254

254-610+

(SW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 15, T.154N., R.28W.)

with Labrador tea and some sedges and cotton grass.

Described By: R. Zarth and R. Fox on September 20, 1976.

Layer

Fibric

Hemic

Bottom

Vegetation: Scattered tamarack; lush understory consists mostly of sedges and bog birch with some pitcher plant.

Microrelief: 60 cm.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox and R. Zarth on September 27, 1976.

ayer	Depth (cm
Hemic	0-488
Sand	488+

Location: 1310 feet S and 1475 feet E of the NW corner of Sec. 22, T.154N., R.28W. (SW1/4 of SW1/4 of NE1/4 of NW1/4, Sec. 22, T.154N., R.28W.)

Vegetation: Scattered tamarack; lush understory consists mostly of bog rosemary and bog birch with sedges and some swamp laurer; ground wave consists of some Sphagnum mosses.

Depth (cm)

0-457 457+

Microrelief: Not recorded.

Depth To Water Table: At surface.

mostly of Sphagnum mosses...

Depth To Water Table: Not recorded.

Microrellef: Not recorded.

Described By: R. Fox and R. Zarth on September 21, 1976.

Laver

Hemic

Medium sand Location: 2500 feet S and 1500 feet E of the NW corner of Sec. 22, T.154N., R.28W. (SW1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 22, T.154N., R.28W.)

Vegetation: Tamarack crown cover of about 35 percent; understory consists of bog birch and bog rosemary with some willow; ground cover consists mostly of Sphagnum mosses with some pitcher plant.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: R. Fox and R. Zarth on September 27, 1976.

Layer	Depth (cm)
Hemic	0-320
Sandy	320+
loam	

Location: 1570 feet N and 1520 feet E of the SW corner of Sec. 22, T.154N., R.28W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 22, T.154N., R.28W.)

Vegetation: Tamarack crown cover of about 35 percent; understory consists of bog birch and bog rosemary with some willow; ground cover consists mostly of Sphagnum mosses with some pitcher plant.

Microrellef: 60 cm.

Depth To Water Table: At surface.

Described By: R. Zarth and R. Fox on September 21, 1976.

Layer	Depth (cm)
Hemic Sandy Ioam	0-178 178+

Location: 825 feet S and 725 feet E of the NW corner of Sec. 23, T.154N., R28W. (NW1/4 of SE1/4 of NW1/4 of NW1/4, Sec. 23, T.154N., R.28W.)

Vegetation: Scattered black spruce; lush understory consists mostly of bog rosemary with sedges and some bog birch; ground cover consists mostly of Sphagnum mosses with some pitcher plant.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: R. Fox and R. Zarth on September 22, 1976.

Layer Depth (cm)

Fibric 0- 38
Hemic 38-472
Pebbly 472+

Location: 1725 feet S and 1275 feet E of the NW corner of Sec. 23, T.154N., R.28W. (SE1/4 of NE1/4 of SW1/4 of NW1/4, Sec. 23, T.154N., R.28W.)

Vegetation: Tamarack crown cover of about 35 percent; lush understory consists mostly of bog rosemary with cattails, bog birch, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrelief: 60 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 23, 1976.

Layer	Depth (cm)
Hemic	0-381
Sand	381+

Location: 2350 feet S and 1150 feet E of the NW corner of Sec. 23, T.154N., R.28W. (SE1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 23, T.154N., R.28W.)

Vegetation: Scattered tamarack; lush understory consists mostly of grasses and leatherleaf with some bog rosemary, Labrador tea, and cattail; ground cover consists mostly of Sphagnum mosses.

Microrelief: 50 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 23, 1976.

Layer	Depth (cm)
Hemic	0-178
Mineral	178+
coil	

Location: 2200 feet N and 1150 feet E of the SW corner of Sec. 23, T.154N., R.28W. (SE1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 23, T.154N., R.28W.)

Vegetation: Scattered tamarack; lush understory consists mostly of grasses with Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrelle: 60 cm

Depth To Water Table: At surface.

Described By: M. Domeier on September 23, 1976.

Layer	Depth (cm)
Hemic	0-50
Sapric	50-56
Coarse	56+
sand	

Location: 2500 feet S and 1750 feet W of the NE corner of Sec. 2, T.155N., R.28W. (SW1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 2, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 65 percent; lush understory consists mostly of leatherleaf with some cotton grass, sedges, and Labrador tea: ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: R. Fox on August 13, 1976.

n)

Location: 1680 feet S and 1150 feet W of the NE corner of Sec. 2, T.155N., R.28W. (NW1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 2, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of cotton grass and sedges with some leatherleaf, swamp laurel, and Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox on August 13, 1976.

Layer	Depth (cr
Hemic	0-246
Sand	246+

Location: 1300 feet N and 2430 feet E of the SW corner of Sec. 2, T.155N., R 28W (NE1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 2, T.155N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of cotton grass with leatherleaf and some swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses and false Solomon's seal.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: R. Fox on August 13, 1976.

Layer	Depth (cm)
Hemic	0-206
Sand	206+

Location: 475 feet N and 1800 feet E of the SW corner of Sec. 2, T.155N., R.28W. (NE1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 2, T.155N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of leatherleaf, sedges, and cotton grass with some bog rosemary and bog birch; ground cover consists mostly of Sphagnum mosses with cranberry, false Solomon's seal and some other mosses.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: R. Fox on August 13, 1976.

Layer	Depth (cm)
Hemic	0-132
Medium	132+
to	
coarse	
sand	

Location: 2125 feet N and 2325 feet W of the SE corner of Sec. 2, T.155N., R.28W. (SE1/4 of NW1/4 of NW1/4 of SE1/4, Sec. 2, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of leatherleaf with some Labrador tea, bog rosemary, swamp laurel, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrelief: 60 cm.

Depth To Water Table: At surface.

Described By: R. Fox on August 13, 1976.

Layer	Depth (cm
Hemic	0-267
Medium	267+
sand	

Location: 60 feet S and 50 feet W of the NE corner of Sec. 2, T.155N., R.28W. (Gov. Lot 1, Sec. 2, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 65 percent with scattered tamarack; understory consists of sedges, Labrador tea, and leatherleaf; ground cover consists of some Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on July 24, 1976.

Depth (cm) Layer 0-152 Hemic Sand 152+

Location: 825 feet S and 540 feet W of the NE corner of Sec. 2, T.155N., R.28W (Gov. Lot 1, Sec. 2, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent; lush understory consists mostly of sedges and cotton grass with some Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry, other mosses, and false Solomon's seal.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: R. Fox on August 13, 1976.

Layer	Depth (cm)
Hemic	0-122
Sand	122+

Location: 2075 feet S and 1200 feet F of the NW corner of Sec. 6, T.155N., R.28W. (NW1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 6, T.155N., R.28W.)

Vegetation: Scattered black spruce; understory consists of Labrador tea and leatherleaf; ground cover consists of Sphagnum mosses and false Solomon's

Microrelief: Not recorded.

Depth To Water Table: 15 cm.

Described By: R. Fox on August 5, 1976.

Layer	Depth (cm)
Hemic	0-91
Sandy	91+
loam	

Location: 2110 feet S and 2260 feet E of the NW corner of Sec. 6, T.155N., B.28W (NE1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 6, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 80 percent; lush understory consists of Labrador tea, leatherleaf, and grasses with some Indian pipe; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: 15 cm.

Described By: R. Fox on August 5, 1976.

Layer	Depth (cm)
Hemic Sapric Sand with calcareous pebbles	0-206 206-229 229+

Location: 2050 feet N and 2300 feet E of the SW corner of Sec. 6, T.155N., R.28W. (SE1/4 of NE1/4 of NE1/4 of SW1/4, Sec. 6, T.155N., R.28W.)

Vegetation: Scattered black spruce and tamarack; understory consists of Labrador tea, swamp laurel, leatherleaf, grasses, and pitcher plant; ground cover consists of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: 5 cm.

Described By: R. Fox on August 5, 1976.

Layer	Depth (cm)
Hemic	0-175
Loamy	175+
sand	

Location: 40 feet N and 2290 feet E of the SW corner of Sec. 6, T.155N., R.28W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 6, T.155N., R.28W.)

Vegetation: Scattered black spruce; understory consists of Labrador tea, leatherleaf, and grasses; ground cover consists of Sphagnum mosses, cranberry, and feather mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on August 5, 1976.

Layer	Depth (cm)
Fibric	0- 18
Hemic	18-120
Loamy	120+
sand	

Location: 110 feet S and 2600 feet W of the NE corner of Sec.6, T.155N., R.28W. (Gov. Lot 2, Sec. 6, T.155N., R.28W.)

Vegetation: Scattered black spruce and tamarack; understory consists of bog birch, Labrador tea, leatherleaf, and sedges; ground cover consists of Sphagnum mosses.

Depth (cm)

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Zarth, R. Fox, and M. Domeier on August 5, 1976.

Layer

Location: 1300 feet S and 2325 feet E of the NW corner of Sec. 6, T.155N., R.28W

Vegetation: Black spruce crown cover of about 95 percent; lush understory consists mostly of Labrador tea with leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: 15 cm.

(Gov. Lot 3, Sec. 6, T.155N., R.28W.)

Described By: R. Zarth, R. Fox, T. Malterer, and M. Domeier on August 5, 1976.

Layer	Depth (cm)
Fibric	0- 23
Hemic	23-200
Sapric	200-213
Loamy	213+
sand	

Location: 2075 feet S and 625 feet E of the NW corner of Sec. 6, T.155N., R.28W. (Gov. Lot 5, Sec. 6, T.155N., R.28W.)

Vegetation: Tamarack crown cover of about 60 percent; understory consists of bog birch with some Labrador tea, sedges, and cotton grass; ground cover consists of some ferns.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on August 5, 1976.

Layer	Depth (cm)
Hemic	0-101
Loamy	101+
sand	

Hemic 0-178 Sapric 178-183 Loamy 183+ sand with calcareous pebbles

Location: 2150 feet S and 2130 feet E of the NW corner of Sec. 7, T.155N., R.28W. (NW1/4 of SE1/4 of SE1/4 of NW1/4, Sec. 7, T.155N., R.28W.)

Vegetation: Scattered black spruce; understory consists of Labrador tea and leatherleaf; ground cover consists of Sphagnum mosses and cranberry.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on August 5, 1976.

Layer Depth (cm)

Hemic 0-198
Clay 198+
with sand varves

Location: 2375 feet S and 275 feet W of the NE corner of Sec. 10, T.155N., R.28W. (SE1/4 of SE1/4 of SE1/4 of NE1/4, Sec. 10, T.155N., R.28W.)

Vegetation: Tamarack crown cover of about 35 percent; understory consists of bog birch and Labrador tea; ground cover consists of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: R. Fox on August 5, 1976.

Layer	Depth (cm)
Hemic	0-182
Sapric	182-193
Sandy	193+
clay	
loam	

Location: 320 feet S and 1200 feet E of the NW corner of Sec. 11, T.155N., R.28W. (NE1/4 of NB1/4 of NW1/4, Sec. 11, T.155N., R.28W.)

Vegetation: Lush understory consists mostly of leatherleaf, sedges, and cotton grass with bog rosemary and some bog birch; ground cover consists mostly of Sphagnum mosses.

Microrelief: 60 cm.

Depth To Water Table: At surface.

Described By: R. Fox on August 5, 1976.

Layer	Depth (cm)
Hemic	0-170
Sandy	170+
clay	
loam	

Location: 1220 feet S and 625 feet E of the NW corner of Sec. 11, T.155N., R.28W. (SE1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 11, T.155N., R.28W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Layer	Depth (cm
Hemic	0-162
Sapric	162-173
Mineral	173+

soil

Location: 300 feet N and 1600 feet E of the SW corner of Sec. 26, T.155N., R.28W. (SW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 26, T.155N., R.28W.)

Vegetation: Consists of tamarack, speckled alder, and grasses.

Microrelief: 8 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer	Depth (cm)
Hemic	0-61
Sand	61+

Location: 1040 feet S and 900 feet W of the NE corner of Sec. 34, T.155N., R.28W. (SE1/4 of SW1/4 of NE1/4 of NE1/4, Sec. 34, T.155N., R.28W.)

Vegetation: Black spruce and paper birch crown cover of about 35 percent each; lush understory consists mostly of speckled alder with sedges; ground cover consists mostly of Sphagnum mosses.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer	Depth (cm)
Sapric Medium sand	0-74 74+

Location: 1090 feet S and 2390 feet E of the NW corner of Sec. 34, T.155N., R.28W. (SE1/4 of SE1/4 of NE1/4 of NW1/4, Sec. 34, T.155N., R.28W.)

Vegetation: Northern white cedar crown cover of about 60 percent with scattered paper birch, balsam fir, and black spruce; understory consists of speckled alder, sedges, and grasses.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer	Depth (cm)	
Hemic	0-41	
Mineral	41+	
soil		

Location: 2590 feet S and 100 feet E of the NW corner of Sec. 34, T.155N., R.28W. (SW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 34, T.155N., R.28W.)

Vegetation: Scattered black spruce and tamarack; understory consists of Labrador tea with some speckled alder; ground cover consists mostly of Sphagnum mosses

Microrelief: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer	Depth (cm)
Sapric	0-112
Fine	112+
sand	

Location: 1775 feet N and 1975 feet E of the SW corner of Sec. 34, T.155N., R.28W. (NW1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 34, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent; understory consists of speckled alder; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer	Depth (cm)
Sapric Sandy	0-79 79+
loam	

Location: 475 feet S and 1430 feet W of the NE corner of Sec. 35, T.155N., R.28W. (SE1/4 of NE1/4 of NW1/4 of NE1/4, Sec. 35, T.155N., R.28W.)

Vegetation: Black spruce and tamarack crown cover of about 35 percent each with scattered balsam fir and paper birch; understory consists of Labrador tea with speckled alder; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer Depth (cm)
Sapric 0-30
Clay 30+
loam

Location: 1100 feet N and 575 feet E of the SW corner of Sec. 35, T.155N., R.28W. (NE1/4 of NE1/4 of SW1/4 of SW1/4, Sec. 35, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 85 percent; understory consists of grasses and Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer Depth (cm)

Hemic 0-89
Sandy 89+
loam
with calcareous
pebbles

Location: 1100 feet N and 1740 feet E of the SW corner of Sec. 35, T.155N., R.28W. (NE1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 35, T.155N., R.28W.)

Location: 1200 feet S and 760 feet E of the NW corner of Sec. 35. T.155N., R.28W.

Vegetation: Scattered black spruce and paper birch; understory consists mostly of

speckled alder with sedges; ground cover consists of some Sphagnum

Depth (cm)

0-46

46+

(SW1/4 of SE1/4 of NW1/4 of NW1/4, Sec. 35, T.155N., R.28W.)

Vegetation: Not recorded.

Microrellef: 30 cm.

mosses.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer

Clay

loam

Depth To Water Table: Not recorded.

Described By: R. Zarth on September 16, 1976.

Depth (cm)
0- 51
51-122
122+

Location: 2200 feet N and 630 feet E of the SW corner of Sec. 36, T.155N., R.28W. (SW1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 36, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent; understory consists of leatherleaf with some Labrador tea, bog rosemary, and swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: R. Fox on September 16, 1976.

Layer Depth (cm)
Fibric 0- 36
Hemic 36-114
Sapric 114-124
Sandy 124+
clay
loam

Location: 710 feet N and 460 feet W of the SE corner of Sec. 36, T.155N., R.28W. (SW1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 36, T.155N., R.28W.)

Vegetation: Not recorded.

Microrellef: 15 cm.

Depth To Water Table: 3 cm.

Described By: R. Zarth on September 9, 1976.

Layer Depth (cm)
Fibric 0- 30
Hemic 30-208
Loam 208+

Location: 1900 feet N and 500 feet E of the SW corner of Sec. 35, T.155N., R.28W. (NE1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 35, T.155N., R.28W.)

Vegetation: Cut over area consists mostly of grasses with speckled alder, leatherleaf, and sedges.

Microrellef: Negligible.

Depth To Water Table: Not recorded.

Described By: M. Domeier on November 4, 1976.

Layer	Depth (cm)
Hemic	0- 75
Sapric	75-114
Sandy	114+
loam	

Location: 1200 feet S and 450 feet W of the NE corner of Sec. 36, T.155N., R.28W. (SW1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 36, T.155N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent; lush understory consists mostly of balsam fir and leatherleaf; ground cover consists of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on September 9, 1976.

Layer	Depth (cm)
Fibric	0- 13
Hemic	13-168
Coarse	168+
sand	

Location: 1140 feet S and 310 feet W of the NE corner of Sec. 3, T.156N., R.28W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 3, T.156N., R.28W.)

Vegetation: Tamarack crown cover of about 50 percent with a black spruce crown cover of about 30 percent; lush understory consists of Labrador tea with some bog birch and ferns; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, false Solomon's seal, and pitcher plant.

Microrellef: 20 cm.

Depth To Water Table: 15 cm.

Described By: B. Wakanabo on August 4, 1977.

Layer	Depth (cm)
Hemic	0-215
Sapric	215-228
Sand	228+

Location: 1310 feet N and 300 feet W of the SE corner of Sec. 3. T.156N., R.28W. (NE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 3. T.156N., R.28W.)

Vegetation: Tamarack crown cover of about 75 percent with a black spruce crown cover of about 20 percent; lush understory consists of Labrador tea with some grasses and bog birch; ground cover consists mostly of Sphagnum mosses with cranberry.

Microrellef: 30 cm.

Depth To Water Table: 20 cm.

Described By: B. Wakanabo on August 4, 1977.

Layer Depth (cm)

Hemic 0-282
Sapric 282-310
Sandy 310+
clay

Location: 1345 feet S and 280 feet W of the NE corner of Sec. 10, T.156N., R.28W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 10, T.156N., R.28W.)

Vegetation: Northern white cedar crown cover of about 60 percent with a black spruce crown cover of about 30 percent with scattered tamarack; lush understory consists mostly of grasses with some dogwood and ferns; ground cover consists of cranberry with some Sphagnum and other mosses.

Microrelief: 40 cm.

Depth To Water Table: 15 cm.

Described By: B. Wakanabo on August 4, 1977.

Layer	Depth (cm)
Hemic Clay Ioam	0-58 58+

Location: 75 feet N and 1200 feet E of the SW corner of Sec. 11, T.156N., R.28W. (SE1/4 of SE1/4 of SW1/4, Sec. 11, T.156N., R.28W.)

Vegetation: Not recorded.
Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described: In July, 1976.

Layer	Depth (cm)
Hemic Clay	0-152 152+
loam	

Location: 50 feet N and 2200 feet E of the SW corner of Sec. 11, T.156N., R.28W. (SW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 11, T.156N., R.28W.)

Vegetation: Not recorded.
Microrelief: Not recorded.
Depth To Water Table: Not recorded.
Described By: Not recorded.

Layer

Hemic

Sand

Depth (cm) 0-182 182+ Location: 125 feet N and 625 feet W of the SE corner of Sec. 11, T. 156N., R.28W. (SE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 11, T.156N., R.28W.)

Location: 75 feet N and 225 feet E of the SW corner of Sec. 11, T.156N., R.28W.

Depth (cm)

0-121

121+

(SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 11, T.156N., R.28W.)

Vegetation: Cut over area surrounded by about 90 percent crown cover of black spruce with scattered tamarack and northern white cedar; sparse understory consists of grasses, raspberry, and goldenrod.

Microrelief: Not recorded.

Vegetation: Not recorded.

Microrelief: Not recorded.

Described: In July, 1976.

Depth To Water Table: Not recorded.

Layer

Hemic Silt

Depth To Water Table: Not recorded

Described By: D. Olson and D. Haverkost on November 13, 1979.

Layer	Depth (cm	
Hemic	0-122	
Sapric	122-152	
Very	152+	
fine		
sand		

Location: 1375 feet S and 200 feet W of the NE corner of Sec. 13, T.156N., R.28W. (SE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 13, T.156N., R.28W.)

Vegetation: Black spruce crown cover of about 50 percent with a northern white cedar crown cover of about 35 percent; understory consists of grasses with some Labrador tea and leatherleaf.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and R. Graber on November 13, 1979.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-213
Fine	213+
sand	

Location: 1450 feet S and 225 feet W of the NE corner of Sec. 15, T.156N., R.28W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 15, T.156N., R.28W.)

Vegetation: Scattered spruce; lush understory consists mostly of grasses with shrubs, Labrador tea, bog birch, and some bog rosemary and ferns; ground cover consists of Sphagnum mosses with some other mosses.

Microrellef: 20 cm.

Depth To Water Table: 50 cm.

Described By: B. Wakanabo on August 3, 1977.

Layer	Depth (cm)
Hemic	0-136
Loamy	136+
sand	

Location: 1485 feet S and 590 feet W of the NE corner of Sec. 15, T.156N., R.28W. (NW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 15, T.156N., R.28W.)

Vegetation: Tamarack crown cover of about 75 percent with scattered cedar; lush understory consists mostly of cotton grass with bog birch and some Labrador tea and bog rosemary; ground cover consists of Sphagnum mosses with some other mosses, false Solomon's seal and cranberry.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: B. Wakanabo on August 3, 1977.

Layer	Depth (cm)
Hemic	0-190
Sand	190+

Location: 75 feet N and 250 feet W of the SE corner of Sec. 15, T.156N., R.28W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 15, T.156N., R.28W.)

Vegetation: Tamarack crown cover of about 40 percent with a black spruce crown cover of about 20 percent; lush understory consists mostly of sedges with some willow and dogwood.

Microrellef: 15 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 21, 1976.

Layer Depth (cm) 0-28 Hemic 28+ Sandy loam

Location: 1525 feet N and 2600 feet E of the SW corner of Sec. 22, T.156N., R.28W. (SE1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 22, T.156N., R.28W.)

Vegetation: Open bog. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: C. Cameron and M. Domeier on July 19, 1976.

Layer	Depth (cm)
Peat	0-30
Sand	30+

Location: 1550 feet N and 225 feet W of the SE corner of Sec. 22, T.156N., R.28W. (SE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 22, T.156N., R.28W.)

Vegetation: Black spruce crown cover of about 50 percent; understory consists of Labrador tea; ground cover consists of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: C. Cameron and M. Domeier on July 19, 1976.

Layer	Depth (cm)
Peat	0-15
Sand	15+

Location: 125 feet S and 50 feet E of the NW corner of Sec. 26, T.156N., R.28W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 26, T.156N., R.28W.)

Depth To Water Table: Not recorded.

Described By: T. Malteror, C. Cameron, and R. Zarth in July, 1976.

Vegetation: Not recorded. Microrellef: Not recorded.

	•
Layer	Depth (cm)
Hemic	0-76
Sand	76+

Location: 500 feet S and 50 feet E of the NW corner of Sec. 26, T.156N., R.28W. (SW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf and Labrador tea.

Microrelief: 45 cm.

Depth To Water Table: Not recorded.

Described: In July, 1976.

Layer	Depth (cm)
Fibric	0- 51
Hemic	51-182
Sapric	182-198
Medium	198+
sand	

Location: 1000 feet S and 50 feet E of the NW corner of Sec. 26, T.156N., R.28W. (SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Not recorded.

Microrellef: 35 cm.

Depth To Water Table: Not recorded.

Described: In July, 1976.

Layer	•	Depth (cm)
Fibric		0- 40
Hemic		40-213
Sapric		213-228
Medium		228+
sand		

Location: 1500 feet S and 50 feet E of the NW corner of Sec. 26, T.156N., R.28W. (NW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Scattered black spruce; understory consists of grasses, leatherleaf, and Labrador tea; ground cover consists of Sphagnum mosses.

Microrellef: 35 cm.

Depth To Water Table: At surface.

Described: In July, 1976.

Laver Depth (cm) Fibric 0- 43 43-274 Hemic 274+ Silt loam

Location: 2000 feet S and 50 feet E of the NW corner of Sec. 26, T.156N., R.28W. (NW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Black spruce crown cover of about 50 percent with scattered tamarack.

Microrellef: 90 cm.

Depth To Water Table: At surface.

Described: In July, 1976.

Layer	Depth (cm)	
Fibric	0- 43	
Hemic	43-304	
Sand	304+	

Location: 2500 feet S and 50 feet E of the NW corner of Sec. 26, T.156N., R.28W. (SW1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described: in July, 1976.

Layer	Depth (cm)
Fibric	0-122
Hemic	122-304
Sapric	304-320
Medium	320+
sand	

Location: 2275 feet N and 2385 feet E of the SW corner of Sec. 26, T.156N., R.28W. (SE1/4 of NE1/4 of NE1/4 of SW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Layer	Depth (cm
Fibric	0- 35
Peat	35-426
Mineral	426+
soil	

Location: 2225 feet N and 50 feet E of the SW corner of Sec. 26, T.156N., R.28W. (SW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Not recorded.
Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described: In July, 1976.

Layer	Depth (cm)
Fibric	0- 56
Hemic	56-304
Sapric	304-335
Sand	335+

Location: 1725 feet N and 50 feet E of the SW corner of Sec. 26, T.156N., R.28W. (NW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Consists mostly of black spruce and Labrador tea with some leatherleaf.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described: In July, 1976.

Layer	Depth (cm)
Fibric	0- 56
-lemic	56-274
Sapric	274~305
Sand	305+

Location: 1265 feet N and 50 feet E of the SW corner of Sec. 26, T.156N., R.28W. (NW1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Not recorded.
Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described: In July, 1976.

Layer	Depth (cm)
Fibric	0-168
Hemic	168-365
Sapric	365-396
Sand	396+

Location: 290 feet N and 1040 feet E of the SW corner of Sec. 26, T.156N., R.28W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 26, T.156N., R.28W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Layer	Depth (cm
Fibric	0-254
Peat	254-548
Mineral	548+
soil	

Location: 2330 feet N and 2300 feet E of the SW corner of Sec. 27, T.156N., R.28W. (NE1/4 of NE1/4 of NE1/4 of SW1/4, Sec. 27, T.156N., R.28W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Depth (cm)
0- 40
40-426
426+

Location: 875 feet N and 1025 feet W of the SE corner of Sec. 27, T.156N., R.28W. (SW1/4 of NW1/4 of SE1/4 of SE1/4, Sec. 27, T.156N., R.28W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Layer	Depth (cm)
Fibric	0-182
Peat	182-457
Mineral soil	457+

Location: 400 feet N and 2500 feet E of the SW corner of Sec. 31, T.156N., R.28W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 31, T.156N., R.28W.)

Vegetation: Not recorded.
Microrelief: Not recorded.
Depth To Water Table: Not recorded.
Described By: Not recorded.

Layer	Depth (cm)
Peat	0-107
Mineral	107+

Location: 200 feet N and 2500 feet E of the SW corner of Sec. 31, T.156N., R.28W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 31, T.156N., R.28W.)

Vegetation: Not recorded.
Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Layer	Depth (cm
Peat	0-152
Mineral	152+
soil	

Location: 1350 feet S and 50 feet W of the NE corner of Sec. 34, T.156N., R.28W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 34, T.156N., R.28W.)

Vegetation: Not recorded. Microrelief: Not recorded.

Depth To Water Table: Not recorded. Described By: Not recorded.

> Layer Depth (cm) Fibric 0-203 Pest 203-457 Mineral 457+ soil

Location: 2200 feet S and 650 feet W of the NE corner of Sec. 34, T.156N., R.28W. (NE1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 34, T.156N., R.28W.)

Vegetation: Not recorded. Microrellef: Not recorded. Depth To Water Table: Not recorded. Described By: Not recorded.

> Layer Depth (cm) Fibric 0-116 Peat 116-365 Mineral 365+ soil

Location: 650 feet S and 2200 feet E of the NW corner of Sec. 34, T.156N., R.28W. (SW1/4 of NE1/4 of NE1/4 of NW1/4, Sec. 34, T.156N., R.28W.)

Vegetation: Not recorded. Microrelief: Not recorded. Depth To Water Table: Not recorded. Described By: Not recorded.

> Layer Depth (cm) Fibric 0- 60 Peat 60-426 Mineral 426+ soil

Location: 2425 feet N and 1075 feet W of the SE corner of Sec. 34, T.156N., R.28W. (NW1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 34, T.156N., R.28W.) Vegetation: Not recorded.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Laye? Depth (cm) Fibric 0- 30 Peat 30-304 Mineral 304+ soit

Location: 1790 feet N and 1475 feet W of the SE corner of Sec. 34, T.156N., R.28W. (NE1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 34, T.156N., R.28W.)

Vegetation: Not recorded. Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Layer	Depth (cm)
Peat Mineral	0-61 61+
soil	- · ·

Location: 500 feet S and 500 feet E of the NW corner of Sec. 35, T.156N., R.28W. (SE1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 35, T.156N., R.28W.)

Vegetation: Not recorded. Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: Not recorded.

Layer	Depth (cm)
Fibric	0-243
Peat	243-518
Mineral	518+
soil	

Location: 1800 feet S and 775 feet W of the NE corner of Sec. 5, T.157N., R.28W. (SE1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 5, T.157N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges and bog birch with some Labrador tea, leatherleaf, horsetail, and grasses.

Microrellef: Not recorded.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on November 13, 1979.

Layer Depth (cm) Hemic 0-106 Mineral 106+ Soil

Location: 2600 feet N and 660 feet E of the SW corner of Sec. 14, T.157N., R.28W. (NE1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 14, T.157N., R.28W.) Vegetation: Scattered northern white cedar, paper birch, and black spruce; lush

understory consists mostly of speckled alder and grasses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 12, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-65 65+

Location: 1575 feet N and 900 feet E of the SW corner of Sec. 15, T.157N., R.28W. (SW1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 15, T.157N., R.28W.)

Vegetation: Northern white cedar crown cover of about 30 percent with scattered black spruce; lush understory consists mostly of grasses with alder and horsetail.

Microrelief: Not recorded.

Depth To Water Table: 10 cm. above surface.

Described By: D. Mellem and D. Haverkost on September 12, 1979.

Depth (cm)
0-145 145+

Location: 950 feet N and 2390 feet E of the SW corner of Sec. 17, T.157N., R.28W. (SE1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 17, T.157N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of bog birch and leatherleaf with some Labrador tea; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and R. Graber on November 13, 1979.

Layer	Depth (cm)
Hemic	0- 75
Sapric	75-122
Clay	122+
loam	

Location: 75 feet N and 25 feet E of the SW corner of Sec. 26, T.157N., R.28W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 26, T.157N., R.28W.)

Vegetation: Consists mostly of northern white cedar.

Microrelief: 5 cm.

Depth To Water Table: Not recorded.

Described By: M. Domeier on October 21, 1976.

Layer	Depth (cm)
Sapric Clay Ioam	0-20 20-35
Sand	35+

Location: 1730 feet S and 1330 feet W of the NE corner of Sec. 30, T.157N., R.28W. (SW1/4 of NW1/4 of SE1/4 of NE1/4, Sec. 30, T.157N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of leatherleaf with bog birch, sedges, and some swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on November 13, 1979.

Layer	Depth (cm
Fibric	0- 61
Hemic	61-152
Mineral	152+
soil	

Location: 300 feet S and 325 feet E of the NW corner of Sec. 30, T.157N., R.28W. (Gov. Lot 1, Sec. 30, T.157N., R.28W.)

Vegetation: Understory consists mostly of sedges with cotton grass, bog birch and some speckled alder and cattails.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on August 2, 1979.

Layer	Depth (cm)
Hemic	0-91
Medium	91+
eand	

Location: 100 feet N and 525 feet E of the SW corner of Sec. 31, T.157N., R.28W. (Gov. Lot 4. Sec. 31, T.157N., R.28W.)

Vegetation: Scattered tamarack; lush understory consists mostly of sedges with some bog rosemary, speckled alder, bog birch, and horsetall; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, cranberry, and pitcher plant,

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on August 1, 1979.

Layer	Depth (cm
Hemic	0- 71
Sapric	71-173
Silty	173+
clay	
loam	

Location: 225 feet N and 1525 feet E of the SW corner of Sec. 6, T.158N., R.28W. (SW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 6, T.158N., R.28W.)

Vegetation: Lush understory consists mostly of sedges with some black spruce, northern white cedar, bog birch, willow, bog rosemary, Labrador tea, leatherleaf, and cattail; ground cover consists of some Sphagnum mosses. Microrelief: 10 cm.

Depth To Water Table: 25 cm.

Described By: D. Olson and N. Aaseng on October 29, 1979.

Layer	Depth (cn
Hemic Silty clay	0-167 167+

Location: 100 feet N and 1865 feet W of the SE corner of Sec. 7, T.158N., R.28W. (SW1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 7, T.158N., R.28W.)

Vegetation: Understory consists mostly of sedges with some bog birch, willow, grasses, and leatherleaf; ground cover consists of some Sphagnum and other mosses.

Microrellef: 10 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on October 29, 1979.

Layer	Depth (cm)
Hemic	0-127
Fine sand	127+

Location: 100 feet S and 150 feet W of the NE corner of Sec. 8, T.158N., R.28W. (NE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 8, T.158N., R.28W.)

Vegetation: Black spruce crown cover of about 65 percent with scattered tamarack; sparse understory consists of sedges, bog rosemary, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and Indian pipe.

Microrelief: 10 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on October 29, 1979.

Layer	Depth (cm)
Hemic	0-233
Clay	233+

Location: 160 feet N and 160 feet W of the SE corner of Sec. 8, T.158N., R.28W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 8, T.158N., R.28W.)

Vegetation: Scattered black spruce and northern white cedar; lush understory consists mostly of grasses and speckled alder.

Microrelief: Negligible.

Depth To Water Table: 15 cm.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Layer	Depth (cm)
Hemic Fine	0-20 20+
sand	207

Location: 1675 feet S and 490 feet W of the NE corner of Sec. 9, T.158N., R.28W. (SW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 9, T.158N., R.28W.)

Vegetation: Black spruce crown cover of about 65 percent; understory consists of Labrador tea with some sedges and leatherleaf; ground cover consists mostly of mosses including Sphagnum mosses, false Solomon's seal and some cranberry and lichens

Microrellef: 20 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Haverkost on November 2, 1979.

Layer	Depth (cn
Hemic Silty clay	0-167 167+

m)

Location: 1975 feet S and 525 feet W of the NE corner of Sec. 13, T.158N., R.28W. (SW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 13, T.158N., R.28W.)

Vegetation: Scattered black spruce and tamarack: lush understory consists mostly of sedges with leatherleaf and some swamp laurel and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry and pitcher plant.

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Haverkost on November 2, 1979.

Layer	Depth (cm)
Hemic	0-180
Sapric	180-225
Hemic	225-330
Sapric	330-390
Fine	390+
sand	
and	
clay	

Location: 475 feet S and 950 feet E of the NW corner of Sec. 20, T.158N., R.28W. (SW1/4 of NE1/4 of NW1/4 of NW1/4, Sec. 20, T.158N., R.28W.)

Vegetation: Scattered black spruce, northern white cedar, and tamarack; sparse understory consists of some sedges, Labrador tea, and dogwood; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not visible.

Described By: D. Olson and D. Mellem on September 19, 1979.

Laver Depth (cm) Sapric 0-76 Sandy 76+ loam

Location: 2350 feet S and 2430 feet W of the NE corner of Sec. 11, T.158N., R.28W. (SW1/4 of SW1/4 of SW1/4 of NE1/4, Sec. 11, T.158N., R.28W.)

Vegetation: Black spruce crown cover of about 85 percent; understory consists of leatherleaf with some swamp laurel and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 2, 1976.

Layer	Depth (cm)
Fibric	0- 46
Hemic	46-125
Sapric	125-150
Hemic	150-323
Clay	323+
loam	

Location: 800 feet N and 1365 feet E of the SW corner of Sec. 14, T.158N., R.28W. (SW1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 14, T.158N., R.28W.)

Vegetation: Black spruce crown cover of about 50 percent; lush understory consists mostly of Labrador tea and grasses; ground cover consists mostly of Sphagnum mosses and cranberry with some false Solomon's seal.

Microrelief: 25 cm.

Depth To Water Table: At surface.

Described By: M. Domeier on September 23, 1976.

Layer	Depth (cm)
Fibric Hemic	0-290 290-549
Mineral soil	549+

Location: 2240 feet S and 900 feet E of the NW corner of Sec. 20. T.158N., R.28W. (NW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 20, T.158N., R.28W.)

Vegetation: Black spruce crown cover of about 50 percent with a tamarack crown cover of about 40 percent; sparse understory consists of some Labrador tea and leatherleaf; ground cover consists of Sphagnum and other mosses.

Microrelief: Negligible.

Depth To Water Table: Below 25 cm.

Described By: D. Olson and D. Mellem on September 19, 1979.

Layer	Depth (cm)
Hemic	0-25
Fine	25+
eand	

Location: 75 feet S and 1625 feet E of the NW corner of Sec. 11, T.158N., R.28W. (NW1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 11, T.158N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; sparse understory consists of some grasses and cattails; ground cover consists mostly of mosses.

Microrellef: Not recorded

Depth To Water Table: At surface.

Described By: D. Olson and R. Graber on November 1, 1979.

Layer	Depth (cm
Hemic	0-244
Silt	244+
and	
very	
fine	
sand	

Location: 645 feet N and 1925 feet E of the SW corner of Sec. 17, T.158N., R.28W. (NW1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 17, T.158N., R.28W.)

Vegetation: Scattered black spruce and tamarack: lush understory consists mostly of grasses; ground cover consists of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not visible.

Described By: D. Oison and D. Mellem on September 19, 1979.

Layer	Depth (cm
Hemic Sandy clay	0-110 110+

Location: 1765 feet N and 885 feet E of the SW corner of Sec. 20, T.158N., R.28W (NW1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 20, T.158N., R.28W.)

Vegetation: Black spruce crown cover of about 65 percent with scattered tamarack; sparse understory consists of some Labrador tea, bog birch, and leatherleaf; ground cover consists mostly of Sphagnum mosses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Olson and D. Mellem on September 19, 1979.

Layer	Depth (cm)
Hemic	0-105
Sapric	105-151
Clay with	151+
calcareous pebbles	

Location: 1235 feet N and 150 feet E of the SW corner of Sec. 25, T.158N., R.28W. (NW1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 25, T.158N., R.28W.)

Vegetation: Scattered black spruce, tamarack, and northern white cedar; lush understory consists mostly of grasses with some willow.

Microrelief: 15 cm.

Depth To Water Table: Below 90 cm.

Described By: D. Olson and D. Mellem on September 11, 1979.

Layer	Depth (cm)
Fibric Fine sand and clay	0-90 90+

Location: 2250 Yeet S and 2525 feet W of the NE corner of Sec. 32, T.158N., R.28W. (NW1/4 of SW1/4 of SW1/4 of NE1/4, Sec. 32, T.158N., R.28W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some swamp laurel and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on November 13, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-281
Sandy	281+
clav	

Location: 1125 feet N and 800 feet W of the SE corner of Sec. 32, T.158N., R.28W. (NE1/4 of NW1/4 of SE1/4 of SE1/4, Sec. 32, T.158N., R.28W.)

Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf with some swamp laurel; ground cover consists mostly of Sphagnum mosses.

Microrellef: 45 cm.

Depth To Water Table: Not recorded.

Described By: D. Mellem and R. Graber on November 13, 1979.

Layer	Depth (cm)
Hemic Sandy	0-122 122+
olaria y	1221

Location: 1650 feet N and 1900 feet W of the SE corner of Sec. 34, T.158N., R.28W. (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 34, T.158N., R.28W.)

Vegetation: Black spruce crown cover of about 75 percent; understory consists of Labrador tea with some sedges, bog rosemary, and leatherleaf; ground cover consists mostly of mosses including Sphagnum mosses with some lingen berry, cranberry, and blueberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Haverkost on November 2, 1979.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-120
Sapric	120-180
Silty	180+
clay	
with	
pebbles	

Location: 75 feet S and 1700 feet E of the NW corner of Sec. 2, T.159N., R.28W. (Gov. Lot 3, Sec. 2, T.159N., R.28W.)

Vegetation: Black spruce crown cover of about 75 percent with an aspen crown cover of about 20 percent; understory consists of some Labrador tea, grasses, and leatherleaf.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem on November 13, 1979.

Layer	Depth (cm)
Hemic Very	0-183 183+
fine	
sand	

Location: 900 feet N and 1450 feet W of the SE corner of Sec. 5, T.159N., R.28W. (SE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 5, T.159N., R.28W.)

Vegetation: Black spruce crown cover of about 60 percent; sparse understory consists of some sedges, balsam fir, and Labrador tea; ground cover consismostly of Sphagnum mosses with some other mosses and bunchberry.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on October 30, 1979.

_ayer	Depth (cm
lemic	0-213
Silty	213+
alay	
oam	

Location: 575 feet S and 2125 feet W of the NE corner of Sec. 6, T.159N., R.28W. (Gov. Lot 2, Sec. 6, T.159N., R.28W.)

Vegetation: Black spruce crown cover of about 35 percent with scattered tamarack; sparse understory consists of some Labrador tea, sedges, and alder; ground cover consists mostly of mosses including Sphagnum mosses with some lingen berry, goldthread, bunchberry, snowberry, and false Solomon's seal.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Fibric Hemic Silty clay	0- 91 91-152 152+

Location: 700 feet S and 100 feet E of the NW corner of Sec. 7, T.159N., R.28W. (Gov. Lot 1, Sec. 7, T.159N., R.28W.)

Vegetation: Understory consists mostly of grasses with scattered poplar.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Hemic	0-180
Silty	180+
clay	

Location: 525 feet N and 1655 feet E of the SW corner of Sec. 9, T.159N., R.28W. (NW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 9, T.159N., R.28W.)

Vegetation: Scattered aspen, balsam fir, tamarack, and black spruce; lush understory consists mostly of grasses with alder and some Labrador tea; ground cover consists of some Sphagnum and other mosses, bunchberry, and raspberry.

Microrelief: Not recorded.

Depth To Water Table: Below 45 cm.

Described By: D. Mellem and N. Aaseng on November 1, 1979.

.ayer	Depth (cm)
Sapric	0-45
Coarse	45+
andy	
lay	

Location: 975 feet N and 150 feet W of the SE corner of Sec. 3, T.152N., R.29W. (NE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 3, T.152N., R.29W.)

Vegetation: Scattered black spruce, tamarack, and northern white cedar; sparse understory consists of some Labrador tea, dogwood, willow, alder, and bog birch: ground cover consists mostly of Sphagnum mosses with some other mosses and pitcher plant.

Microrellef: 20 cm.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 30, 1979.

Laver Depth (cm) 0-272 Hemic Fine 272+ and medium sand

Location: 1100 feet N and 2220 feet E of the SW corner of Sec. 5, T.152N., R.29W.

Vegetation: Northern white cedar crown cover of about 45 percent with scattered

tamarack and black spruce; sparse understory consists of some willow.

dogwood, and horsetall; ground cover consists of some mosses and cranberry.

Depth (cm)

0-105

105 +

(NW1/4 of NE1/4 of SE1/4 of SW1/4, Sec. 5, T.152N., R.29W.)

Described By: D. Mellem and D. Haverkost on August 29, 1979.

Layer

Hemic

Fine

sand

Location: 330 feet N and 50 feet W of the SE corner of Sec. 3, T.152N., R.29W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 3, T.152N., R.29W.)

Vegetation: Scattered northern white cedar, ash, and black spruce; sparse understory consists of some grasses, Labrador tea, willow, dogwood, and daisies; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 30, 1979.

Layer	Depth (cm
Hemic Sandy clay with	0-210 210+

Location: 1760 feet S and 300 feet E of the NW corner of Sec. 6, T.152N., R.29W. (Gov. Lot 5, Sec. 6, T.152N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with bog birch and some tamarack, dogwood, Labrador tea, cotton grass, ferns, and cattails. Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Olson and D. Mellem on August 3, 1979.

Layer Depth (cm) Hemic 0-150 Sandv 150 +loam

Location: 1450 feet S and 1350 feet W of the NE corner of Sec. 7, T.152N., R.29W. (NE1/4 of NE1/4 of SW1/4 of NE1/4, Sec. 7, T.152N., R.29W.)

Location: 1745 feet S and 2050 feet W of the NE corner of Sec. 5, T.152N., R.29W.

Vegetation: Scattered black spruce, tamarack, and northern white cedar: understory consists of bog birch with some Labrador tea. leatherleaf, bog

Depth (cm)

0-190

190÷

rosemary, and daisies; ground cover consists of Sphagnum mosses.

(SE1/4 of NW1/4 of SW1/4 of NE1/4, Sec. 5, T.152N., R.29W.)

Described By: D. Mellem and D. Haverkost on August 29, 1979.

Laver

Hemic

Fine

sandy

clav

and medium

Vegetation: Northern white cedar crown cover of about 70 percent with scattered balsam fir, ash, and black spruce; understory consists of grasses with some speckled alder and dogwood; ground cover consists of some mosses.

Microrellef: 30 cm.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Depth To Water Table: 15 cm.

Described By: D. Mellem and D. Haverkost on August 29, 1979.

Layer	Depth (cm)
Hemic Fine	0-45 45+
sand	

Location: 1050 feet S and 250 feet E of the NW corner of Sec. 7, T.152N., R.29W. (Gov. Lot 1, Sec. 7, T.152N., R.29W.)

Vegetation: Northern white cedar crown cover of about 40 percent with scattered tamarack; understory consists of sedges with some Labrador tea, alder, balsam fir, and horsetail; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 29, 1979.

Laver Depth (cm) Hemic 0-45 45+ Fine sand

Location: 755 feet S and 75 feet W of the NE corner of Sec. 10, T.152N., R.29W. (NE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 10, T.152N., R.29W.)

Vegetation: Northern white cedar crown cover of about 40 percent with about 40 percent crown cover of ash; sparse understory consists of some balsam fir; grasses, and horsetail; ground cover consists mostly of mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 30, 1979.

Layer	Depth (cm	
Hemic	0-90	
Coarse	90+	
medium		
and		
fine		
sandy		
clav		

Location: 1375 feet N and 1325 feet W of the SE corner of Sec. 12, T.153N., R.29W. (SE1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 12, T.153N., R.29W.)

Vegetation: Scattered tamarack; sparse understory consists of some grasses, bog rosemary, willow, bog birch, fern, and cattail; ground cover consists of some Sphagnum mosses, other mosses, and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 22, 1979.

Layer	Depth (cm)
Hemic	0-191
Fine	191+
and	
medium	
sand	

Location: 100 feet N and 1400 feet W of the SE corner of Sec. 12, T.153N., R.29W. (SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 12, T.153N., R.29W.)

Vegetation: Scattered black spruce; understory consists of some bog birch, sedges, Labrador tea, leatherleaf, cattails, grasses, and ferns; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

other mosses and cranberry.

Depth To Water Table: At surface.

Microrellef: 30 cm.

Solomon's seal. Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 22, 1979.

Layer	Depth (cm
Hemic Sandy clay	0-252 252+

Location: 2300 feet S and 2000 feet E of the NW corner of Sec. 17, T.153N., R.29W.

Vegetation: Black spruce crown cover of about 65 percent; lush understory con-

sists mostly of Labrador tea with some grasses, leatherleaf, swamp laurel, and

cotton grass; ground cover consists mostly of Sphagnum mosses with some

(SE1/4 of SW1/4 of SE1/4 of NW1/4, Sec. 17, T.153N., R.29W.)

Described By: D. Mellem and D. Haverkost on August 21, 1979.

Layer

Hemic Silty clay Location: 1200 feet S and 775 feet E of the NW corner of Sec. 17, T.153N., R.29W. (SW1/4 of SE1/4 of NW1/4 of NW1/4, Sec. 17, T.153N., R.29W.)

Vegetation: Scattered tamarack; understory consists mostly of sedges with some grasses, willow, bog birch, Labrador tea, leatherleaf, bog rosemary, and horsetail; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 21, 1979.

Layer	Depth (cm)
Hemic	0-105
Fine	105+
and	
medium	
sand	

Location: 1200 feet N and 2580 feet W of the SE corner of Sec. 17, T.153N., R.29W. (SW1/4 of SW1/4 of NW1/4 of SE1/4, Sec. 17, T.153N., R.29W.)

Vegetation: Black spruce crown cover of about 20 percent with scattered tamarack; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, sedges, cotton grass, cattails, and bog birch; ground cover consists mostly of Sphagnum mosses with some other mosses and false Solomon's seal.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 21, 1979.

Depth (cm)		
0-145	Layer	Depth (cm)
145+	Fibric	0- 25
145+		
	Hemic	25-170
	Fine	170+
	and	
	medium	1
	sandy	•
	clay	

Location: 1230 feet S and 1290 feet E of the NW corner of Sec. 19, T.153N., R.29W. (SW1/4 of SW1/4 of NE1/4 of NW1/4, Sec. 19, T.153N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists mostly of leatherleaf with some cotton grass; ground cover consists mostly of Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem and D. Olson on August 3, 1979.

Layer Depth (cm)

Hemic 0-290
Fine 290+
and medium

Location: 50 feet N and 1200 feet W of the SE corner of Sec. 18, T.153N., R.29W. (SE1/4 of SE1/4 of SW1/4 of SE1/4, Sec. 18, T.153N., R.29W.)

Vegetation: Scattered black spruce; sparse understory consists of some Labrador

Described By: D. Mellem and D. Haverkost on August 21, 1979.

tea, leatherleaf, swamp laurel, bog rosemary, and cotton grass; ground cover

consists mostly of Sphagnum mosses with some cranberry and false

Fibric 0- 76 Hemic 76-427 Medium 427+ sand Location: 2275 feet S and 400 feet E of the NW corner of Sec. 17, T.153N., R.29W. (NE1/4 of SW1/4 of SW1/4 of NW1/4, Sec. 17, T.153N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of some Labrador tea, leatherleaf, swamp laurel, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 21, 1979.

Layer	Depth (cm)
Hemic	0-230
Sapric	230-235
Fine	235+
sand	

Location: 1560 feet N and 250 feet E of the SW corner of Sec. 18, T.153N., R.29W. (Gov. Lot 3, Sec. 18, T.153N., R.29W.)

Vegetation: Scattered black spruce; understory consists of some grasses, Labrador tea, swamp laurel, bog rosemary, and sedges.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem and D. Haverkost on August 21, 1979.

Layer	Depth (cm
Hemic	0-150
Fine	150+
and	
medium	
sand	

Location: 1375 feet N and 590 feet E of the SW corner of Sec. 19, T.153N., R.29W. (Gov. Lot 3, Sec. 19, T.153N., R.29W.)

Vegetation: Lush understory consists mostly of bog birch with bog rosemary and grasses.

Microrelief: Negligible.

Depth To Water Table: 30 cm. above surface.

Described By: D. Mellem and D. Olson on August 3, 1979.

Layer	Depth (cm
Hemic Fine	0-208 208+
eand	200 1

Location: 75 feet S and 1605 feet E of the NW corner of Sec. 20, T.153N., R.29W. (NW1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 20, T.153N., R.29W.)

Vegetation: Black spruce crown cover of about 30 percent; understory consists of some black spruce, tamarack, Labrador tea, leatherleaf, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 21, 1979.

Layer Depth (cm)
Hemic 0-243
Loam 243+

Location: 2560 feet N and 440 feet E of the SW corner of Sec. 21, T.153N., R.29W. (NE1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 21, T.153N., R.29W.)

Vegetation: Scattered tamarack crown cover; lush understory consists mostly of grasses and bog birch with some speckled alder and willow.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Meilem and D. Olson on August 3, 1979.

Layer	Depth (cm)
Sapric	0-61
Sand	61+

pth (cm)
Layer Depth (cm)

and aquatic plants.

Depth To Water Table: At surface.

Microrelief: Negligible.

Location: 655 feet S and 590 feet E of the NW corner of Sec. 26, T.153N., R.29W. (NE1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 26, T.153N., R.29W.)

Vegetation: Scattered black spruce crown cover; sparse understory consists of some black spruce, tamarack, northern white cedar, willow, bog birch, Labrador tea, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrellef: 30 cm.

cranberry.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 30, 1979.

Layer	Depth (cm)
Hemic	0-236
Sapric	236-244
Silty	244+
clay	

Location: 1750 feet S and 260 feet E of the NW corner of Sec. 26, T.153N., R.29W. (SW1/4 of NW1/4 of SW1/4 of NW1/4, Sec. 26, T.153N., R.29W.)

Vegetation: Scattered tamarack and black spruce crown cover; understory consists of some bog birch, tamarack, black spruce, Labrador tea, bog rosemary, ferns, and horsetail; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and pitcher plant.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 30, 1979.

Layer	Depth (cm)
Hemic Silty	0-223 223+
clay	

Location: 275 feet N and 200 feet E of the SW corner of Sec. 33. T.153N., R.29W.

(SW1/4 of SW1/4 of SW1/4, Sec. 33, T.153N., R.29W.)

Vegetation: Scattered black spruce, tamarack, and northern white cedar; understory consists mostly of bog birch with some Labrador tea; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 29, 1979.

Layer Depth (cm)

Hemic 0-90
Silty 90+
clay
with
mica

Location: 1585 feet N and 100 feet E of the SW corner of Sec. 33, T.153N., R.29W.

Vegetation: Scattered black spruce, tamarack, and northern white cedar; sparse

understory consists of some bog birch, dogwood, speckled alder, Labrador

tea, and reeds; ground cover consists mostly of Sphagnum mosses with some

(SW1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 33, T.153N., R.29W.)

Described By: D. Mellem and D. Haverkost on August 29, 1979.

Layer Depth (cm)
Hemic 0-135
Silty 135+
clay

Location: 2225 feet N and 80 feet E of the SW corner of Sec. 26, T.153N., R.29W. (SW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 26, T.153N., R.29W.)

Location: 1555 feet S and 1075 feet E of the NW corner of Sec. 24, T.153N., R.29W.

Vegetation: Scattered black spruce: lush understory consists mostly of sedges with

some bog birch and Phragmites; ground cover consists of some pitcher plant

0-244

244+

(NE1/4 of NE1/4 of SW1/4 of NW1/4, Sec. 24, T.153N., R.29W.)

Described By: D. Mellem and D. Olson on August 3, 1979.

Hemic

sand

Medium

Vegetation: Tamarack crown cover of about 40 percent with scattered black spruce; lush understory consists mostly of grasses and speckled alder with some bog birch, Labrador tea, and leatherleaf; ground cover consists of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 30, 1979.

Layer	Depth (cm)
Hemic	0-120
Sifty	120+
clay	

Location: 1450 feet S and 10 feet W of the NE corner of Sec. 34, T.153N., R.29W. (NE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 34, T.153N., R.29W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists of sedges with some bog birch, leatherleaf, Labrador tea, swamp laurel, ferns, and grasses; ground cover consists of some false Solomon's seal and cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 30, 1979.

Layer	Depth (cm
Hemic	0-150
Silty	150+
clav	

Location: 260 feet N and 1380 feet E of the SW corner of Sec. 35, T.153N., R.29W. (SW1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 35, T.153N., R.29W.)

Vegetation: Scattered tamarack and black spruce; sparse understory consists of some bog birch, Labrador tea, bog rosemary, cattail, and horsetail; ground cover consists mostly of Sphagnum mosses with some pitcher plant, cranberry, and false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on August 30, 1979.

Layer	Depth (cm)
Fibric	0- 45
Hemic	45-300
Silty	300+
clay	

Location: 1450 feet N and 250 feet W of the SE corner of Sec. 9 T.154N., R.29W. (SE1/4 of SE1/4 of NE1/4 of SE1/4, Sec. 9, T.154N., R.29W.)

Vegetation: Tamarack crown cover of about 25 percent; lush understory consists mostly of grasses with some willow.

Microrelief: Negligible.

Depth To Water Table: Not recorded.

Described By: D. Mellem and D. Olson on August 3, 1979.

Layer	Depth (cm)
Hemic Sandy	0-213 213+
loam	

Location: 2370 feet S and 830 feet E of the NW corner of Sec. 14, T.154N., R.29W. (SW1/4 of SE1/4 of SW1/4 of NW1/4, Sec. 14, T.154N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with some tamarack, northern white cedar, bog birch, horsetail, and Phragmites.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Olson on August 3, 1979.

Layer	Depth (cm)
Hemic	0-274
Fibric	274-427
Loam	427+

Location: 1260 feet N and 1850 feet E of the SW corner of Sec. 14, T.154N., R.29W. (NE1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 14, T.154N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with bog birch, Phragmites and some tamarack; ground cover consists of some pitcher plant and aquatic plants.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Olson on August 3, 1979.

Layer	Depth (cm)
Hemic Bottom unknown	0-488+

Location: 2360 feet S and 1055 feet W of the NE corner of Sec. 2, T.155N., R.29W. (SW1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 2, T.155N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of swamp laurel with some leatherleaf and sedges; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 21, 1979.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-135
Silty	135+
clay	
with	
calcareous	
pebbles	

Location: 1480 feet N and 1780 feet E of the SW corner of Sec. 2, T.155N., R.29W. (SE1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 2, T.155N., R.29W.)

Vegetation: Black spruce crown cover of about 50 percent; understory consists of some Labrador tea, leatherleaf, black spruce, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 21, 1979.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-135
Silty	135+
clay	

Location: 2550 feet N and 700 feet E of the SW corner of Sec. 2, T.155N., R.29W. (NW1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 2, T.155N., R.29W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 21, 1979.

Layer	Depth (cm)	
Hemic Silty clay	0-90 90+	

Location: 355 feet N and 620 feet W of the SE corner of Sec. 2, T.155N., R.29W. (NE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 2, T.155N., R.29W.)

Vegetation: Scattered black spruce; understory consists of some Labrador tea, leatherleaf, swamp laurel, and grasses; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 21, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-115
Sapric	115-120
Fine	120+
and	
medium	
sand	

Location: 1875 feet S and 2375 feet E of the NW corner of Sec. 5, T.155N., R.29W. (SE1/4 of NE1/4 of SE1/4 of NW1/4, Sec. 5, T.155N., R.29W.)

Vegetation: Scattered black spruce; understory consists of some Labrador tea, leatherleaf, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal.

Microrellet: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem, D. Olson, and D. Haverkost on July 27, 1979.

Layer	Depth (cn
Fibric	0- 50
Hemic	50-244
Sand	244+

Location: 1865 feet N and 535 feet E of the SW corner of Sec. 5, T.155N., R.29W. (NF1/4 of SW1/4 of NW1/4 of SW1/4, Sec. 5, T.155N., R.29W.)

Vegetation: Black spruce crown cover of about 50 percent; understory consists of leatherleaf with some black spruce, Labrador tea, and cotton grass; ground cover consists mostly of Sphagnum mosses with some false Solomoh's seal and cranberry.

Microrellef: 25 cm.

Depth To Water Table: 10 cm.

seal and cranberry.

Depth To Water Table: At surface.

Microrellef: 20 cm.

and blueberry.

Depth To Water Table: 10 cm.

Microrellef: 15 cm.

Described By: D. Mellem, D. Olson, and D. Haverkost on July 26, 1979.

Laver Depth (cm) Fibric 0-122 122-238 Hemic Sandy 238+ loam

Location: 1875 feet N and 1650 feet W of the SE corner of Sec. 5, T.155N., R.29W. (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 5, T.155N., R.29W.)

Vegetation: Black spruce crown cover of about 45 percent; understory consists of some black spruce. Labrador tea, swamp laurel, leatherleaf, and cotton grass: ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem, D. Olson, and D. Haverkost on July 26, 1979.

Layer	Depth (cm)
Hemic	0-213
Sandy	213+
loom '	

Location: 1875 feet N and 2050 feet E of the SW corner of Sec. 6, T.155N., R.29W.

Vegetation: Scattered black spruce: sparse understory consists of some cotton grass, leatherleaf, and swamp laurel; ground cover consists mostly of

Microrelief: 10 cm.

Layer Depth (cm) **Fibric** 0- 75 Hemic 75-250 Sapric 250-270 Coarse 270+ sand

Location: 1860 feet N and 1225 feet W of the SE corner of Sec. 6. T.155N., R.29W.

Vegetation: Black spruce crown cover of about 30 percent; lush understory con-

sists mostly of Labrador tea with some leatherleaf and cotton grass; ground

cover consists mostly of Sphagnum mosses with some false Solomon's seal

229+

(NW1/4 of SW1/4 of NE1/4 of SE1/4, Sec. 6, T.155N., R.29W.)

Described By: D. Mellem, D. Oison, and D. Haverkost on July 26, 1979.

Layer

Hemic

Sandy

loam

Location: 1800 feet S and 10 feet W of the NE corner of Sec. 6, T.155N., R.29W.

Vegetation: Scattered black spruce: sparse understory consists of some sedges.

swamp laurel, Labrador tea, leatherleaf, bog rosemary, and cotton grass;

ground cover consists mostly of Sphagnum mosses with some false Solomon's

(SE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 6, T.155N., R.29W.)

Described By: D. Mellem, D. Olson, and D. Haverkost on July 27, 1979.

(NW1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 6, T.155N., R.29W.)

Sphagnum mosses with some false Solomon's seal.

Depth To Water Table: At surface.

Described By: D. Mellem, D. Olson, and D. Haverkost on July 26, 1979.

Layer	Depth (cm
Hemic	0-138
Sand	138+

Location: 100 feet S and 150 feet E of the NW corner of Sec. 6, T.155N., R.29W. (Gov. Lot 4, Sec. 6, T.155N., R.29W.)

Vegetation: Scattered black spruce; sparse understory consists of some sedges, bog rosemary, swamp laurel, leatherleaf, Labrador tea, and cotton grass; ground cover consists mostly of Sphagnum mosses with some cranberry. Microrellef: 35 cm.

Depth (cm)

0-270

270 +

Depth To Water Table: At surface.

Described By: D. Mellem, D. Olson, and D. Haverkost on July 26, 1979.

Layer Depth (cm) Hemic 0-229 Sandy loam with pebbles Location: 200 feet S and 425 feet E of the NW corner of Sec. 33, T.155N., R.29W. (NE1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 33, T.155N., R.29W.)

Vegetation: Scattered black spruce; understory consists of bog birch, leatherleaf, and cotton grass with some willow; ground cover consists mostly of Sphagnum

Microrellef: 60 cm.

Layer	Depth (cm)
Hemic	0-143
Silt	143+
loam	
with	
pebbles	

Hemic 0-150 Sand 150 +

Layer

(Gov. Lot 6, Sec. 6, T.155N., R.29W.)

and cranberry.

Depth To Water Table: At surface.

Microrellef: 10 cm.

with calcareous

Described By: D. Mellem, D. Olson, and D. Haverkost on July 28, 1979.

Location: 2050 feet N and 25 feet E of the SW corner of Sec. 6, T.155N., R.29W.

Vegetation: Scattered black spruce and tamarack; understory consists of cotton grass with some sedges, swamp laurel, leatherleaf, and Labrador tea; ground

cover consists mostly of Sphagnum mosses with some false Solomon's seal

Depth (cm)

Location: 1750 feet S and 2375 feet W of the NE corner of Sec. 6, T.155N., R.29W.

Vegetation: Scattered black spruce; understory consists of some Labrador tea,

leatherleaf, swamp laurel, and cotton grass; ground cover consists mostly of

Depth (cm)

0-213

213+

(SW1/4 of NW1/4 of SW1/4 of NE1/4, Sec. 6, T.155N., R.29W.)

Described By: D. Mellem, D. Olson, and D. Haverkost on July 27, 1979.

Sphagnum mosses with some false Solomon's seal.

Layer

Hemic

Sand

Microrellef: 20 cm.

Depth To Water Table: At surface.

pebbles

mosses.

Depth To Water Table: 15 cm.

Described By: T. Malterer, R. Zarth, and M. Domeier on August 3, 1976.

Location: 850 feet S and 2000 feet E of the NW corner of Sec. 33. T.155N.. R.29W (NW1/4 of SE1/4 of NE1/4 of NW1/4. Sec. 33. T.155N.. R.29W.)

Vegetation: Scattered tamarack; understory consists of bog birch, bog rosemary, leatherleaf, and cotton grass with some cattails and willow: ground cover consists mostly of Sphagnum mosses.

Microrellef: 60 cm.

Depth To Water Table: 15 cm.

Described By: T. Malterer, R. Zarth, and M. Domeier on August 3, 1976.

Layer Depth (cm)
Hemic 0-236
Clay 236+
loam

Location: 2100 feet S and 1010 feet W of the NE corner of Sec. 34, T.155N., R.29W. (NE1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 34, T.155N., R.29W.)

Vegetation: Tamarack crown cover of about 40 percent; understory consists of cotton grass with some leatherleaf and bog birch; ground cover consists of some Sphaonum mosses.

Microrelief: Not recorded.

Depth To Water Table: At surface.

Described By: T. Malterer, R. Zarth, and M. Domeier on August 3, 1976.

	m)
Fibric 0- 61 Hemic 61-305 Sand 305+	

Location: 1050 feet S and 50 feet E of the NW corner of Sec. 34, T.155N., R.29W. (SW1/4 of SW1/4 of NW1/4 of NW1/4, Sec. 34, T.155N., R.29W.) Vegetation: Consists of tamarack, bog rosemary, leatherleaf, bog birch, and cotton

Vegetation: Consists of tamarack, bog rosemary, leatherleaf, bog birch, and cot grass; ground cover consists mostly of Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: 15 cm.

Described By: T. Malterer, R. Zarth, and M. Domeier on August 3, 1976.

Layer	Depth (cm)
Hemic	0-228
Silty	228+
clay	
loam	
with	
pebbles	

Location: 2100 feet N and 2425 feet E of the SW corner of Sec. 35, T.155N., R.29W. (SE1/4 of NE1/4 of NE1/4 of SW1/4, Sec. 35, T.155N., R.29W.)

Vegetation: Scattered black spruce, tamarack, and northern white cedar; understory consists of bog birch, cotton grass, and grasses; ground cover consists of some pitcher plant.

Microrellef: Not recorded.

Depth To Water Table: At surface.

Described By: T. Malterer, R. Zarth, and M. Domeier on August 3, 1976.

Layer	Depth (cm)
Hemic Clay	0-259 259+
Clay	2097

Location: 1875 feet N and 2425 feet E of the SW corner of Sec. 35, T.155N., R.29W. (NE1/4 of SE1/4 of NE1/4 of SW1/4, Sec. 35, T.155N., R.29W.)

Vegetation: Crown cover consists of black spruce and cedar with scattered tamarack; understory consists of Labrador tea; ground cover consists of some cranberry.

Microrellef: Not recorded.

Depth To Water Table: 20 cm.

Described By: T. Malterer, R. Zarth, and M. Domeier on August 3, 1976.

Layer	Depth (cm)
Hemic	0-295
Clay	295-315
Sandy	315+
inam	

Location: 2425 feet N and 710 feet E of the SW corner of Sec. 35, T.155N., R.29W. (NW1/4 of NE1/4 of NW1/4 of SW1/4, Sec. 35, T.155N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of Labrador tea, bog rosemary, leatherleaf, and bog birch; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: T. Malterer, R. Zarth, and M. Domeier on August 3, 1976.

Layer	Depth (cm)
Hemic Sandy clay loam	0-457 457+

Location: 420 feet S and 990 feet W of the NE corner of Sec. 5, T.156N., R.29W. (SW1/4 of NW1/4 of NE1/4 of NE1/4, Sec. 5, T.156N., R.29W.)

Vegetation: Scattered black spruce; lush understory consists mostly of cotton grass with some bog birch, leatherleaf, and bog rosemary; ground cover consists of some pitcher plant.

Microrellef: Not recorded.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on August 1, 1979.

Hemic 0-254 Sandy 254+ loam Location: 250 feet S and 250 feet E of the NW corner of Sec. 6, T.156N., R.29W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 6, T.156N., R.29W.)

Vegetation: Lush understory consists mostly of cotton grass, cattails, and bog birch with some dogwood and leatherleaf; ground cover consists of some Sphagnum mosses and cranberry.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on August 1, 1979.

Layer	Depth (cm)
Hemic Sandy	0-211 211+
loem	

Location: 60 feet N and 790 feet W of the SE corner of Sec. 7, T.156N., R.29W. (SE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 7, T.156N., R.29W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges with some leatherleaf, Labrador tea, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and D. Mellem on August 1, 1979.

n)

Location: 20 feet N and 1990 feet E of the SW corner of Sec. 8, T.156N., R.29W. (SE1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 8, T.156N., R.29W.)

Vegetation: Lush understory consists mostly of sedges with some black spruce; ground cover consists mostly of Sphagnum mosses.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Olson and D. Haverkost on August 1, 1979.

Depth (cm) Laver Fibric 0- 91 Hemic 91-396 Sand 396+

Location: 75 feet N and 1910 feet E of the SW corner of Sec. 10, T.156N., R.29W.

Vegetation: Lush understory consists mostly of sedges with some black spruce, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of

Depth (cm)

0- 61

61-389 389-414

414+

(SE1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 10, T.156N., R.29W.)

Sphagnum mosses with some cranberry.

Described By: T. Malterer and D. Mellem on August 1, 1979.

Layer

Fibric Hemic

Sapric Sandy

loam

Microrellef: 15 cm.

Depth To Water Table: At surface.

Location: 75 feet N and 1000 feet W of the SE corner of Sec. 8, T.156N., R.29W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 8, T.156N., R.29W.)

Vegetation: Sparse understory consists of some sedges, leatherleaf, black spruce. and tamarack; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrellef: 10 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and D. Mellem on August 1, 1979.

Layer	Depth (cn
Fibric	0- 15
Hemic	15-302
Sapric	302-325
Silty	325+
clay	
loam	

Location: 100 feet N and 1975 feet W of the SE corner of Sec. 10, T.156N., R.29W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 10, T.156N., R.29W.)

Vegetation: Not recorded.

Depth To Water Table: Not recorded.

Microrelief: Not recorded.

Described By: D. Olson and D. Haverkost on August 1, 1979.

Depth (cm)
0- 91
91-335
335+

Location: 90 feet N and 110 feet W of the SE corner of Sec. 11, T.156N., R.29W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 11, T.156N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of some sedges, bog birch, leatherleaf, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: T. Malterer and D. Mellem on August 1, 1979.

Laver Depth (cm) Hemic 0-244 Sapric 244-272 Loamy 272+ sand

Location: 400 feet N and 710 feet W of the SE corner of Sec. 12, T.156N., R.29W. (NE1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 12, T.156N., R.29W.)

Vegetation: Lush understory consists mostly of sedges with wild rice, cattails, bog birch, and some tamarack, leatherleaf, bog rosemary, horsetail, fern, and pitcher plant.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on August 1, 1979.

Layer	Depth (cm)
Hemic	0-264
Sand	264+

Location: 50 feet N and 2350 feet W of the SE corner of Sec. 9, T.156N., R.29W. (SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 9, T.156N., R.29W.)

Vegetation: Sparse understory consists of some bog rosemary; ground cover consists mostly of Sphagnum mosses with some pitcher plant.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on August 1, 1979.

Layer	Depth (cm)
Fibric	0- 38
Hemic	38-330
Sandy	330+
loam	•

Location: 110 feet N and 50 feet W of the SE corner of Sec. 10, T.156N., R.29W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 10, T.156N., R.29W.)

Vegetation: Understory consists of sedges with some black spruce, tamarack, bog birch, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and D. Mellem on August 1, 1979.

Layer	Depth (cm)
Hemic	0-246
Fine	246+
medium	
and	
coarse	
Saliu	

Location: 610 feet S and 2560 feet W of the NE corner of Sec. 14, T.156N., R.29W. (SW1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 14, T.156N., R.29W.)

Vegetation: Not recorded.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Olson and D. Haverkost on August 1, 1979.

Layer	Depth (cm)
Sapric	0- 91
Hemic	91-323
Sand	323+

Location: 1325 feet N and 330 feet E of the SW corner of Sec. 23, T.156N., R.29W. (NW1/4 of NW1/4 of SW1/4 of SW1/4. Sec. 23, T.156N., R.29W.)

Vegetation: Understory consists mostly of grasses and cattails with some bog birch, tamarack, willow, and bog rosemary; ground cover consists of some masses.

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: T. Malterer and D. Mellem on August 2, 1979.

Layer Depth (cm)
Hemic 0-300
Sandy 300+
loam

Location: 425 feet S and 90 feet E of the NW corner of Sec. 25, T.156N., R.29W. (SW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 25, T.156N., R.29W.)

Vegetation: Scattered black spruce and tamarack crown cover; understory consists mostly of black spruce, tamarack, and grasses with some willow and dogwood; ground cover consists mostly of Sphagnum and other mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 21, 1979.

yer	Depth (cm)
e	0-70 70-75 75+
nd	

Layer Depth (cm)
Hemic 0-254

Described By: D. Olson and D. Haverkost on August 2, 1979.

grass and some bog birch and bog rosemary.

Fine

sand

Location: 930 feet S and 375 feet W of the NE corner of Sec. 27, T.156N., R.29W.

Vegetation: Scattered tamarack; understory consists mostly of cattails with cotton

254 +

(NW1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 27, T.156N., R.29W.)

Location: 750 feet N and 1325 feet W of the SE corner of Sec. 32, T.156N., R.29W. (SE1/4 of NE1/4 of SW1/4 of SE1/4, Sec. 32, T.156N., R.29W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some sedges, Labrador tea, leatherleaf, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal and cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Olson, D. Mellem, and D. Haverkost on July 27, 1979.

Hemic 0-165 Sandy 165+ loam Location: 525 feet N and 125 feet E of the SW corner of Sec. 7, T.157N., R.29W. (Gov. Lot 4, Sec. 7, T.157N., R.29W.)

Vegetation: Lush understory consists mostly of grasses and cattails; ground cover consists of some IIIy pads.

Microrellef: Negligible.

Depth To Water Table: 30 cm. above surface.

Described By: T. Malterer and D. Mellem on August 2, 1979.

Layer Depth (cm)
Hemic 0-15
Medium 15+
and
fine
sand

Location: 300 feet N and 260 feet W of the SE corner of Sec. 10, T.157N., R.29W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 10, T.157N., R.29W.)

Vegetation: Lush understory consists mostly of sedges with some grasses, speckled alder, willow, dogwood, cattail, and raspberry.

Microrelief: Negligible.

Microrelief: 10 cm.

Depth To Water Table: At surface.

Depth To Water Table: Not visible.

Described By: D. Olson and D. Haverkost on August 2, 1979.

Layer	Depth (cm)
Hemic	0-61
Sand	61+

Location: 205 feet N and 225 feet E of the SW corner of Sec. 12, T.157N., R.29W. (SW1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 12, T.157N., R.29W.)

Vegetation: Not recorded. Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: T. Malterer and D. Mellem on August 2, 1979.

Sapric 0-20 Loamy 20+ sand Location: 220 feet S and 625 feet W of the NE corner of Sec. 16, T.157N., R.29W. (NW1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 16, T.157N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with speckled alder and some bog birch, gooseberry, and raspberry.

Microrelief: 15 cm.

Denth To Water Table: At surface.

Described By: T. Malterer and D. Mellem on August 2, 1979.

Layer Depth (cm)

Hemic 0-46

Fine 46+

and

medium

sand

Location: 475 feet S and 330 feet E of the NW corner of Sec. 17, T.157N., R.29W. (SW1/4 of NW1/4 of NW1/4, Sec. 17, T.157N., R.29W.)

Vegetation: Scattered tamarack; lush understory consists mostly of sedges with some speckled alder, bog birch, and willow.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on August 2, 1979.

Layer	Depth (cm)
Hemic Sandy clay	0-86 86+

Vegetation: Lush understory consists mostly of speckled alder, dogwood, and grasses with some leatherleaf.

Microrellef: 10 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on November 13, 1979.

Layer Depth (cm) Hemic 0-91 Sand 91+

Vegetation: Understory consists of bog birch with some black spruce, leatherleaf, Labrador tea, swamp laurel, cotton grass, and grasses; ground cover consists mostly of Sphagnum mosses with some cranberry and false Solomon's seal

Location: 150 feet N and 500 feet W of the SE corner of Sec. 33, T.157N., R.29W.

(SW1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 33, T.157N., R.29W.)

Microrelief: Not recorded.

Depth To Water Table: Not recorded.

Described By: T. Malterer and D. Mellem on August 1, 1979.

Laver Depth (cm) Hemic 0-130 Sanric 130-143 Loamy 143+ sand

Layer Depth (cm) Hemic 0-107 Loamy 107+ sand

Location: 1800 feet S and 200 feet W of the NE corner of Sec. 24, T.157N., R.29W.

Vegetation: Lush understory consists mostly of grasses with speckled alder and

(SE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 24, T.157N., R.29W.)

Described By: T. Malterer and D. Mellem on August 2, 1979.

Location: 1240 feet S and 210 feet W of the NE corner of Sec. 34, T.157N., R.29W. (SE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 34, T.157N., R.29W.)

Vegetation: Lush understory consists of bog birch, sedges, and leatherleaf. Microrellef: 20 cm.

Depth To Water Table: Not recorded.

some willow and goldenrod.

Depth To Water Table: At surface

Microrelief: 10 cm.

Described By: D. Mellem and R. Graber on November 13, 1979.

Layer Depth (cm) Hemic 0-122 Loam 122+

Location: 2500 feet S and 825 feet W of the NE corner of Sec. 28, T.157N., R.29W. (SE1/4 of SW1/4 of SE1/4 of NE1/4, Sec. 28, T.157N., R.29W.)

Vegetation: Scattered tamarack and black spruce; understory consists of some grasses, sedges, Labrador tea, and bog birch; ground cover consists mostly of Sphagnum mosses.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on November 13, 1979.

Layer	Depth (cm
Hemic	0-117
Sand	117+

Location: 160 feet N and 350 feet E of the SW corner of Sec. 35, T.157N., R.29W. (SE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 35, T.157N., R.29W.)

Vegetation: Lush understory consists mostly of sedges with leatherleaf and some bog birch, speckled alder, willow, and tamarack; ground cover consists mostly of Sphagnum mosses with some false Solomon's seal.

Microrellef: 20 cm.

Depth To Water Table: At surface.

Described By: D. Mellem on August 1, 1979.

Layer	Depth (cm
Hemic	0-152
Sapric	152-168
Sandy	168+
clay	
loam	

Location: 625 feet S and 225 feet W of the NE corner of Sec. 36, T.157N., R.29W. (SE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 36, T.157N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with bog birch, leatherleaf, and some bog rosemary; ground cover consists of some mosses. Microrellef: 38 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and D. Mellem on August 2, 1979.

Laver Depth (cm) Hemic 0-130 Loamy 130 +sand

Location: 1725 feet N and 1450 feet E of the SW corner of Sec. 1, T.158N., R.29W. (NW1/4 of SW1/4 of NE1/4 of SW1/4, Sec. 1, T.158N., R.29W.)

Vegetation: Understory consists of leatherleaf with some black spruce, bog rosemary, Labrador tea, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, pitcher plant, and lichens.

Microrellef: 50 cm.

Depth To Water Table: 20 cm.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer Depth (cm) Hemic 0-198 Silty 198+ clay

Location: 450 feet N and 1745 feet E of the SW corner of Sec. 2, T.158N., R.29W. (NE1/4 of SW1/4 of SE1/4 of SW1/4, Sec. 2, T.158N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf with some Labrador tea, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and lichens.

Microrellef: 40 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Olson on October 30, 1979.

Layer	Depth (cm)
Hemic	0-198
Clay	198+

Location: 1085 feet S and 2010 feet W of the NE corner of Sec. 5, T.158N., R.29W. (SE1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 5, T.158N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of some leatherleaf, swamp laurel, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 25 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer	Depth (cm
Fibric	0- 61
Hemic	61-170
Clay	170+

Location: 125 feet S and 2040 feet W of the NE corner of Sec. 6, T.158N., R.29W. (Gov. Lot 2, Sec. 6, T.158N., R.29W.)

Vegetation: Understory consists of willow with some bog birch, aspen, and cattail.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Olson and D. Haverkost on September 19, 1979.

Layer	Depth (cm)
Sapric	0-84
Clay	84+

Location: 2410 feet S and 1330 feet W of the NE corner of Sec. 6, T.158N., R.29W. (SE1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 6, T.158N., R.29W.)

Vegetation: Tamarack crown cover of about 30 percent with scattered aspen; lush understory consists mostly of bog birch and bog rosemary with some willow, Labrador tea, leatherleaf, grasses, and Phagmites; ground cover consists mostly of Sphagnum mosses.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on September 19, 1979.

Layer	Depth (cm
Hemic	0-198
Sapric	198-199
Silty	199+
clav	

Location: 720 feet S and 2075 feet W of the NE corner of Sec. 7, T.158N., R.29W. (NE1/4 of SW1/4 of NW1/4 of NE1/4, Sec. 7, T.158N., R.29W.)

Vegetation: Lush understory consists mostly of bog birch and leatherleaf with some poplar, alder, Labrador tea, and grasses.

Microrellef: 15 cm.

Depth To Water Table: 15 cm.

Described By: D. Mellem and D. Haverkost on September 18, 1979.

Layer	Depth (cm)
Hemic	0-166
Sapric	166-170
Silty	170+
clav	

Location: 750 feet S and 1400 feet W of the NE corner of Sec. 7, T.158N., R.29W. (NE1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 7, T.158N., R.29W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists of bog birch and bog rosemary with some leatherieaf, grasses, and Labrador tea; ground cover consists of Sphagnum and other mosses.

Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on September 18, 1979.

Layer	Depth (cm)
Hemic	0-165
Sapric	165-173
Silty	173+
clay	

Location: 190 feet N and 10 feet W of the SE corner of Sec. 7, T.158N., R.29W. (SE1/4 of SE1/4 of SE1/4 of SE1/4. Sec. 7, T.158N., R.29W.)

Vegetation: Scattered tamarack and black spruce; lush understory consists mostly of sedges with some grasses, willow, leatherleaf, and ferns; ground cover consists of mosses with some false Solomon's seal.

Microrelief: 15 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Layer	Depth (cm)
Fibric Silty clay with pebbles	0-151 151+

Location: 125 feet S and 70 feet W of the NE corner of Sec. 8, T.158N., R.29W. (NE1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 8, T.158N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with some bog birch, alder, willow, sedges, leatherleaf, ferns, and aster; ground cover consists of some mosses.

Microrelief: 10 cm.

Depth To Water Table: 35 cm.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer	Depth (cm
Hemic Sand	0-244 244+

Location: 495 feet N and 550 feet W of the SE corner of Sec. 8, T.158N., R.29W. (NW1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 8, T.158N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with willow and some cattails.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Layer	Depth (cm)
Hemic	0-151
Sapric	151-166
Loam	166+

Location: 45 feet N and 1300 feet W of the SE corner of Sec. 9, T.158N., R.29W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 9, T.158N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with some sedges and willow; ground cover consists of some mosses.

Microrellef: 5 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer	Depth (cm)
Hemic	0-152
Silty	152+
clay	
loam	

Lecation: 75 feet N and 1000 feet W of the SE corner of Sec. 11, T.158N., R.29W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 11, T.158N., R.29W.)

Vegetation: Lush understory consists mostly of bog birch, grasses, and leatherleaf. Microrelief: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Layer	Depth (cm
Sapric	0- 30
Hemic	30-151
Silty	151+
clay	

Location: 135 feet N and 190 feet W of the SE corner of Sec. 12, T.158N., R.29W. (SE1/4 of SE1/4 of SE1/4 of SE1/4. Sec. 12, T.158N., R.29W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of grasses with leatherleaf and some willow and cattail.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Layer	Depth (cm)
Hemic	0-210
Loam	210+

Location: 1900 feet S and 140 feet W of the NE corner of Sec. 13, T.158N., R.29W. (SE1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 13, T.158N., R.29W.)

Vegetation: Lush understory consists mostly of sedges and grasses with some willow and aster; ground cover consists of some mosses.

Microrellef: 10 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer	Depth (cm)
Hemic	0-183
Sandy	183+
loom	

Location: 175 feet N and 700 feet E of the SW corner of Sec. 20, T.158N., R.29W. (SE1/4 of SW1/4 of SW1/4 of SW1/4, Sec. 20, T.158N., R.29W.)

Vegetation: Not recorded.
Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Described By: D. Mellem and D. Haverkost on September 19, 1979.

Depth (cm)
0-20 20+

Location: 900 feet N and 575 feet E of the SW corner of Sec. 21, T.158N., R.29W. (SE1/4 of NW1/4 of SW1/4 of SW1/4, Sec. 21, T.158N., R.29W.)

Vegetation: Lush understory consists mostly of grasses.

Microrellef: 5 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer	Depth (cm)
Hemic	0-5
Loam	5+

Location: 1850 feet S and 2340 feet W of the NE corner of Sec. 31, T.158N., R.29W. (SE1/4 of NW1/4 of SW1/4 of NE1/4, Sec. 31, T.158N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with some willow, horsetail, cattail, and cotton grass; ground cover consists of some Sphagnum mosses and other mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: T. Malterer and D. Mellem on August 2, 1979.

Layer	Depth (cm)
Sapric Fine and medium sand	0-8 8+

Location: 20 feet N and 280 feet E of the SW corner of Sec. 31, T.158N., R.29W. (Gov. Lot 4, Sec. 31, T.158N., R.29w.)

Vegetation: Lush understory consists mostly of sedges with some willow, speckled alder, and cattails.

Microrellef: Negligible.

Depth To Water Table: At surface.

Described By: D. Olson and D. Haverkost on August 2, 1979.

Sapric 0-122 Sand 122+ Location: 500 feet S and 150 feet W of the NE corner of Sec. 1, T.159N., R.29W. (Gov. Lot 1, Sec. 1, T.159N., R.29W.)

Vegetation: Lush understory consists mostly of grasses with some alder, dogwood, poplar, ferns, and aster.

Microrellef: 5 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Mellem on November 1, 1979.

Layer	Depth (cm)
Hemic Medium sand with pebbles	0-15 15+

Location: 2000 feet N and 360 feet E of the SW corner of Sec. 1. T.159N., R.29W. (SE1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 1, T.159N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of cotton grass with some leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some cranberry, pitcher plant, and false Solomon's seal.

Microrelief: 15 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Olson on October 30, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-335
Clay	335+
loam	

Location: 375 feet S and 210 feet W of the NE corner of Sec. 2, T.159N., R.29W. (Gov. Let 1, Sec. 2, T.159N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists mostly of sedges with some leatherleaf and bog rosemary; ground cover consists mostly of Sphagnum mosses with some pitcher plant.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm
Hemic	0- 60
Fibric	60-180
Hemic	180-318
Silt	318+

Location: 100 feet S and 90 feet E of the NW corner of Sec. 8, T.159N., R.29W. (NW1/4 of NW1/4 of NW1/4 of NW1/4, Sec. 8, T.159N., R.29W.)

Vegetation: Scattered black spruce; lush understory consists mostly of cotton grass with some leatherleaf, Labrador tea, and swamp laurel; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrellef: 25 cm.

Depth To Water Table: 35 cm.

Vegetation: Consists mostly of grasses.

Depth To Water Table: At surface.

Microrellef: 10 cm.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-229
Very	229+
fine	
sand	

Location: 1115 feet N and 710 feet W of the SE corner of Sec. 10, T.159N., R.29W.

Depth (cm)

60+

(NE1/4 of NW1/4 of SE1/4 of SE1/4, Sec. 10, T.159N., R.29W.)

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer

Hemic

Location: 60 feet N and 50 feet W of the SE corner of Sec. 5, T.159N., R.29W. (SE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 5, T.159N., R.29W.)

Vegetation: Scattered black spruce; sparse understory consists of some swamp laurel, sedges, grasses, leatherleaf, and Labrador tea; ground cover consists mostiv of mosses with some Sphagnum mosses.

Microrelief: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Layer	Depth (cm)
Fibric	0- 30
Hemic	30-151
Silty	151+
clav	

Location: 235 feet N and 810 feet E of the SW corner of Sec. 8, T.159N., R.29W. (SW1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 8, T.159N., R.29W.)

Vegetation: Scattered black spruce; understory consists mostly of leatherleaf with some swamp laurel, cotton grass, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrellef: 20 cm.

Depth To Water Table: 40 cm.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-259
Clay	259+
loom	

Location: 75 feet N and 2525 feet E of the SW corner of Sec. 11, T.159N., R.29W. (SE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 11, T.159N., R.29W.)

Vegetation: Scattered tamarack; lush understory consists mostly of sedges with some bog birch, speckled alder, swamp laurel, leatherleaf, ferns, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 10 cm.

Depth To Water Table: 10 cm.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Hemic	0-172
√ery	172+
ine	
sand	
and	
tlis	

Location: 1190 feet S and 1690 feet W of the NE corner of Sec. 7, T.159N., R.29W. (SW1/4 of SE1/4 of NW1/4 of NE1/4, Sec. 7, T.159N., R.29W.)

Vegetation: Scattered black spruce; sparse understory consists of some Labradortea, leatherleaf, swamp laurel, bog rosemary, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Layer	Depth (cm)
Fibric	0-135
Hemic	135-196
Loam	196+

Location: 1225 feet S and 130 feet W of the NE corner of Sec. 10, T.159N., R.29W. (SE1/4 of SE1/4 of NE1/4 of NE1/4, Sec. 10, T.159N., R.29W.)

Vegetation: Black spruce crown cover of about 60 percent with scattered tamarack; lush understory consists mostly of grasses with some willow. Microrellef: 45 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-181 181+

Location: 650 feet S and 2100 feet W of the NE corner of Sec. 12, T.159N., R.29W. (SE1/4 of NW1/4 of NW1/4 of NE1/4, Sec. 12, T.159N., R.29W.)

Vegetation: Black spruce crown cover of about 45 percent; sparse understory consists of some black spruce and grasses; ground cover consists of some Sphagnum and other mosses.

Microrellef: 15 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Hemic Siltv	0-120 120+
clay	1201

Location: 875 feet N and 2050 feet E of the SW corner of Sec. 12, T.159N., R.29W. (SE1/4 of NW1/4 of SE1/4 of SW1/4. Sec. 12, T.159N., R.29W.)

Vegetation: Scattered black spruce; lush understory consists mostly of grasses with some bog birch, bog rosemary, and cotton grass; ground cover consists of some mosses.

Microrellef: 15 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer Depth (cm)
Hemic 0-240
Silty 240+
clay

Location: 750 feet N and 1230 feet E of the SW corner of Sec. 20, T.159N., R.29W.

Vegetation: Scattered black spruce; lush understory consists of leatherleaf and

cotton grass with some Labrador tea and swamp laurel; ground cover consists

mostly of Sphagnum mosses with some other mosses, cranberry, and false

Depth (cm)

0- 61

61-162

162+

(SE1/4 of NE1/4 of SW1/4 of SW1/4, Sec. 20, T.159N., R.29W.)

Described By: D. Olson and N. Aaseng on October 29, 1979.

Laver

Fibric

Clay

iosm

Hemic

Location: 810 feet N and 145 feet W of the SE corner of Sec. 15, T.159N., R.29W. (SE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 15, T.159N. R.29W.)

Vegetation: Scattered black spruce; understory consists of sedges with some leatherleaf, speckled alder, grasses, and cattall; ground cover consists mostly of Sphagnum mosses with some other mosses and false Solomon's seal.

Microrelief: 10 cm.

Depth To Water Table: 5 cm.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Hemic	0-61
Sapric	61-91
Sandy	91+
clay	
loam	

Location: 140 feet N and 640 feet W of the SE corner of Sec. 22, T.159N., R.29W. (SW1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 22, T.159N., R.29W.)

Vegetation: Scattered tamarack; lush understory consists mostly of leatherleaf with some Labrador tea and black spruce; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 45 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Hemic	0-90
Clay	90+

Location: 1690 feet N and 1640 feet W of the SE corner of Sec. 23, T.159N., R.29W. (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 23, T.159N., R.29W.)

Location: 1375 feet S and 225 feet W of the NE corner of Sec. 18, T.159N., R.29W.

Vegetation: Scattered black spruce; sparse understory consists of some Labrador

tea, leatherleaf, swamp laurel, bog rosemary, and sedges; ground cover con-

Depth (cm)

0- 90

90-302

302+

(NW1/4 of NE1/4 of SE1/4 of NE1/4, Sec. 18, T.159N., R.29W.)

sists mostly of Sphagnum mosses with some cranberry.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Laver

Fibric

Hemic

Sandy

clay

Vegetation: Scattered black spruce; sparse understory consists of some leatherleaf, Labrador tea, swamp laurel, bog rosemary, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrelief: 30 cm.

Microrellef: Not recorded.

Depth To Water Table: Not recorded.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Hemic	0-181
Silty	181+
clay	

Location: 1830 feet S and 90 feet W of the NE corner of Sec. 24, T.159N., R.29W. (Gov. Lot 2, Sec. 24, T.159N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of Labrador tea with some bog birch, willow, leatherleaf, swamp laurel, bog rosemary, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses, and lichens.

Microrellef: 30 cm.

Solomon's seal.

Depth To Water Table: Below 162 cm.

Microrellef: 25 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Hemic	0-122
Sandy	122+
clay	
loam	

Location: 850 feet N and 1380 feet E of the SW corner of Sec. 24, T.159N., R.29W. (NE1/4 of SE1/4 of NW1/4 of SW1/4, Sec. 24, T.159N., R.29W.)

Vegetation: Scattered black spruce; lush understory consists mostly of leatherleaf with some cotton grass and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-183
Clay	183+

Location: 80 feet S and 1090 feet E of the NW corner of Sec. 25, T.159N., R.29W. (NE1/4 of NE1/4 of NW1/4 of NW1/4, Sec. 25, T.159N., R.29W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of leatherleaf with some Labrador tea, bog rosemary, cotton grass, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and lichens.

Microrelief: 35 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-183 183+

Location: 2240 feet S and 1440 feet W of the NE corner of Sec. 32, T.159N., R.29W. (NE1/4 of SE1/4 of SW1/4 of NE1/4, Sec. 32, T.159N., R.29W.)

Vegetation: Scattered black spruce and tamarack; sparse understory consists of some Labrador tea, leatherleaf, swamp laurel, sedges, and grasses; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrellef: 30 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Depth (cm) Layer 0- 75 Fibric 75-175 Hemic Silty 175 +clav

Location: 460 feet N and 2575 feet E of the SW corner of Sec. 13, T.160N., R.29W.

Labrador tea; ground cover consists mostly of Sphagnum mosses with some

Depth (cm)

0-165

165 +

(NE1/4 of SE1/4 of SE1/4 of SW1/4, Sec. 13, T.160N., R.29W.) Vegetation: Black spruce crown cover of about 70 percent; understory consists of

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Laver

Hemic

Sandy

clav

cranberry and false Solomon's seal.

Depth To Water Table: At surface.

Microrelief: 30 cm.

Location: 700 feet N and 360 feet W of the SE corner of Sec. 34, T.159N., R.29W. (SE1/4 of NE1/4 of SE1/4 of SE1/4, Sec. 34, T.159N., R.29W.)

Vegetation: Black spruce crown cover of about 80 percent with scattered tamarack and poplar: Jush understory consists mostly of leatherleaf with Labrador tea: ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 30 cm.

Depth To Water Table: Below 151 cm.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Hemic Silty clay	0-151 151+

(NE1/4 of NW1/4 of SW1/4 of SE1/4, Sec. 14, T.160N., R.29W.)

Vegetation: Understory consists mostly of sedges with some black spruce. tamarack, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal.

Microrelief: 30 cm.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Depth (cm) Layer Fibric 0-120 Hemic 120-195 Sandy 195+

Location: 1275 feet N and 2030 feet W of the SE corner of Sec. 14, T.160N., R.29W.

and cranberry.

Depth To Water Table: At surface.

Location: 2000 feet N and 850 feet W of the SE corner of Sec. 16, T.160N., R.29W. (SE1/4 of NW1/4 of NE1/4 of SE1/4, Sec. 16, T.160N., R.29W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some bog rosemary, swamp laurel, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and pitcher plant.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-244
Clay	244+

Location: 120 feet N and 1060 feet W of the SE corner of Sec. 21, T.160N., R.29W. (SW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 21, T.160N., R.29W.)

Vegetation: Scattered black spruce; lush understory consists mostly of sedges with some bog rosemary, swamp laurel, and leatherleaf; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 20 cm.

Depth To Water Table: At surface.

Described By: D. Oison and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Fibric	0- 61
Hemic	61-106
Silty	106+
clay	

Location: 360 feet N and 1035 feet W of the SE corner of Sec. 9, T.160N., R.29W. (NW1/4 of SW1/4 of SE1/4 of SE1/4, Sec. 9, T.160N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of some Labrador tea, leatherleaf, bog rosemary, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and lichens.

Microrellef: 30 cm.

Depth To Water Table: 30 cm.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Hen :	0-101
Clay	101+

Location: 1825 feet N and 1975 feet W of the SE corner of Sec. 15, T.160N., R.29W. (NW1/4 of SE1/4 of NW1/4 of SE1/4, Sec. 15, T.160N., R.29W.)

Vegetation: Scattered tamarack and black spruce; lush understory consists mostly of sedges with some swamp laurel; ground cover consists mostly of Sphagnum mosses with some cranberry.

Microrelief: 40 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Fibric	0- 90
Hemic	90-180
Sapric	180-195
Silty	195+
clay	

Location: 350 feet N and 1065 feet E of the SW corner of Sec. 22, T.160N., R.29W. (NE1/4 of SE1/4 of SW1/4 of SW1/4, Sec. 22, T.160N., R.29W.)

Vegetation: Scattered black spruce; understory consists of leatherleaf with some Labrador tea, swamp laurel, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

_ayer	Depth (cm)
ibric	0- 30
lemic	30-151
Silt	151+
oam	

Location: 560 feet N and 510 feet W of the SE corner of Sec. 22, T.160N., R.29W. (NE1/4 of SE1/4 of SE1/4 of SE1/4, Sec. 22, T.160N., R.29W.)

Vegetation: Scattered black spruce; understory consists of leatherleaf with some Labrador tea, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses and lichens.

Microrellef: 25 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer Depth (cm)
Hemic 0-315
Clay 315+

Location: 1625 feet S and 2115 feet W of the NE corner of Sec. 23, T.160N., R.29W. (NE1/4 of NW1/4 of SW1/4 of NE1/4, Sec. 23, T.160N., R.29W.)

Vegetation: Scattered tamarack and black spruce; lush understory consists mostly of sedges with some leatherleaf and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses, cranberry, and false Solomon's seal.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Hemic	0-183
Clay	183+
loam	

Location: 110 feet S and 660 feet W of the NE corner of Sec. 25, T.160N., R.29W. (NW1/4 of NE1/4 of NE1/4 of NE1/4, Sec. 25, T.160N., R.29W.)

Vegetation: Scattered black spruce, aspen, and tamarack; lush understory consists of bog birch and leatherleaf with some sedges, bog rosemary, Labrador tea, and grasses: ground cover consists of mosses.

Microrellef: 20 cm.

Depth To Water Table: Not recorded.

Described By: D. Olson and N. Aaseng on October 30, 1979.

Layer	Depth (cm)
Hemic Silty	0-167 167+
clay	

Location: 760 feet S and 365 feet W of the NE corner of Sec. 26, T.160N., R.29W. (NW1/4 or SE1/4 of NE1/4 of NE1/4, Sec. 26, T.160N., R.29W.)

Vegetation: Scattered tamarack and black spruce; understory consists of sedges with some leatherleaf, grasses, swamp laurel, bog rosemary, and Labrador tea; ground cover consists mostly of Sphagnum mosses with some other mosses and pitcher plant.

Microrelief: 45 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-270
Silty	270+
clay	

Location: 2550 feet N and 245 feet E of the SW corner of Sec. 26, T.160N., R.29W. (NW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 26, T.160N., R.29W.)

Vegetation: Scattered black spruce and tamarack; understory consists of leatherleaf with some Labrador tea, sedges, and cotton grass; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Depth (cm
0-200
200-240
240+

Location: 1225 feet N and 1320 feet E of the SW corner of Sec. 27, T.160N., R.29W. (NW1/4 of NW1/4 of SE1/4 of SW1/4, Sec. 27, T.160N., R.29W.)

Vegetation: Lush understory consists mostly of sedges with some leatherleaf, bog rosemary, swamp laurel, and scattered black spruce; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, and cranberry.

Microrelief: 35 cm.

Depth To Water Table: 5 cm.

Described By: N. Aaseng and D. Olson on October 30, 1979.

Layer	Depth (cm)
Hemic	0-122
Loam	122+

Location: 50 feet S and 1975 feet E of the NW corner of Sec. 32, T.160N., R.29W. (NW1/4 of NW1/4 of NE1/4 of NW1/4, Sec. 32, T.160N., R.29W.)

Vegetation: Scattered black spruce, tamarack, jack pine, and poplar; sparse understory consists of some sedges, Labrador tea. leatherleaf, swamp laurel, and cotton grass; ground cover consists mostly of Sphagnum mosses are cranberry with some other mosses.

Microrelief: 45 cm.

Depth To Water Table: Not visible.

Described By: D. Mellem and D. Haverkost on October 29, 1979.

Layer	Depth (cm)
Fibric	0- 60
Hemic	60-135
Silty clay	135+

Location: 490 feet N and 2510 feet W of the SE corner of Sec. 32, T.160N., R.29W. (SW1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 32, T.160N., R.29W.)

Vegetation: Scattered black spruce; lush understory consists mostly of cotton grass with some leatherleaf, swamp laurel, Labrador tea, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrelief: 18 cm.

Depth To Water Table: 10 cm.

Described By: N. Aaseng and D. Olson on October 29, 1979.

Layer	Depth (cm
Fibric	0- 61
Hemic	61-152
Sapric	152-168
Very	168+
fine	
sandy	
loam	

Location: 125 feet N and 2350 feet W of the SE corner of Sec. 34, T.160N., R.29W. (SE1/4 of SW1/4 of SW1/4 of SE1/4, Sec. 34, T.160N., R.29W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists of Labrador tea and leatherleaf with some swamp laurel, bog rosemary, cotton grass, and sedges; ground cover consists mostly of Sphagnum mosses with some other mosses and cranberry.

Microrelief: 20 cm.

Depth To Water Table: Not recorded.

Described By: N. Aaseng and D. Olson on October 30, 1979.

.ayer	Depth (cm)
femic Silty Iay	0-213 213+

Location: 2250 feet N and 250 feet E of the SW corner of Sec. 35, T.160N., R.29W. (SW1/4 of NW1/4 of NW1/4 of SW1/4, Sec. 35, T.160N., R.29W.)

Vegetation: Scattered black spruce and tamarack; lush understory consists mostly of sedges with some bog birch, willow, leatherleaf, swamp laurel, and bog rosemary; ground cover consists mostly of Sphagnum mosses with some other mosses.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Olson on October 30, 1979.

Layer Depth (cm) Hemic 0-244 Silty 244+ clay

Location: 2425 feet N and 490 feet W of the SE corner of Sec. 35, T.160N., R.29W. (NW1/4 of NE1/4 of NE1/4 of SE1/4, Sec. 35, T.160N., R.29W.)

Vegetation: Lush understory consists mostly of sedges with some leatherleaf, bog rosemary, and scattered black spruce; ground cover consists mostly of Sphagnum mosses with some other mosses, false Solomon's seal, cranberry, and pitcher plant.

Microrellef: 30 cm.

Depth To Water Table: At surface.

Described By: D. Mellem and D. Haverkost on October 30, 1979.

Layer	Depth (cm)
Fibric	0-241
Hemic	241-271
Silty	271+
clay	

Location: 770 feet S and 1450 feet E of the NW corner of Sec. 36, T.160N., R.29W. (NW1/4 of SW1/4 of NE1/4 of NW1/4, Sec. 36, T.160N., R.29W.)

Vegetation: Understory consists of sedges with some bog rosemary, leatherleaf, cotton grass, bog birch, and scattered black spruce; ground cover consists mostly of Sphagnum mosses with some other mosses and pitcher plant. Microrellef: 50 cm.

Depth To Water Table: At surface.

Described By: N. Aaseng and D. Olson on October 30, 1979.

Layer	Depth (cm)
Hemic	0-290
Silty	290+
clay	

Severson, Inventors Koochiching County, 940 Ċ of rest resources

