

Effects of Continuous Fishing

MN DEPT OF NATURAL RESOURCES
SH328 .I58 no.378 1984
Thorn, William - Effects of continuous fishing on t



3 0318 00018 6171

Section of Fisheries INVESTIGATIONAL REPORT

No. 378

EFFECTS OF CONTINUOUS FISHING ON THE WALLEYE
AND SAUGER POPULATION IN POOL 4, MISSISSIPPI RIVER

AUGUST 1984

LIBRARY
Dept of Natural Resources
500 Lafayette Road
St. Paul, MN 55155-4021

Minnesota

DNR
SH
328
.I58
no.378
1984

Division of Fish and Wildlife

This document is made available electronically by the Minnesota Legislative Reference Library as part of an ongoing digital archiving project. <http://www.leg.state.mn.us/lrl/lrl.asp>
(Funding for document digitization was provided, in part, by a grant from the Minnesota Historical & Cultural Heritage Program.)

EFFECTS OF CONTINUOUS FISHING ON THE WALLEYE AND SAUGER POPULATION
IN POOL 4, MISSISSIPPI RIVER¹

by

William C. Thorn

ABSTRACT

A 15 year (1967-81) continuous fishing season during 15 years (1967-81) for walleye and sauger in Pool 4 of the Mississippi River had no adverse effects on either population. Although fishing pressure increased as a result of the spring season and during the rest of the year, test netting indicated that the abundance of walleye after 15 years of continuous fishing was similar to the abundance before spring fishing was allowed. The spring fishery accounted for 16.4% of the annual fishing pressure and 17.3% of the annual walleye harvest. The average estimated number of walleye harvested and the catch rate during 1977-81 was 22,457 and 24.0 hr/fish as compared to 1962 at 19,881 and 25.6 hr/fish, respectively. The average weight of walleye harvested in spring was 0.93 kg compared to 0.73 kg for the entire season. The average weight of walleye harvested for the full season since 1967 has varied between 0.65 and 0.80 kg with no trend being evident. The abundance and harvest of sauger has fluctuated widely since 1967 and appears to be unrelated to changes in fishing pressure. Total harvest and fishing success for both walleye and sauger was dependent on year-class strength.

¹ This project was funded in part by Federal Aid Fish Restoration (Dingell-Johnson) Program. Completion report, Study 116 D-J Project F-26-R Minnesota.

Harvest and catch rates of walleye were correlated with the combined abundance of age II and III fish and for sauger with the combined abundance of age classes I-III. Walleye year-class strength was highly correlated with water levels during the spawning and incubation period in April-May. Growth rates of walleye and sauger in Pool 4 are rapid with high rates of total and natural mortality. Both species reach maturity two to three years earlier than populations in northern Minnesota. The combination of rapid growth, high natural mortality and early maturity allows a fast turnover of the Pool 4 populations and thus they could probably withstand even greater fishing pressure without adverse effects on the population.

INTRODUCTION

A closed fishing season during the pre-spawning and spawning periods for game fish has been a traditional fish management tool. In 1967, Wisconsin initiated a continuous season for game fish on its portion of the Mississippi River where it forms the Minnesota-Wisconsin boundary. Minnesota's season remained closed from mid-February to May 1. Confusion for anglers and enforcement problems prompted Minnesota to allow continuous fishing on the Minnesota-Wisconsin boundary waters of the Mississippi River starting in 1969. This was the only area in Minnesota where continuous fishing for walleye (Stizostedion vitreum vitreum) and sauger (Stizostedion canadense) was allowed. Anglers now fish during winter and spring whenever the river is ice-free.

The spring fishery (March-April) was concentrated in the tailwaters of the navigational dams. After several years, local anglers in both states became concerned about or opposed to continuous fishing. The major reasons were the ethics of fishing before and during spawning and resentment toward non-local anglers. Many of the spring anglers were from the Twin Cities metropolitan area.

The Minnesota Department of Natural Resources (MDNR) initiated a creel census at Locks and Dams (L&D) #3 and 4 during March and April in 1968. Results through 1974 were reported by Sternberg (1974b) who recommended further creel census and fish population assessment and noted that fishing would be prohibited within 300 feet of L&D #3 and 4 during March and April starting in 1975. This restriction was based on a MDNR Commissioner's Order allowing the establishment of fish refuges. Wisconsin concurred with the restriction.

Available data included three annual estimates of total fishing pres-

sure and harvest in Pool 4 and indices of population abundance from gill netting since 1965. Creel surveys were conducted under the auspices of the Upper Mississippi River Conservation Committee (UMRCC) at 5 year intervals beginning in 1962 (Daley and Skrypek 1964; Sternberg 1969, 1974a).

Angler concern and opposition to spring fishing persisted. In response, MDNR initiated this 5 year study of walleye and sauger populations of Pool 4 in the Mississippi River. The objective was to determine if these populations were adversely affected by continuous fishing and if more restrictive regulations would be necessary.

STUDY AREA

Pool 4 of the Mississippi River extends from L&D #3 near Red Wing, Minnesota, downstream to L&D #4 at Alma, Wisconsin (Fig. 1). The pool is 71 km long and covers 15,702 ha. Lake Pepin is located in the middle of the pool and is about 35 km long, 2.8 km wide and covers 10,118 ha. There are about 17.7 river km upstream and downstream from Lake Pepin to the locks and dams.

Although the Mississippi River flows through it, Lake Pepin has many characteristics of an inland lake. It is subject to wave action, has little current and has an average depth of 6.4 m. Mean Secchi disc readings in Lake Pepin in 1978-79 were less than 1.0 m. Transparency in the lower portion of the lake was twice that of the upper portion (1.0 m vs. 0.5 m) (Minnesota Pollution Control Agency, personal communication 1981.)

METHODS

Fish populations in Lake Pepin have been assessed with standard experimental gill nets 76 m long by 1.8 m deep with sections of 38, 51, 64, 76 and 102 mm bar mesh. Twenty netting stations were sampled during September-October since 1965 with the smallest mesh set at the 3.1 m contour and




Figure 1. Pool 4, Mississippi River.

larger meshes extending into deeper water. Analysis of the data from 1965-1976 indicated that larger walleye were not being adequately sampled. A 567 m commercial seine with 25 mm bar mesh was used during 1978 and 1979 but also failed to adequately sample larger walleye. Larger walleye were successfully sampled in Lake Pepin by electrofishing during September-October 1979-81 with a boom-type A.C. electrofisher. Gill nets adequately sampled the most abundant sauger age-classes and use of this gear was continued to assess abundance of sauger and young walleye (ages I and II). Data from gill nets were used to develop an index of abundance for walleye and sauger similar to that described by Carlander et al. (1960). Length of all walleye and sauger was measured and a representative sample of weights and scales was taken.

Walleye and sauger were tagged to determine movement and rates of exploitation. During the study, 5,113 walleye and 2,115 sauger were tagged. Minimum tagging size was 356 mm for walleye and 305 mm for sauger. Floy Model FD-67 tags were used in 1976 and FD-68B tags in 1977-80¹. Fish were to be tagged during April at L&D #3 and September-October in Lake Pepin to determine if Lake Pepin fish were being harvested by spring anglers. Signs were posted at access areas advertising a \$50 reward for "lucky tags" as an inducement for voluntary tag returns. Most fish for tagging were captured by electrofishing, however, sauger could not be captured in Lake Pepin by this method. Small numbers of sauger, during the summers of 1978 and 1979, and walleye in 1978 were tagged from seine catches in Lake Pepin.

Angling pressure for walleye and sauger and harvest of walleye and sauger were estimated during 1977-81. A creel survey year began 1 March

¹ Mention of a specific product does not constitute endorsement of that product.

and ended 28 February. Study years in this report are designated by the year they began. A non-uniform probability creel survey (Fleener 1971) and completed trip interviews were used to estimate angling trips per week. A survey clerk counted all boats and anglers completing fishing trips during two 4 h periods/day and 5 days/week at access sites frequented by walleye and sauger fishermen. The clerk interviewed as many anglers as possible during the 4 h periods. "Months" did not conform to a calendar month but a constant number of weeks per month was maintained during the study. All "months" contained four weeks except May, July, October and January which had five. The weekly trip estimate was multiplied by the number of weeks/month to estimate the number of trips/month. This estimate was then multiplied by the average hours/trip for the month to estimate angling hours for the month. The catch/trip was calculated from completed trip interviews and was used to estimate monthly harvest. Monthly estimates of pressure and harvest were summed to calculate annual totals. Walleye and sauger were measured and weighed when possible and a representative sample of scales was taken each year.

The UMRCC annual surveys estimated pressure and harvest based on stratified, instantaneous angler counts and completed and incompleted trip interviews (Daley and Skrypek 1964). The survey year began 1 April and ended 30 March. Walleye and sauger fishing pressure were not initially estimated and were extrapolated from total trips, the percent of anglers seeking these species and the average trip length. Fish were measured and weighed and scales were taken from a representative number of fish.

The annual spring creel surveys at L&D #3 estimated fishing pressure and harvest for walleye and sauger through instantaneous angler counts and fishermen interviews (Sternberg 1974b). The Prairie Island Nuclear Plant

monitoring group provided similar data and estimates for 1975 and 1976 (S. Gustafson, MN Dept. Nat. Res., personal communication 1977).

Rates of exploitation were calculated from voluntary tag returns within one year of tagging. A tag loss of 15% (T.C. Osborn, MN Dept. Nat. Res., personal communication 1983) and a nonreporting rate of 50% (T. Bolland, Iowa Cons. Comm., personal communication 1983) were used to adjust rates of exploitation. Survival rates were calculated from catch curves (Ricker 1975). Scales were read by two readers and a nomograph was used for back-calculating growth to the last annulus.

RESULTS

Population Assessment

The walleye population of Pool 4 appears to have been stable since 1965 (Table 1). Catch per unit effort (CPUE) has not changed significantly for the total or age I walleye ($r = -0.314$ and 0.187 , respectively, $P > 0.05$). The low CPUE in 1978 was probably due to inadequate sampling of age I fish which appeared to be of near normal abundance at age II.

The sauger population of Pool 4 has fluctuated since 1965 (Table 2). Sauger abundance indicates three periods of relative stability: 1965-68; 1970-74; and 1975-81. Abundance increased markedly in 1968 and 1969 and decreased in 1970 and 1975. The mean 1975-81 CPUE was 34% higher than in years before continuous fishing began (1965-66).

Fishing Pressure

Fishing pressure for walleye and sauger on Pool 4 has increased since 1962. The percentage of fishermen seeking walleye and sauger increased from 36% in 1962-63, to 45% in 1967-68, to 57% in 1972-73 (Daley and Skrypek 1964; Sternberg 1969, 1974a) with the corresponding estimates of walleye-sauger fishing trips being 40,597, 76,212 and 82,261, respectively. Estimated

Table 1. Abundance of walleye in fall gill nets (N = 20), Pool 4, Mississippi River, 1965-1981.

| Year | Total | Age 0 | Age I | Age II | Age II+ |
|------|-------|-------|-------|--------|---------|
| 1965 | 53 | 2 | 11 | 16 | 24 |
| 1966 | 69 | 3 | 40 | 16 | 10 |
| 1967 | 72 | - | 17 | 37 | 18 |
| 1968 | 66 | 1 | 14 | 36 | 15 |
| 1969 | 75 | 1 | .6 | 17 | 51 |
| 1970 | 80 | 5 | 28 | 6 | 41 |
| 1971 | 71 | 4 | 22 | 26 | 19 |
| 1972 | 55 | 1 | 24 | 13 | 17 |
| 1973 | 73 | 6 | 40 | 20 | 7 |
| 1974 | 54 | 1 | 34 | 13 | 6 |
| 1975 | 47 | 7 | 13 | 16 | 11 |
| 1975 | 62 | 4 | 29 | 7 | 22 |
| 1977 | 60 | - | 18 | 25 | 17 |
| 1978 | 31 | - | .3 | 13 | 15 |
| 1979 | 68 | 4 | 17 | 17 | 30 |
| 1980 | 68 | 3 | 39 | 5 | 21 |
| 1981 | 61 | - | 36 | 16 | 9 |
| Mean | 63 | 2 | 23 | 18 | 20 |

Table 2. Abundance of sauger in fall gill nets (N = 20), Pool 4, Mississippi River, 1965-1981.

| Year | Total | I | II | III | IV | V | VI | VII+ |
|------|-------|-----|-----|-----|-----|-----|----|------|
| 1965 | 388 | 52 | 169 | 112 | 33 | 20 | 1 | 1 |
| 1966 | 368a | 68 | 63 | 141 | 55 | 30 | 9 | 1 |
| 1967 | 362 | 119 | 157 | 35 | 29 | 12 | 9 | 1 |
| 1968 | 760 | 45 | 346 | 315 | 37 | 11 | 5 | 1 |
| 1969 | 1,253 | 117 | 184 | 757 | 175 | | | 20 |
| 1970 | 873 | 271 | 86 | 172 | 248 | 86 | 5 | 5 |
| 1971 | 952 | 262 | 404 | 83 | 89 | 102 | 10 | 2 |
| 1972 | 807 | 170 | 294 | 229 | 28 | 55 | 24 | 7 |
| 1973 | 869 | 148 | 341 | 253 | 88 | 20 | 18 | 1 |
| 1974 | 1,005 | 432 | 243 | 182 | 92 | 28 | 10 | 18 |
| 1975 | 580a | 53 | 326 | 118 | 27 | 32 | 4 | 19 |
| 1976 | 771a | 166 | 162 | 300 | 92 | 23 | 11 | 15 |
| 1977 | 649 | 186 | 189 | 53 | 167 | 21 | 22 | 11 |
| 1978 | 568 | 35 | 318 | 123 | 40 | 32 | 4 | 16 |
| 1979 | 234 | 39 | 29 | 117 | 28 | 13 | 3 | 5 |
| 1980 | 598a | 184 | 126 | 99 | 130 | 39 | 11 | 5 |
| 1981 | 523 | 202 | 145 | 71 | 12 | 73 | 15 | 5 |
| Mean | 680 | 150 | 211 | 186 | 81 | 37 | 10 | 8 |

a Includes 1-4 age 0.

walleye-sauger fishing trips during the present study averaged 129,814 (+13,399) and ranged from 97,811 (+13,773) to 189,122 (+28,298) (Table 3).

During the present study, fishing pressure for walleye and sauger decreased after the first year and fluctuated little the final four years. Estimated pressure for the 5 years averaged 32.1 hr/ha and ranged from 25.2 to 47.8 (Table 3). The spring fishery accounted for 16.4% of the annual pressure and 12.1% of the trips.

Spring fishing pressure (Table 4) has increased significantly since 1968 ($r = 0.638$, $P < 0.05$). Pressure increased through 1973 after which it has been relatively stable. This spring pressure is assumed to be entirely for walleye and sauger. Spring pressure and mean water level were not correlated ($r = -0.309$, $P > 0.05$) but fishing pressure in 1969 and 1971 was considerably reduced due to high water.

Annual fishing pressure for walleye and sauger has increased more rapidly than can be accounted for by the spring fishery. By 1972-73, the spring fishery had increased to 15,500 trips, about 20% of the annual total. During the present study, the mean estimate of spring fishing trips was similar (15,642) but had decreased to 12% of the annual estimate. The annual estimate of trips increased 42% between 1967-68 and 1977-71.

Walleye Fishery

Success of the walleye fishery appeared to be related to abundance of walleye. Harvest and catch rates of walleye during 1967-68, 1972-73 and 1977-81 were correlated with abundance of age II and III walleye ($r = 0.899$, $P < 0.01$ and $r = 0.891$, $P < 0.01$, respectively) (Figs. 2 and 3). Age II and III walleye were predominant in the creel during three of four years (Table 5).

Walleye catch rates have fluctuated since 1962-63 and during the

Table 3. Estimated fishing pressure (manhours) for walleye and sauger,
Pool 4, Mississippi River, 1977-81.

| Year | Trips | Standard error | Manhours (mh) | Manhours/ha |
|------|---------|-------------------|------------------|-------------|
| 1977 | 189,122 | 14,438 | 751,004 | 47.8 |
| 1978 | 124,704 | 7,874 | 522,641 | 33.3 |
| 1979 | 108,421 | 6,102 | 432,212 | 27.5 |
| 1980 | 97,811 | 7,027 | 396,020 | 25.2 |
| 1981 | 129,012 | 6,836 | 479,738 | 30.6 |
| Mean | 129,814 | 6,836 | 503,820 | 32.1 |

Table 4. Estimated fishing pressure (mh) for walleye and sauger, mean water level (ft msl), harvest, mean weight (kg) and catch rate (hrs/fish), spring creel census near Lock and Dam #3, Mississippi River, 1968-81.

| Year | Manhours | Water level | Walleye | | | Sauger | | |
|------|----------|-------------|---------|-----------------|------------|--------|-----------------|------------|
| | | | Number | Mean weight(kg) | Hours/fish | Number | Mean weight(kg) | Hours/fish |
| 1968 | 22,350 | 668.5 | 4,389 | 0.99 | 5.4 | 6,955 | 0.40 | 3.2 |
| 1969 | 3,540 | 675.1 | 297 | 0.83 | 12.0 | 1,323 | 0.35 | 2.7 |
| 1970 | 41,067 | 669.9 | 5,874 | 0.95 | 6.4 | 13,332 | 0.41 | 3.3 |
| 1971 | 17,455 | 673.6 | 1,835 | 0.89 | 9.7 | 6,629 | 0.58 | 3.0 |
| 1972 | 43,766 | 673.4 | 3,048 | 0.91 | 11.2 | 21,103 | 0.54 | 2.3 |
| 1973 | 75,051 | 673.9 | 4,115 | 0.68 | 18.2 | 35,755 | 0.47 | 2.3 |
| 1974 | 100,935 | 671.6 | 6,049 | 0.74 | 16.7 | 36,514 | 0.59 | 3.0 |
| 1975 | 63,737 | 670.5 | 3,190 | 0.92 | 19.6 | 15,040 | 0.62 | 4.2 |
| 1976 | 73,556 | 672.4 | 2,260 | 0.96 | 33.3 | 33,156 | 0.67 | 2.2 |
| 1977 | 118,019 | 668.6 | 5,969 | 1.16 | 19.6 | 30,658 | 0.58 | 3.8 |
| 1978 | 102,832 | 672.1 | 6,895 | 1.31 | 10.9 | 13,507 | 0.57 | 7.6 |
| 1979 | 68,270 | 673.7 | 2,844 | 0.75 | 23.8 | 8,411 | 0.42 | 8.1 |
| 1980 | 45,199 | 670.9 | 1,037 | 1.11 | 43.5 | 6,750 | 0.47 | 6.7 |
| 1981 | 78,029 | 669.4 | 2,694 | 0.82 | 28.6 | 9,316 | 0.49 | 8.4 |
| MEAN | 60,986 | -- | 3,607 | 0.93 | 18.5 | 17,034 | 0.51 | 3.5 |




Figure 2. Linear regression of harvest of walleye and sauger on abundance of walleye (ages II and III) and sauger (ages I-III); Pool 4, Mississippi River; 1967, 1972, 1977-81.




Figure 3. Linear regression of catch rate (fish/hr) on abundance of walleye (ages II and III) and sauger (ages I-III); Pool 4, Mississippi River; 1967, 1972, 1977-81.

Table 5. Catch rate (hrs/fish), estimated harvest (standard error), yield (kg/ha) and mean weight (kg) for walleye and sauger, annual creel censuses, Pool 4, Mississippi River.

| Year | Hours/Fish | | Harvest | | Yield(kg/ha) | | Mean weight(kg) | |
|---------|------------|--------|---------------|---------------|--------------|--------|-----------------|--------|
| | Walleye | Sauger | Walleye | Sauger | Walleye | Sauger | Walleye | Sauger |
| 1962-63 | 25.6 | 12.3 | 19,881 | 39,624 | a | 1.2 | a | 0.47 |
| 1967-68 | 12.2 | 6.5 | 49,553 | 79,083 | 2.5 | 2.3 | 0.80 | 0.46 |
| 1972-73 | 14.3 | 3.7 | 33,362 | 129,529 | 1.6 | 3.2 | 0.72 | 0.39 |
| 1977 | 22.6 | 8.3 | 33,276(3,465) | 90,893(8,071) | 1.7 | 3.0 | 0.80 | 0.51 |
| 1978 | 17.9 | 7.9 | 29,166(3,594) | 66,539(4,905) | 1.4 | 1.7 | 0.75 | 0.40 |
| 1979 | 25.5 | 13.4 | 16,979(1,431) | 32,446(2,262) | 0.7 | 0.8 | 0.65 | 0.39 |
| 1980 | 28.2 | 8.5 | 14,455(1,689) | 48,640(3,933) | 0.7 | 1.4 | 0.80 | 0.44 |
| 1981 | 25.6 | 10.9 | 18,405(1,290) | 47,173(3,037) | 0.8 | 1.3 | 0.66 | 0.42 |
| 1977-81 | | | | | | | | |
| Mean | 24.0 | 9.8 | 22,457(1,791) | 57,138(5,134) | 1.1 | 1.6 | 0.73 | 0.43 |

a Mean weight not available.

present study were similar to those of 1962-63 (Table 5). Catch rates increased from 25.6 hrs/walleye in 1962, to 12.2 in 1967-68, to 14.3 in 1972-73 and decreased to a 5 year mean of 24.0 hrs/walleye in 1977-81. Monthly catch rates during 1977-81 varied with the best success in May and June (Table 6). These months accounted for 43.3% of the annual harvest compared to 17.3% during the spring fishery.

The spring catch rate during 1968-81 averaged 18.5 hrs/walleye (Table 4) but a significant decrease has occurred ($r = 0.764$, $P < 0.01$). Spring catch rates were not correlated with fishing pressure ($r = 0.272$, $P > 0.05$).

Angler harvest of walleye varied with the largest harvest recorded in 1967-68 (Table 5). The 1979-81 harvests were similar to that of 1962-63. Average harvest and yield for 1977-81 was 22,457 (± 3510) walleye and 1.1 kg/ha. Annual harvest was correlated with pressure during 1977-81 ($r = 0.901$, $P < 0.05$).

The spring walleye harvest during 1968-81 has fluctuated and was dependent on fishing pressure. The average spring harvest was 3,607 walleye (Table 4) and has not changed significantly ($r = 0.002$, $P > 0.05$). Spring harvest was correlated with pressure ($r = 0.653$, $P < 0.05$) but not with mean spring water level ($r = 0.454$, $P > 0.05$) or population abundance as indicated by gill net catches of the previous fall ($r = 0.113$, $P > 0.05$). Spring harvest was correlated with annual harvest during 1977-81 ($r = 0.940$, $P < 0.05$).

The mean weight of walleye creeled has varied little since 1967 (Table 5). The mean weight of the full season for 1977-81 was 0.73 kg and did not change significantly ($r = -0.496$, $P > 0.05$). The mean weight of walleye in the spring creel was 0.93 kg and has not changed significantly ($r = 0.227$, $P > 0.05$) from early creel surveys (Table 4). Mean weight of creeled walleye

Table 6. Seasonal catch rates (hours/fish), harvest (% of annual) and mean weight (kg) of walleye and sauger, Pool 4, Mississippi River, 1977-81.

| Month | Hours/Fish | | Harvest(%) | | Mean weight(kg) | |
|-----------|------------|--------|------------|--------|-----------------|--------|
| | Walleye | Sauger | Walleye | Sauger | Walleye | Sauger |
| March | 21.3 | 7.4 | 9.5 | 9.9 | 1.06 | 0.53 |
| April | 34.9 | 8.0 | 7.8 | 14.1 | 0.94 | 0.50 |
| May | 16.1 | 10.9 | 21.8 | 14.3 | 0.67 | 0.39 |
| June | 14.6 | 9.5 | 21.5 | 12.5 | 0.60 | 0.34 |
| July | 31.1 | 17.1 | 9.7 | 6.4 | 0.60 | 0.30 |
| August | 72.2 | 26.4 | 9.6 | 8.6 | 0.67 | 0.31 |
| September | 51.7 | 17.3 | 8.3 | 11.1 | 0.51 | 0.34 |
| October | 23.4 | 5.2 | 9.5 | 13.7 | 0.58 | 0.45 |
| November | 22.3 | 5.7 | 1.4 | 2.1 | 0.54 | 0.47 |
| December | 26.6 | 13.5 | 0.4 | 1.3 | — | 0.26 |
| January | 132.0 | 9.7 | 0.6 | 3.1 | 0.48 | 0.31 |
| February | 28.4 | 7.3 | 1.9 | 2.9 | 0.98 | 0.46 |

varied by month but generally was greatest during spring (including February when the river was ice-free) (Table 6).

The lower mean weights of walleye between individual years during spring fishing was the result of age I+ fish (immature) entering the creel. The percentage of age I walleye harvested in the spring was positively correlated with year-class strength (percent of age I walleye in previous fall gill nets) (Fig. 4) and mean weight in the spring creel was negatively correlated with the harvest of age I walleye (Fig. 5).

Young walleye were predominant in the anglers' catch. Age-classes I-III were most common in the annual creel (Table 7) and few walleye older than age V were observed. Walleye older than age V were also relatively uncommon in the spring creel of which 26.6% were age I+. The mean percentage of "large" (≥ 610 mm) walleye in the spring creel (1968-74, 1977-81) was 3.7% compared to 3.0% in the annual creel (1977-81). There was a strong correlation between percentage of "large" walleye observed in the two creels during 1977-81 ($r = 0.962$, $P < 0.01$).

Sauger Fishery

Success of the sauger fishery was also related to sauger abundance. Harvest and catch rates of sauger during 1967-68, 1972-73 and 1977-81 were correlated with abundance of age I, II and III sauger ($r = 0.849$, $P < 0.05$ and $r = 0.872$, $P < 0.01$; respectively) (Figs. 2 and 3).

Sauger catch rates have fluctuated with the best catch rates recorded in 1967-68 and 1972-73 (Table 5). The 1977-81 mean was higher than recorded in 1962-63. The highest catch rates were noted in October-November (Table 6).

Sauger catch rates in spring were relatively stable during 1968-76, then decreased and stabilized during 1977-81 (Table 4). Spring catch rates




Figure 4. Linear regression of percent age I walleye spring creel on percent age I walleye in fall gillnets, 1968-81; Pool 4, Mississippi River (1978 data excluded, no spring creel data available for 1975-76).




Figure 5. Linear regression of mean weight (kg) of walleye on percentage of age I walleye in the spring creel; Pool 4, Mississippi River; 1968-81 (no data 1975-76).

Table 7. Age distribution of walleye and sauger in the angler's creel, Pool 4, Mississippi River, 1978-81.
Percent in parentheses.

| Year | Sample Size | Age | | | | | | | | |
|---------|----------------|-----------|-------------|-------------|-----------|----------|---------|---------|--------|--------|
| | | I | II | III | IV | V | VI | VII | VIII | IX |
| Walleye | | | | | | | | | | |
| 1978 | 746 | 23(3.1) | 508(68.1) | 91(12.2) | 90(12.1) | 18(2.4) | 3(0.4) | 6(0.8) | 3(0.4) | 4(0.5) |
| 1979 | 1,029 | 81(7.9) | 668(64.9) | 148(14.4) | 58(5.6) | 34(3.3) | 13(1.3) | 16(1.5) | 9(0.9) | 2(0.2) |
| 1980 | 620 | 111(17.9) | 184(29.6) | 236(38.0) | 36(5.8) | 37(6.0) | 7(1.1) | 6(1.0) | 3(0.5) | |
| 1981 | 1,321 | 687(52.0) | 250(18.9) | 204(15.4) | 130(9.8) | 20(1.5) | 23(1.7) | 4(0.3) | 2(0.2) | 1(TR.) |
| Sauger | | | | | | | | | | |
| 1978 | 2,001 | 101(5.0) | 1,123(56.1) | 444(22.2) | 211(10.5) | 69(3.5) | 28(1.4) | 22(1.1) | 3(0.2) | |
| 1979 | 2,168 | 93(4.3) | 1,004(46.3) | 820(37.8) | 129(6.0) | 98(4.5) | 13(0.6) | 9(0.4) | 2(0.1) | |
| 1980 | 2,722 | 192(7.1) | 750(27.6) | 1,383(50.8) | 262(9.6) | 86(3.2) | 35(1.3) | 13(0.4) | | 1(TR) |
| 1981 | 3,592 | 730(20.3) | 1,418(39.5) | 696(19.4) | 541(15.1) | 130(3.6) | 48(1.3) | 24(0.7) | 5(0.1) | |

were not strongly correlated with increased fishing pressure ($r = 0.304$, $P > 0.05$). The decrease in catch rate between the periods 1968-76 and 1977-81 appears to be the result of increasing pressure and decreasing abundance.

Sauger harvest varied with and was related to fishing pressure. The largest harvest was recorded in 1972-73 and the smallest in 1962-63 (Table 5). Average harvest and yield during 1977-81 were 57,138 fish and 1.6 kg/ha. During 1977-81, harvest was correlated with pressure ($r = 0.924$, $P < 0.05$). Similar harvests were recorded for the months of March-June and September-October (Table 6).

Sauger harvests during the spring were greatest during 1972-78 (Table 4). The spring sauger harvest was also correlated with fishing pressure ($r = 0.670$, $P < 0.01$) but not with mean spring water level ($r = 0.028$, $P > 0.05$). The best harvest years (1972-78) appear to coincide with high abundance but spring sauger harvest was not correlated with the gill net CPUE of the previous year ($r = 0.247$, $P > 0.05$) (Table 2).

The mean weight of sauger caught has varied little since 1962-63 with the average during 1977-81 being 0.43 kg (Table 5). Mean weight of sauger in the spring creel was 0.51 kg and has not changed significantly during the years 1968-81 ($r = 0.323$, $P > 0.05$). Highest mean weights were recorded during 1971-78.

Ages II and III sauger were predominant in the annual creel during 1978-81 but age I were numerous in 1981 as a result of a strong 1980 year class (Table 7).

Walleye and Sauger Movements

Tag returns indicated that walleye from Lake Pepin migrated upstream prior to spawning and moved back downstream immediately after spawning.

During the spring tailwater fishing season in 1977, anglers returned tags from 36 walleye tagged the previous September-October in Lake Pepin. Electrofishing in the same vicinity in spring 1977, 1978 and 1980 captured 12, 19 and 9 walleye, respectively which were tagged the previous September-October in Lake Pepin. The post-spawning fishery in 1978 (May-June in Lake Pepin) accounted for 63% of the first year tag returns of walleye tagged prior to spawning.

Walleye did not move extensively within Pool 4. Average distance moved by walleye tagged in Lake Pepin during 1976 and 1977 and caught by anglers in Lake Pepin was 8.0 and 6.4 km, respectively. Recaptures of tagged walleye by electrofishing in Lake Pepin indicated an average movement of 3.2 km. Few recaptures were reported from upstream or downstream of Pool 4 (Table 8).

Walleye movement upstream out of Pool 4 increased with elevated water levels in spring. The largest percentage of walleye tag returns reported upstream of L&D #3 was spring 1978 during the highest level water recorded during this study (Tables 4 and 8). A weak correlation was found between the percentage of walleye tags returned from upstream of Pool 4 and the mean April water level during the spring of tagging or the spring immediately after tagging ($r = 0.799$, $P < 0.10$).

Movement patterns of sauger appeared to be similar to those of walleye. Most sauger tagged at L&D #3 were taken downstream in Lake Pepin during May-June and few sauger were recaptured other than in Pool 4 (Table 8). Upstream movement prior to spawning was indicated from spring angling near L&D #3 in 1979 and 1980 when 2 tagged sauger were recaptured from the small numbers that were tagged the previous summer in Lake Pepin. Also during spring electrofishing near L&D #3 in 1980, 3 sauger were recaptured that had

Table 8. Location of tag returns from walleye and sauger tagged in Pool 4, expressed as a percentage of the total, Mississippi River, 1976-80.

| Location of Tag Return | Season Tagged | | | | | | |
|----------------------------|----------------|----------------|----------------|----------------|----------------|--------------|--------------|
| | Fall 1976 | Spring 1977 | Fall 1977 | Spring 1978 | Summer 1978 | Fall 1979 | Fall 1980 |
| Walleye (N=708) | | | | | | | |
| Pool 3 and St. Croix R. | 3 | 5 | 8 | 11 | 3 | 8 | 1 |
| Pool 4 | 95 | 90 | 91 | 84 | 94 | 84 | 96 |
| Pool 5 and Chippewa R. | 2 | 5 | 1 | 3 | 3 | 8 | 1 |
| | Spring 1978 | Summer 1978 | Spring 1979 | Summer 1979 | Spring 1980 | | |
| Sauger (N=522) | | | | | | | |
| Pool 3 and St. Croix R. | 7 | 0 | 13 | 0 | 2 | | |
| Pool 4 | 92 | 100 | 86 | 100 | 97 | | |
| Pool 5 and Chippewa R. | 1 | 0 | | 0 | 1 | | |

been tagged the previous summer in Lake Pepin.

Sauger also appeared to exhibit more upstream movement in years of higher water levels. Sauger were not tagged in a sufficient number of years to allow statistical analysis, however, the percentage of sauger tags returned from upstream of Pool 4 was considerably greater for fish tagged in spring of 1978 and 1979 than in 1980 (7% and 13% vs. 2%). Mean April-May water levels in 1978 and 1979 were the highest recorded during 1977-81.

Vital Statistics

Annual mortality of walleye and sauger was calculated to be greater than 50%. Annual mortality of walleye estimated from a catch curve based on creel fish from 1978-81 (ages II-IX) was 55% (Table 7). Annual mortality of sauger based on gill net catches (ages III-VI) was estimated to be 63% for the 1962-75 year classes (from Table 2) and 59% for age classes II-VII based on creel fish for the years 1978-81 (Table 7).

Adjusted rates of exploitation were greater for sauger than for walleye. Rates of exploitation for walleye from 1976 through 1980 were 25.2%, 26.8%, 33.6%, 29.5% and 33.0% with an average of 29.6%. Rates of exploitation for sauger tagged during the springs of 1978-80 and summers of 1978-79 were 32.6%, 33.8%, 58.0%, 27.6% and 37.2% with an average of 37.8%.

The annual expectation of death from natural causes was similar for the two species with natural mortality for walleye estimated to be 25% and for sauger 23%. Natural mortality for walleye was in the higher portion of the range reported by Ney (1978) for exploited populations. Conditional natural mortality rates (Ricker 1975) were 36 and 37% for walleye and sauger, respectively.

Walleye grew faster than sauger in Pool 4, both species grew faster than the Minnesota average and females of both species grew faster than

males and lived longer (Table 9). The Brody growth coefficients (K) (Ricker 1975) for walleye in Lake Pepin were at least twice that of Lake of the Woods on the Canadian border:

| | <u>Male</u> | <u>Female</u> |
|-------------------|-------------|---------------|
| Lake Pepin | 0.329 | 0.310 |
| Lake of the Woods | 0.177 | 0.107 |

Growth increments of walleye and sauger have not changed significantly since 1957 (Tables 10 and 11).

Year-class strength of walleye but not sauger was related to water level during spawning and incubation. The number of age I walleye in gill nets for the 1964-80 year-classes was correlated with mean water levels during 20 April-10 May ($Y = 2.007X - 1330.935$, $r = 0.628$, $P < 0.01$). A similar correlation did not exist for age I sauger (1964-80 year-classes) or age III sauger (1962-78 year-classes) (age at which fully recruited to gill net) in the gill net and water level during April 27-May 17 ($r = 0.131$ and 0.266 , respectively, $P > 0.05$).

DISCUSSION

During 15 years of continuous fishing (1967-1981) for walleye and sauger on Pool 4 of the Mississippi River (Lake Pepin) fishing pressure has increased but neither species has been adversely affected. The walleye population has remained stable while the sauger population has widely fluctuated.

The spring fishing season has added fishing pressure but pressure during the rest of the year has also increased. The estimated fishing pressure for walleye and sauger on Pool 4 of 32.1 hr/ha was similar to or greater than pressure on major Minnesota walleye lakes. Osborn (in press) and Maloney (1978) both estimated about 32 hr/ha for Lakes Winnibigoshish

Table 9. Growth of walleye and sauger (mm) sampled during creel census, 1979-81, (sex undetermined) and during spring electrofishing (sexes separate), 1979-80 in Pool 4, Mississippi River. Growth back-calculated to last annulus only.

| Year | I | II | III | IV | V | VI | VII | VIII | IX |
|--------------------------------|-----|-----|-----|-----|-----|-----|-----|------|-----|
| Walleye | | | | | | | | | |
| 1979-81 average | 168 | 290 | 396 | 485 | 533 | 592 | 650 | 688 | 683 |
| Minnesota average ^b | 130 | 229 | 312 | 384 | 439 | 488 | 531 | 561 | 594 |
| Male 1979-80 | | 295 | 386 | 457 | 511 | 533 | 561 | | |
| Female 1979-80 | | 330 | 437 | 511 | 589 | 620 | 660 | 661 | 699 |
| Sauger | | | | | | | | | |
| 1979-81 average | 142 | 257 | 330 | 394 | 442 | 481 | 505 | 489 | |
| Minnesota average ^b | 109 | 201 | 264 | 300 | 330 | 361 | 363 | | |
| Male 1979-80 | | 259 | 330 | 386 | 409 | 455 | | | |
| Female 1979-80 | | 262 | 438 | 411 | 465 | 478 | 511 | 559 | |

^b From Scidmore (1970).

Table 10. Growth increments (mm) of walleye and sauger collected by creel census (1957, 1967-68, 1979-81), gill nets (1965-68, 1978-81) and spring electrofishing (male and female; 1968, 1979-80), Pool 4, Mississippi River.

| Year | Gear | Year of Growth | | | | | | |
|----------------|----------------|----------------|-----|-----|----|----|----|-----|
| | | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| WALLEYE | | | | | | | | |
| 1957 | Creel census | 150 | 129 | 107 | 69 | 41 | 53 | 25 |
| 1967-68 | Creel census | 165 | 119 | 122 | 66 | 36 | 61 | 46 |
| 1979-81 | Creel census | 168 | 122 | 106 | 89 | 48 | 59 | 58 |
| 1965-68 | Gill net | 163 | 137 | 111 | 72 | | | |
| 1978-81 | Gill net | 160 | 132 | 89 | 79 | | | |
| 1968 Male | Electrofishing | | | 97 | 71 | 25 | 43 | 25 |
| 1979-80 Male | Electrofishing | | | 91 | 71 | 54 | 22 | 28 |
| 1968 Female | Electrofishing | | | 84 | 76 | 53 | 20 | 122 |
| 1979-80 Female | Electrofishing | | | 97 | 84 | 78 | 31 | 40 |
| SAUGER | | | | | | | | |
| 1957 | Creel census | 135 | 94 | 78 | 51 | | | |
| 1967-68 | Creel census | 145 | 106 | 79 | 54 | 73 | 13 | |
| 1979-81 | Creel census | 142 | 115 | 73 | 64 | 59 | 47 | |
| 1965-68 | Gill net | 127 | 119 | 92 | 63 | 49 | 28 | |
| 1978-81 | Gill net | 137 | 125 | 73 | 54 | 56 | 22 | |
| 1968 Male | Electrofishing | | | 79 | 58 | 25 | 13 | |
| 1979-80 Male | Electrofishing | | | 74 | 53 | 23 | 46 | |
| 1968 Female | Electrofishing | | | 91 | 59 | 43 | 23 | |
| 1979-80 Female | Electrofishing | | | 86 | 63 | 54 | 13 | |

Table 11. Calculated and tabular values of t (t and t_{05}) of differences in growth increments of walleye and sauger, 1957-81, Pool 4, Mississippi River.

| | 1967-68 | | 1979-81 | | 1978-81 | | 1979-80 Male | | 1979-80 Female | |
|----------------|---------------------|-----------------------|---------------------|-----------------------|-----------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| | <u>Creel Census</u> | | <u>Creel Census</u> | | <u>Gill Net</u> | | <u>Electrofishing</u> | | <u>Electrofishing</u> | |
| | <u>t</u> | <u>t₀₅</u> | <u>t</u> | <u>t₀₅</u> | <u>t</u> | <u>t₀₅</u> | <u>t</u> | <u>t₀₅</u> | <u>t</u> | <u>t₀₅</u> |
| Walleye | | | | | | | | | | |
| 1957 | -0.224 | 2.179 | -0.450 | 2.179 | | | | | | |
| Creel Census | | | | | | | | | | |
| 1967-68 | | | -0.206 | 2.179 | | | | | | |
| Creel Census | | | | | | | | | | |
| 1965-68 | | | | | 0.212 | 2.447 | | | | |
| Gill net | | | | | | | | | | |
| 1968 Male | | | | | | | -0.052 | 2.306 | | |
| Electrofishing | | | | | | | | | | |
| 1968 Female | | | | | | | | | 0.235 | 2.306 |
| Electrofishing | | | | | | | | | | |
| Sauger | | | | | | | | | | |
| 1957 | -0.248 | 2.447 | -0.355 | 2.447 | | | | | | |
| Creel census | | | | | | | | | | |
| 1967-68 | | | -0.167 | 2.228 | | | | | | |
| Creel census | | | | | | | | | | |
| 1965-68 | | | | | 0.075 | 2.228 | | | | |
| Gill net | | | | | | | | | | |
| 1968 Male | | | | | | | -0.285 | 2.728 | | |
| Electrofishing | | | | | | | | | | |
| 1968 Female | | | | | | | | | 0.000 | 2.228 |
| Electrofishing | | | | | | | | | | |

and Mille Lacs, respectively, while Schupp (1972) reported a 3 year range of 17.4-19.1 hr/ha for Leech Lake.

Walleye fisheries in northern Minnesota lakes subjected to increasing fishing pressure have produced higher yields, a reduced average size and a decline in catch rate (Johnson and Johnson 1971; Maloney 1978; Osborn, in press). Percid populations may exhibit several responses to overexploitation: increased variability of recruitment, increased growth rates and reduced age at maturity (Spangler et al. 1977). None of these responses were evident in Pool 4. Harvest and catch rates were dependent on abundance of walleye and sauger. Average weight in the creel and growth rates of both species have not changed and overexploitation was not indicated.

Annual rates of mortality and exploitation for walleye and sauger were comparable to those of other exploited populations and do not appear excessive. Ney (1978) stated that annual mortality of exploited walleye populations may range up to 57% and that natural mortality may range from 5-30%. This study estimated rates of annual mortality, natural mortality and exploitation as 55%, 25% and 30%, respectively. Recent estimates of walleye annual mortality in Minnesota range from 37% at Leech Lake (Schupp 1972) to 59% for three upper Mississippi River reservoir lakes (Strand 1980). Minimum (unadjusted) rates of exploitation as high as 35% have been reported for walleye in Minnesota lakes (Strand 1980; Osborn, in press). Annual rates of mortality and exploitation for exploited sauger populations of 45% and 10.4-33%, and 87.7% and 22.1% for Missouri River and Tennessee River impoundments, respectively, have been reported (Nelson 1969; Nelson and Walburg 1977; Fitz and Holbrook 1978; Hackney and Holbrook 1978). Rates of annual mortality and exploitation for sauger in this study were 59-63% and 38%.

Angling success for both walleye and sauger was greatly influenced by year-class strength. Strong year-classes of walleye have been shown to dominate the angler's catch in other Minnesota waters (Schupp 1974; Osborn, in press). Serns and Kempinger (1981) found an inverse relationship between exploitation rate and the mean length and age of walleye in Escanaba Lake, Wisconsin.

The most important physical factors affecting walleye reproductive success are water temperature, wind and water level or flow rates (Koonce et al. 1977). Spangler et al. (1977) showed a positive relationship between flow rate and year-class strength for river spawning walleye populations. A positive association between walleye year-class strength and water level at spawning time has been reported for Missouri River reservoirs (Nelson and Walburg 1977) and for lakes with marshes used for spawning (Priegel 1970). It is possible that spring water levels on the Mississippi River may make available both river and marsh spawning habitat and provide sufficient flow rates for adequate reproduction.

Movement of some Lake Pepin walleye upstream to spawning areas and back to Lake Pepin summer feeding grounds appears to be characteristic of many walleye populations (Colby et al. 1979). Movement within the lake was minimal. Schupp (1972) found that walleye tagged in Leech Lake and caught during the summer were generally taken less than 6.7 km from the place of capture.

The fluctuation of the sauger population in Pool 4 was probably due to abiotic factors. The largest increase in sauger abundance coincided with the initiation of continuous fishing. Schupp and Macins (1977) reported that the sauger population in Lake of the Woods fluctuated widely and independently of commercial fishing. Abundance of sauger in Tennessee River

reservoirs fluctuated and year-class strength was attributed to abiotic effects (Hackney and Holbrook 1978).

An examination of biological and physical characteristics indicate that Lake Pepin is a more favorable environment for sauger than for walleye. Sauger have been predominant in both the gill nets and the anglers' catch. Clady (1978) theorized that the presence of sauger usually indicated a reduced walleye population and that the fish community is more diverse in lakes with a well established sauger population than in a lake where the walleye is the predominant percid. At least 85 species of fish have been reported in Pool 4 (Rasmusson 1979). Colby et al. (1979) stated that an inverse relationship exists between walleye vulnerability to angling and abundance of forage fish and, therefore, low catch rates for walleye in the Mississippi River would be expected. Physiology of the percid eye favors sauger over walleye in turbid water (Ali and Ancil 1977). Schlick (1978) concluded that waters with Secchi disc transparencies less than 0.9 m are best suited for sauger.

Anglers opposed to spring fishing assume that all Lake Pepin walleye and sauger spawn in the tailwaters of Lock and Dam 3. This is probably not a valid assumption. Holzer and VonRuden (1982) described off-channel spawning habitat for walleye in Pool 8 as reed canary grass located in flooded timber. Skrypek (1962) netted Lake Pepin during the spawning season and found "green", "ripe" and "spent" females of both species. He concluded that an abundance of excellent walleye spawning grounds existed and that spawning appeared to be scattered throughout the lake. Priegel (1970) found that not all walleye in Lake Winnebago moved upstream into the Wolf and Fox Rivers to spawn but a substantial number spawned successfully over a large length of shoreline in the lake.

Several walleye populations in Wisconsin have been subjected to continuous fishing for some years. Liberalized regulations have been in effect on Escanaba Lake since 1946 and no detrimental effects upon the fish populations have been noted (Kempinger and Carline 1977). The walleye population of Lake Winnebago has been fished during a spawning migration into the Fox and Wolf Rivers for many years, where Priegel (1970) conducted a three-year tagging study and concluded that high fishing pressure prior to the spawning season had little impact on reproductive success in the Wolf River or the population. Holzer and VonRuden (1981) concluded that exploitation of walleye on the Mississippi River did not indicate overfishing and that the continuous season should be continued.

A closed season for walleye prior to and during spawning in inland Minnesota waters has long been in force. Walleye populations in northern Minnesota lakes are characterized by slow growth, late maturity (V-VII) and low natural mortality (5-10%). A closed season may benefit populations with these characteristics. However, Mississippi River walleye in the Minnesota-Wisconsin boundary waters grow rapidly, mature earlier (III-IV) and have a high natural mortality rate. In a population with these parameters, more fishing pressure apparently will not have an adverse effect on the population because of high natural mortality rates. Because of this, it appears that a continuous season would have no adverse effect on the walleye population.

Fish that grow fast and approach their ultimate length quickly are likely to have a high natural mortality rate and fish that grow slowly are likely to have a low natural mortality rate (Gulland 1973). Maturity of walleye is determined by length not age (Ney 1978). Walleye in Lake Pepin are about 100 mm longer than in northern Minnesota lakes by age IV and

mature at ages III-IV. Rates of exploitation in Lake Pepin and many northern Minnesota lakes do not differ greatly (30-35%) but natural mortality rates in Lake Pepin (25%) may be several times greater than in northern lakes. The rapid growth rates combined with a high rate of natural mortality assures a high turnover rate for the population. Under these circumstances, it is unlikely that a reduced season length would "save" walleye in sufficient numbers that could be detected by anglers.

The walleye and sauger populations appear capable of sustaining the harvest and yield recorded in this study and the biological characteristics of these species indicate that a continuous walleye-sauger fishing season can be continued on the Mississippi River without adversely affecting either population.

MANAGEMENT IMPLICATIONS

1. Biological parameters of the walleye population including rapid population turnover because of fast growth, early maturity and relatively high rates of natural mortality differ from those of Minnesota inland walleye lakes and are the basis for the difference in management.
2. Angling success for walleye and sauger is dependent upon abundance which appears to be dependent upon environmental or abiotic factors and neither show any symptoms of overexploitation.
3. Lake Pepin (66% of the Pool 4 area) is a more favorable habitat for sauger than for walleye. This is reflected in both populations and fishery where the sauger are predominant and the walleye tends to be part of an incidental fishery.

4. The walleye and sauger populations of Pool 4 of the Mississippi River have not been adversely affected by 15 years of continuous fishing and more restrictive regulations are not necessary.
5. Annually monitor Lake Pepin walleye and sauger populations.

LITERATURE CITED

- Ali, M.I., and M. Anctil. 1977. Retinal structure and function in the walleye (Stizostedion vitreum vitreum) and sauger (S. canadense). J. Fish. Res. Board Can. 34:1467-1474.
- Carlander, K.D., R.R. Whitney, E.B. Speaker, and K. Madden. 1960. Evaluation of walleye fry stocking in Clear Lake, Iowa, by alternate-year planting. Trans. Am. Fish. Soc. 89:249-254.
- Clady, M.D. 1978. Structure of fish communities in lakes that contain yellow perch, sauger, and walleye populations. Pages 100-108 in R.L. Kendall, ed. Selected coolwater fishes of North America. Am. Fish. Soc. Sp. Publ. No. 11. Washington, D.C.
- Colby, P.J., R.E. McNichol, and R.A. Ryder. 1979. Synopsis of biological data on the walleye Stizostedion v. vitreum (Mitchell 1818). F.A.O. Fisheries Synopsis No. 119. Food and Agriculture Organization of the United Nations, Rome, Italy. 139 pp.
- Daley, S.A., and J. Skrypek. 1964. Angler creel census of Pools 4 and 5 of the Mississippi River, Goodhue and Wabasha Counties, Minnesota, in 1962-63. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep. No. 277: 49 pp.
- Fitz, R.B., and J.A. Holbrook II. 1978. Sauger and walleye in Norris Reservoir, Tennessee. Pages 82-88 in R.L. Kendall, ed. Selected coolwater fishes of North America. Am. Fish. Soc. Sp. Publ. No. 11. Washington, D.C.
- Fleener, G.G. 1971. Recreational use of the Platte River, Missouri. Pages 63-78 in E. Schneberger and J.L. Funk, ed. Stream channelization: a symposium. N. Cent. Div., Am. Fish. Soc. Spec. Pub. No. 2. Omaha, Neb.

- Gulland, J.A. 1973. Manual of methods for fish stock assessment. Part 1. Fish Population Analysis. F.A.O. Rome. 154 pp.
- Hackney, P.A., and J.A. Holbrook II. 1978. Sauger, walleye, and yellow perch in the southeastern United States. Pages 74-81 in R.L. Kendall, ed. Selected coolwater fishes of North America. Am. Fish. Soc. Sp. Publ. No. 11. Washington, D.C.
- Holzer, J.A., and K.L. VonRuden. 1981. Determining walleye movement and exploitation rates in the Upper Mississippi River (Summary Report). Pages 73-81 in Mississippi River Work Unit Annual Report 1980-81. La Crosse, Wisc.
- Holzer, J.A., and K.L. VonRuden. 1982. Determine walleye spawning movements in Pool 8 of the Mississippi River (summary report). Pages 1-15 in Mississippi River Work Unit annual report 1981-82. La Crosse, Wisc.
- Johnson, F.H., and M.W. Johnson. 1971. Characteristics of the 1957-1958 and 1939 sport fishery on Lake Winnibigoshish and connecting waters with special emphasis on the walleye population and catch. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep. No. 312: 30 pp.
- Kempinger, J.J., and R.F. Carline. 1977. Dynamics of the walleye (Stizostedion vitreum vitreum) population in Escanaba Lake, Wisconsin, 1955-72. J. Fish. Res. Board Can. 34:1800-1811.
- Koonce, J.F., T.B. Bagenal, R.F. Carline, K.E.F. Hokanson, and M. Nagiec. 1977. Factors influencing year-class strength of percids: A summary and a model of temperature effects. J. Fish. Res. Board Can. 34:1900-1909.

- Maloney, J. 1978. A qualitative and quantitative census of Mille Lacs, 1976-1977. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Fish Management Rep. No. 7: 18 pp.
- Nelson, W.R. 1969. Biological characteristics of the sauger population in Lewis and Clark Lake. Bur. Sport Fish. Wildl. (U.S.) Tech. Pap. 21: 11 p.
- Nelson, W.R., and C.H. Walburg. 1977. Population dynamics of yellow perch (Perca flavescens), sauger (Stizostedion canadense), and walleye (S. vitreum vitreum) in four mainstem Missouri River reservoirs. J. Fish Res. Board Can. 34:1748-1763.
- Ney, J.J. 1978. A synoptic review of yellow perch and walleye biology. Pages 1-12 in R.L. Kendall, ed. Selected coolwater fishes of North America. Am. Fish. Soc. Sp. Publ. No. 11. Washington, D.C.
- Osborn, T.C. (In Press). The sportfishery of Lake Winnibigoshish and connected waters. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep.
- Priegel, G.R. 1970. Reproduction and early life history of the walleye in the Lake Winnebago region. Wisc. Dept. Nat. Res. Tech. Bull. 45: 105 pp.
- Rasmussen, J.L. 1979. A compendium of fishery information on the Upper Mississippi River. Upper Mississippi River Conservation Committee. 259 pp.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Board Can. 191: 382 pp.
- Schlick, R.O. 1978. Management for walleye or sauger, South Basin, Lake Winnipeg. Pages 266-269 in R.L. Kendall, ed. Selected coolwater fishes of North America. Am. Fish. Soc. Sp. Publ. No. 11. Washington, D.C.

- Schupp, D.H. 1972. The walleye fishery of Leech Lake, Minnesota. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep. No. 317: 11 pp.
- Schupp, D.H. 1974. The fish population structure and angling harvest of Lake of the Woods, Minnesota, 1968-1970. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep. No. 324: 17 pp.
- Schupp, D.H., and V. Macins. 1977. Trends in percid yields from Lake of the Woods, 1888-1973. J. Fish. Res. Board Can. 34:1784-1791.
- Scidmore, W.J. 1970. Manual of instructions for lake survey. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Spec. Pub. No. 1: 47 pp.
- Serns, S.L., and J.J. Kempinger. 1981. Relationship of angler exploitation to size, age, and sex of walleye in Escanaba Lake, Wisconsin. Trans. Am. Fish. Soc. 110:216-220.
- Skrypek, J. Minnesota walleye and sauger spawning investigations - Lake Pepin. Page 72 in Proceedings of the eighteenth annual meeting, Upper Mississippi River Conservation Committee. 161 pp.
- Spangler, G.R., N.R. Payne, J.E. Thorpe, J.M. Byrne, H.A. Regier, and W.J. Christie. 1977. Responses of percids to exploitation. J. Fish. Res. Board Can. 34:1983-1988.
- Sternberg, R.B. 1969. Angler creel census of Pools 4 and 5 of the Mississippi River, Goodhue and Wabasha counties, Minnesota, in 1967-68. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep. No. 306: 53 pp.
- Sternberg, R.B. 1974a. Angler creel census of Pools 4 and 5 of the Mississippi River, 1972-73. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep. No. 331: 48 pp.

Sternberg, R.B. 1974b. Assessment of continuous walleye-sauger fishing at Lock and Dam No. 3 and 4 of the Mississippi River, 1968-74. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep. No. 332: 25 pp.

Strand, R.F. 1980. The walleye sport fishery in three Upper Mississippi Reservoir lakes: Cass, Andrusia, and Big Wolf, 1971-75. Minn. Dept. Nat. Res., Div. Fish Wildl., Sect. Fish. Invest. Rep. No. 368: 38 pp.

Table A1. Estimated fishing trips and standard error (upper) and manhours (lower), Pool 4, Mississippi River, 1977-81.

| | <u>1977</u> Mar 77-Feb 78 | <u>1978</u> Mar 78-Feb 79 | <u>1979</u> Mar 79-Feb 80 | <u>1980</u> Mar 80-Feb 81 | <u>1981</u> Mar 81-Feb 82 | MEAN | MEAN % |
|-----------|--------------------------------|-------------------------------|-------------------------------|-------------------------------|--------------------------------|-------------------------------|--------------------|
| March | 6,412 (2,157) (35,266) | 12,387 (2,021) (64,412) | 6,148 (1,013) (29,510) | 2,962 (567) (12,734) | 6,795 (1,044) (39,411) | 6,941 (1,363) (36,267) | 5.4 (7.2) |
| April | 15,046 (3,467) (82,753) | 7,248 (1,460) (38,414) | 8,426 (1,514) (38,760) | 6,012 (762) (32,465) | 6,775 (878) (38,618) | 8,701 (1,461) (46,202) | 6.7 (9.2) |
| May | 27,821 (4,583) (116,848) | 18,267 (2,498) (76,721) | 15,289 (3,061) (44,338) | 19,864 (4,478) (75,483) | 17,140 (2,784) (61,704) | 19,676 (1,939) (62,516) | 15.2 (12.4) |
| June | 19,688 (5,360) (84,658) | 16,783 (2,792) (65,453) | 15,564 (2,335) (59,143) | 13,671 (2,382) (56,051) | 16,292 (2,204) (58,651) | 16,400 (874) (64,791) | 12.7 (12.9) |
| July | 17,623 (3,685) (68,730) | 12,720 (2,504) (52,152) | 18,049 (2,284) (75,806) | 9,660 (1,899) (39,606) | 14,560 (1,724) (50,960) | 14,522 (1,399) (57,451) | 11.2 (11.4) |
| August | 40,566 (8,569) (121,698) | 19,965 (4,026) (71,874) | 17,896 (2,384) (83,322) | 16,043 (2,181) (62,568) | 29,128 (3,590) (110,686) | 24,720 (4,074) (90,030) | 19.1 (17.9) |
| September | 32,639 (6,894) (130,556) | 22,780 (3,995) (95,676) | 13,969 (2,314) (53,082) | 7,721 (1,124) (33,972) | 18,480 (3,532) (57,876) | 19,118 (3,758) (74,232) | 14.8 (14.7) |

| | | | | | | | |
|----------|----------------------------------|---------------------------------|---------------------------------|--------------------------------|---------------------------------|---------------------------------|------------------|
| October | 12,864 (2,431) (52,742) | 6,905 (1,413) (27,620) | 7,529 (1,337) (27,104) | 11,522 (2,132) (42,631) | 10,300 (1,447) (36,050) | 9,824 (1,023) (37,229) | 7.6 (7.4) |
| November | 1,967 (621) (8,261) | 2,332 (901) (9,095) | 1,480 (468) (4,884) | a1,764 (1,242) (6,742) | 1,278 (354) (4,729) | 1,764 (1,242) (6,742) | 1.3 (1.3) |
| December | 7,503 (2,455) (21,759) | 1,460 (352) (4,818) | 359 (189) (682) | a2,369 (2,489) (6,834) | 154 (90) (77) | 2,369 (2,489) (6,834) | 1.8 (1.3) |
| January | 4,694 (1,015) (18,307) | 2,164 (486) (6,925) | 2,708 (850) (10,561) | 3,431 (686) (12,695) | 3,859 (1,180) (11,963) | 3,371 (394) (12,090) | 2.6 (2.4) |
| February | 2,299 (444) (9,426) | 1,693 (372) (9,481) | 1,004 (268) (5,020) | 2,792 (608) (14,239) | 2,311 (556) (9,013) | 2,020 (276) (9,436) | 1.6 (1.9) |
| TOTAL | 189,122 (14,438) (751,004) | 124,704 (7,874) (522,641) | 108,421 (6,102) (432,212) | 97,811 (7,027) (396,020) | 127,012 (6,836) (479,738) | 129,426 (7,086) (503,820) | |

a No census; mean of other four years used.

Table A2. Estimated walleye harvest, (standard error in parentheses),
Pool 4, Mississippi River, 1977-81.

| | 1977 | 1978 | 1979 | 1980 | 1981 | MEAN |
|-----------|-------------------|-------------------|-------------------|--------------------------|-------------------|-------------------|
| March | 1,461 (506) | 5,693 (2,761) | 1,696 (306) | 503 (107) | 1,380 (238) | 2,147 (813) |
| April | 4,508 (1,105) | 1,202 (286) | 1,148 (244) | 534 (84) | 1,314 (280) | 1,741 (630) |
| May | 6,734 (1,403) | 6,686 (1,124) | 2,245 (615) | 5,335 (1,421) | 3,462 (742) | 4,892 (796) |
| June | 6,838 (2,238) | 5,269 (1,578) | 5,963 (1,025) | 2,634 (614) | 3,494 (676) | 4,840 (696) |
| July | 1,535 (507) | 3,356 (795) | 2,983 (534) | 854 (267) | 2,155 (408) | 2,177 (410) |
| August | 2,997 (1,087) | 2,418 (657) | 776 (225) | 518 (155) | 1,777 (256) | 1,697 (422) |
| September | 4,324 (1,130) | 2,100 (502) | 1,051 (273) | 367 (135) | 1,478 (371) | 1,864 (605) |
| October | 4,072 (893) | 1,027 (277) | 598 (182) | 2,453 (552) | 2,472 (345) | 2,124 (550) |
| November | 420 (170) | 229 (114) | 284 (129) | 320 ^a (36) | 345 (121) | 320 (28) |
| December | 47 (47) | 341 (120) | 0 (0) | 100 ^a (70) | 10 (10) | 100 (56) |
| January | 49 (26) | 133 (80) | 143 (74) | 156 (56) | 151 (75) | 126 (18) |
| February | 291 (92) | 712 (173) | 92 (35) | 681 (99) | 367 (112) | 429 (106) |
| Total | 33,276 (3,465) | 29,166 (3,594) | 16,979 (1,431) | 14,455 (1,689) | 18,405 (1,290) | 22,457 (1,791) |

^a No census; mean of other four years used.

Table A3. Estimated catch per trip of walleye (standard error in parentheses), Pool 4, Mississippi River, 1977-81.

| | 1977 | 1978 | 1979 | 1980 | 1981 | Mean |
|-----------|----------------|----------------|----------------|----------------|----------------|----------------|
| March | 0.23 (0.02) | 0.46 (0.21) | 0.28 (0.02) | 0.17 (0.02) | 0.20 (0.02) | 0.27 (0.05) |
| April | 0.30 (0.03) | 0.16 (0.02) | 0.14 (0.02) | 0.09 (0.01) | 0.19 (0.03) | 0.18 (0.03) |
| May | 0.24 (0.03) | 0.37 (0.04) | 0.15 (0.02) | 0.27 (0.04) | 0.20 (0.03) | 0.24 (0.03) |
| June | 0.35 (0.07) | 0.31 (0.03) | 0.38 (0.03) | 0.19 (0.03) | 0.21 (0.03) | 0.29 (0.03) |
| July | 0.09 (0.02) | 0.26 (0.04) | 0.16 (0.02) | 0.09 (0.02) | 0.15 (0.02) | 0.15 (0.03) |
| August | 0.07 (0.02) | 0.12 (0.02) | 0.04 (0.01) | 0.03 (0.01) | 0.06 (0.00) | 0.07 (0.01) |
| September | 0.13 (0.02) | 0.09 (0.02) | 0.08 (0.02) | 0.05 (0.02) | 0.08 (0.01) | 0.09 (0.01) |
| October | 0.32 (0.04) | 0.14 (0.03) | 0.08 (0.02) | 0.21 (0.03) | 0.24 (0.03) | 0.20 (0.04) |
| November | 0.21 (0.06) | 0.10 (0.03) | 0.19 (0.06) | a a | 0.27 (0.06) | 0.19 (0.03) |
| December | 0.01 (0.01) | 0.23 (0.06) | 0.00 (0.00) | a a | 0.07 (0.06) | 0.08 (0.04) |
| January | 0.01 (0.01) | 0.06 (0.04) | 0.05 (0.02) | 0.04 (0.01) | 0.04 (0.02) | 0.04 (0.01) |
| February | 0.13 (0.03) | 0.42 (0.04) | 0.09 (0.03) | 0.24 (0.03) | 0.16 (0.03) | 0.21 (0.05) |
| ANNUAL | 0.17 (0.01) | 0.23 (0.06) | 0.14 (0.01) | 0.14 (0.01) | 0.16 (0.01) | 0.17 (0.02) |

a No census.

Table A4. Average size (kg) of walleye (standard error in parentheses) in angler's creel, Pool 4, Mississippi River, 1977-81.

| | 1977 | 1978 | 1979 | 1980 | 1981 | Mean |
|-----------|----------------|---------------|---------------|---------------|---------------|---------------|
| March | 1.1 (0.10) | 1.5 (0.13) | 0.8 (0.05) | 1.1 (0.09) | 0.8 (0.04) | 1.1 (0.12) |
| April | 1.3 (0.18) | 0.8 (0.11) | 0.7 (0.06) | 1.2 (0.10) | 0.8 (0.06) | 0.9 (0.10) |
| May | 0.6 (0.07) | 0.8 (0.05) | 0.7 (0.05) | 0.6 (0.04) | 0.6 (0.04) | 0.7 (0.04) |
| June | 0.8 (0.09) | 0.4 (0.03) | 0.5 (0.02) | 0.7 (0.08) | 0.5 (0.03) | 0.6 (0.07) |
| July | — — | 0.6 (0.06) | 0.6 (0.04) | 0.7 (0.08) | 0.5 (0.03) | 0.6 (0.04) |
| August | 0.8 (0.12)) | 0.6 (0.09) | 1.1 (0.23) | 0.2 (0.04) | 0.6 (0.06) | 0.7 (0.12) |
| September | 0.5 (0.12) | 0.4 (0.06) | 0.6 (0.10) | 0.5 (0.09) | 0.5 (0.07) | 0.5 (0.03) |
| October | 0.6 (0.07) | 0.4 (0.06) | 0.6 (0.07) | 0.6 (0.04) | 0.7 (0.06) | 0.6 (0.05) |
| November | 0.5 (0.15) | 0.3 (0.04) | 0.7 (0.18) | — — | 0.6 (0.07) | 0.5 (0.09) |
| December | — — | 0.3 (0.06) | — — | — — | 0.2 (0.00) | 0.3 (0.03) |
| January | — — | 0.4 (0.05) | 0.6 (0.06) | 0.4 (0.04) | 0.3 (0.07) | 0.5 (0.04) |
| February | 1.3 (0.19) | 1.1 (0.07) | 1.1 (0.23) | 0.7 (0.06) | 0.7 (0.08) | 1.0 (0.10) |
| Mean | 0.8 (0.04) | 0.8 (0.05) | 0.7 (0.04) | 0.8 (0.06) | 0.7 (0.04) | 0.7 (0.03) |

Table A5. Estimated sauger harvest (standard error in parentheses), Pool 4, Mississippi River, 1977-81.

| | 1977 | 1978 | 1979 | 1980 | 1981 | Mean |
|-----------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| March | 8,622 (2,923) | 8,392 (1,420) | 3,171 (566) | 1,280 (267) | 6,796 (1,082) | 5,652 (1,310) |
| April | 22,036 (5,155) | 5,115 (1,102) | 5,240 (1,057) | 5,470 (735) | 2,520 (380) | 8,076 (3,158) |
| May | 16,417 (3,089) | 7,102 (1,225) | 3,062 (474) | 10,096 (2,467) | 4,285 (887) | 8,192 (2,136) |
| June | 5,486 (1,755) | 8,327 (1,578) | 7,400 (1,266) | 8,137 (1,704) | 6,338 (1,103) | 7,138 (484) |
| July | 6,018 (1,660) | 3,317 (824) | 2,528 (487) | 3,042 (787) | 3,334 (565) | 3,648 (546) |
| August | 3,797 (1,250) | 11,810 (2,760) | 2,276 (474) | 1,798 (401) | 4,923 (1,133) | 4,921 (1,618) |
| September | 9,860 (2,388) | 13,437 (2,636) | 2,406 (554) | 1,053 (256) | 4,990 (1,044) | 6,349 (2,079) |
| October | 12,282 (2,527) | 4,357 (982) | 4,054 (866) | 10,556 (2,131) | 7,931 (1,302) | 7,836 (1,464) |
| November | 1,338 (447) | 1,249 (543) | 1,216 (483) | 1,184a (76) | 933 (286) | 1,184 (61) |
| December | 1,852 (686) | 832 (251) | 17 (19) | 722a (360) | 185 (130) | 722 (288) |
| January | 1,958 (489) | 430 (147) | 731 (293) | 2,320 (523) | 3,477 (1,122) | 1,783 (495) |
| February | 1,227 (279) | 2,171 (500) | 345 (117) | 2,982 (269) | 1,461 (396) | 1,637 (398) |
| TOTAL | 90,893 (8,071) | 66,539 (4,905) | 32,446 (2,262) | 48,640 (3,933) | 47,173 (3,037) | 57,138 (5,134) |

a No census; mean of four other years used.

Table A6. Estimated catch per trip of sauger (standard error in parentheses), Pool 4, Mississippi River, 1977-81.

| | 1977 | 1978 | 1979 | 1980 | 1981 | Mean |
|-----------|----------------|----------------|----------------|----------------|----------------|----------------|
| March | 1.34 (0.06) | 0.68 (0.03) | 0.52 (0.04) | 0.43 (0.04) | 1.00 (0.04) | 0.79 (0.15) |
| April | 1.46 (0.06) | 0.71 (0.06) | 0.62 (0.06) | 0.91 (0.04) | 0.37 (0.03) | 0.82 (0.16) |
| May | 0.59 (0.05) | 0.39 (0.04) | 0.20 (0.03) | 0.51 (0.05) | 0.25 (0.03) | 0.39 (0.02) |
| June | 0.28 (0.05) | 0.50 (0.05) | 0.48 (0.04) | 0.60 (0.07) | 0.39 (0.04) | 0.45 (0.05) |
| July | 0.34 (0.06) | 0.26 (0.04) | 0.14 (0.02) | 0.32 (0.05) | 0.23 (0.03) | 0.26 (0.03) |
| August | 0.09 (0.02) | 0.59 (0.07) | 0.13 (0.02) | 0.11 (0.02) | 0.17 (0.01) | 0.22 (0.08) |
| September | 0.30 (0.04) | 0.59 (0.05) | 0.17 (0.03) | 0.14 (0.03) | 0.27 (0.03) | 0.29 (0.07) |
| October | 0.96 (0.08) | 0.60 (0.07) | 0.54 (0.06) | 0.92 (0.08) | 0.77 (0.07) | 0.76 (0.07) |
| November | 0.68 (0.14) | 0.54 (0.12) | 0.82 (0.19) | —a —a | 0.73 (0.10) | 0.69 (0.05) |
| December | 0.25 (0.04) | 0.57 (0.11) | 0.06 (0.05) | —a —a | 1.20 (0.58) | 0.52 (0.22) |
| January | 0.42 (0.05) | 0.20 (0.05) | 0.27 (0.07) | 0.68 (0.07) | 0.90 (0.10) | 0.49 (0.12) |
| February | 0.53 (0.06) | 1.28 (0.09) | 0.34 (0.08) | 1.07 (0.08) | 0.63 (0.08) | 0.77 (0.16) |
| ANNUAL | 0.60 (0.02) | 0.58 (0.02) | 0.36 (0.01) | 0.57 (0.02) | 0.58 (0.01) | 0.54 (0.04) |

a No census; mean of other four years used.

Table A7. Average size (kg) of sauger (standard error in parentheses) in angler's creel, Pool 4, Mississippi River, 1977-81.

| | 1977 | 1978 | 1979 | 1980 | 1981 | Mean |
|-----------|---------------|---------------|---------------|---------------|---------------|---------------|
| March | 0.6 (0.03) | 0.6 (0.06) | 0.6 (0.02) | 0.5 (0.02) | 0.5 (0.02) | 0.5 (0.02) |
| April | 0.6 (0.18) | 0.6 (0.03) | 0.4 (0.01) | 0.5 (0.01) | 0.6 (0.01) | 0.5 (0.03) |
| May | 0.4 (0.03) | 0.4 (0.02) | 0.3 (0.01) | 0.4 (0.02) | 0.4 (0.02) | 0.4 (0.02) |
| June | 0.4 (0.07) | 0.3 (0.01) | 0.3 (0.01) | 0.3 (0.02) | 0.3 (0.01) | 0.3 (0.02) |
| July | — — | 0.2 (0.02) | 0.3 (0.02) | 0.3 (0.01) | 0.3 (0.01) | 0.3 (0.02) |
| August | 0.3 (0.08) | 0.3 (0.01) | 0.4 (0.02) | 0.3 (0.02) | 0.3 (0.01) | 0.3 (0.02) |
| September | 0.4 (0.04) | 0.3 (0.01) | 0.3 (0.01) | 0.3 (0.02) | 0.4 (0.01) | 0.3 (0.02) |
| October | 0.5 (0.02) | 0.3 (0.02) | 0.4 (0.02) | 0.6 (0.01) | 0.4 (0.01) | 0.4 (0.04) |
| November | 0.4 (0.06) | 0.4 (0.06) | 0.5 (0.03) | — — | 0.5 (0.03) | 0.5 (0.03) |
| December | 0.3 (0.04) | 0.2 (0.02) | 0.3 (0.00) | — — | 0.2 (0.02) | 0.3 (0.02) |
| January | 0.2 (0.03) | 0.2 (0.02) | 0.5 (0.09) | 0.3 (0.02) | 0.3 (0.01) | 0.3 (0.05) |
| February | 0.6 (0.03) | 0.6 (0.01) | 0.4 (0.03) | 0.4 (0.01) | 0.4 (0.02) | 0.5 (0.05) |
| MEAN | 0.5 (0.01) | 0.4 (0.01) | 0.4 (0.01) | 0.4 (0.01) | 0.4 (0.01) | 0.4 (0.02) |

Table A8. Length-frequency distribution of walleye in the creel, Pool 4, Mississippi River, 1977-1981.

| Size group | 1977 | 1978 | 1979 | 1980 | 1981 |
|------------|------|------|-------|------|-------|
| 175-200 | | | | 2 | |
| 201-225 | | | | 2 | 1 |
| 226-250 | 24 | 16 | 1 | 3 | 9 |
| 251-275 | 25 | 38 | 13 | 2 | 24 |
| 276-300 | 28 | 61 | 48 | 17 | 58 |
| 301-325 | 25 | 107 | 83 | 49 | 150 |
| 326-350 | 49 | 86 | 150 | 92 | 212 |
| 351-375 | 29 | 66 | 198 | 93 | 208 |
| 376-400 | 16 | 60 | 158 | 73 | 99 |
| 401-425 | 35 | 55 | 100 | 89 | 104 |
| 426-450 | 23 | 63 | 64 | 55 | 90 |
| 451-475 | 31 | 37 | 49 | 42 | 77 |
| 476-500 | 19 | 34 | 27 | 24 | 52 |
| 501-525 | 23 | 27 | 31 | 25 | 30 |
| 526-550 | 16 | 29 | 31 | 11 | 20 |
| 551-575 | 5 | 28 | 25 | 6 | 23 |
| 576-600 | 8 | 18 | 23 | 6 | 5 |
| 601-625 | 5 | 9 | 10 | 8 | 8 |
| 626-650 | 1 | 1 | 8 | 9 | 3 |
| 651-675 | 3 | 6 | 4 | 7 | 7 |
| 676-700 | 1 | 3 | 1 | 1 | 4 |
| 701-725 | 2 | 2 | 2 | 2 | 1 |
| 726-750 | 1 | 1 | 2 | 1 | |
| TOTAL | 369 | 747 | 1,027 | 618 | 1,185 |

Table A9. Length-frequency distribution of sauger in creel, Pool 4, Mississippi River, 1977-1981.

| Size group | 1977 | 1978 | 1979 | 1980 | 1981 |
|------------|-------|-------|-------|-------|-------|
| 175-200 | | | | 1 | |
| 201-225 | 8 | 5 | | 1 | 12 |
| 226-250 | 52 | 87 | 18 | 18 | 115 |
| 251-275 | 74 | 187 | 102 | 129 | 316 |
| 276-300 | 94 | 283 | 340 | 267 | 526 |
| 301-325 | 98 | 411 | 559 | 414 | 485 |
| 326-350 | 155 | 285 | 406 | 583 | 436 |
| 351-375 | 178 | 209 | 271 | 554 | 524 |
| 376-400 | 182 | 174 | 165 | 367 | 524 |
| 401-425 | 124 | 150 | 131 | 209 | 321 |
| 426-450 | 78 | 107 | 89 | 137 | 186 |
| 451-475 | 40 | 51 | 48 | 52 | 88 |
| 476-500 | 19 | 37 | 26 | 29 | 41 |
| 501-525 | 4 | 10 | 11 | 11 | 14 |
| 526-550 | 4 | 5 | 2 | 4 | 3 |
| 551-575 | 2 | 1 | 1 | 1 | 1 |
| 576-600 | | 1 | | | |
| 601-625 | | 1 | | | |
| TOTAL | 1,112 | 2,004 | 2,169 | 2 777 | 3,592 |

Table A10. The number of walleye and sauger tagged in Pool 4, Mississippi River, 1976-80, the number and percent returned and percent first year returns.

| Year tagged | Number tagged | Number returned | Percent returned | Number first year returns | Percent first year returns |
|--------------|---------------|-----------------|------------------|---------------------------|----------------------------|
| Walleye | | | | | |
| Spring, 1976 | 302 | 37 | 12.3 | | |
| Fall, 1976 | 1,186 | 163 | 13.7 | 127 | 77.9 |
| Spring, 1977 | 213 | 20 | 9.4 | | |
| Fall, 1977 | 1,231 | 203 | 16.5 | 147 | 72.4 |
| Spring, 1978 | 168 | 214 | 18.3 | 167 | 78.0 |
| Summer, 1978 | 135 | 37 | 27.4 | | |
| Fall, 1979 | 343 | 51 | 14.9 | 43 | 84.3 |
| Fall, 1980 | 535 | 75 | 14.0 | 75 | 100.0 |
| TOTAL | 5,113 | 800 | 15.6 | | |
| Sauger | | | | | |
| Spring, 1978 | 426 | 81 | 19.0 | 59 | 72.3 |
| Summer, 1978 | 171 | 38 | 22.2 | 20 | 52.6 |
| Spring, 1979 | 251 | 64 | 25.5 | 36 | 57.1 |
| Summer, 1979 | 83 | 31 | 16.9 | 29 | 93.5 |
| Spring, 1980 | 1,084 | 308 | 28.4 | 267 | 87.0 |
| TOTAL | 2,115 | 522 | 24.7 | | |

Table All. Return of walleye tagged in Pool 4, Mississippi River, 1976-81.

| Year | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | TOTAL |
|-----------|------|------|------|------|------|------|------|------|-------|------|------|------|-------|
| A - 1976 | | | | | | | | | | 13 | 2 | 3 | 18 |
| 1977 | 2 | 8 | 12 | 24 | 29 | 25 | 6 | 1 | 2 | 2 | 0 | 0 | 111 |
| 1978 | 0 | 1 | 1 | 0 | 5 | 5 | 4 | 2 | 0 | 1 | 4 | 0 | 23 |
| 1979 | 0 | 1 | 1 | 1 | 1 | 2 | 1 | 0 | 0 | 1 | 0 | 0 | 8 |
| 1980 | 0 | 0 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| B - 1977 | | | | | | | | | 1 | 8 | 4 | 0 | 13 |
| 1978 | 0 | 4 | 2 | 9 | 48 | 33 | 24 | 9 | 6 | 2 | 1 | 2 | 140 |
| 1979 | 0 | 3 | 2 | 5 | 5 | 11 | 7 | 2 | 1 | 1 | 0 | 0 | 37 |
| 1980 | 0 | 1 | 0 | 2 | 1 | 0 | 1 | 1 | 0 | 0 | 3 | 0 | 9 |
| 1981 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| C - 1978 | | | | 4 | 65 | 39 | 24 | 12 | 7 | 3 | 1 | 0 | 155 |
| 1979 | 1 | 3 | 2 | 6 | 9 | 9 | 6 | 3 | 1 | 4 | 0 | 1 | 45 |
| 1980 | 0 | 0 | 3 | 2 | 3 | 1 | 2 | 0 | 0 | 1 | 0 | 0 | 12 |
| 1981 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 2 |
| G - 1979 | | | | | | | | | | 1 | 0 | 0 | 1 |
| 1980 | 2 | 0 | 0 | 1 | 16 | 12 | 11 | 0 | 0 | 1 | 0 | 0 | 43 |
| 1981 | 0 | 0 | 2 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 7 |
| H - 1980 | | | | | | | | | | 5 | 1 | 2 | 8 |
| 1981 | 2 | 3 | 1 | 2 | 10 | 9 | 20 | 9 | 11 | 0 | 0 | 0 | 67 |
| 1982 | 0 | 0 | 1 | 2 | 2 | | | | | | | | 5 |
| Total No. | 7 | 24 | 28 | 58 | 196 | 150 | 108 | 40 | 30 | 43 | 16 | 8 | 708 |
| Total % | 1.0 | 3.4 | 3.9 | 8.2 | 27.7 | 21.2 | 15.3 | 5.6 | 4.2 | 6.1 | 2.3 | 1.1 | |

A - Tagged in Lake Pepin fall 1976.

B - Tagged in Lake Pepin fall 1977.

C - Tagged at L&D 3, spring 1978.

G - Tagged in Lake Pepin, fall 1979.

H - Tagged in Lake Pepin, fall 1980.

Table A12. Return of sauger tagged in Pool 4, Mississippi River, 1978-81.

| Year | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | TOTAL |
|-----------|------|------|------|------|------|------|------|------|-------|------|------|------|-------|
| A - 1978 | | | | 1 | 22 | 13 | 5 | 7 | 5 | 2 | 1 | 0 | 56 |
| 1979 | 0 | 2 | 1 | 0 | 0 | 6 | 2 | 5 | 0 | 0 | 0 | 0 | 16 |
| 1980 | 0 | 0 | 0 | 0 | 6 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 7 |
| 1981 | 1 | 2 | 2 | 0 | 1 | | | | | | | | 6 |
| B - 1978 | | | | | | 1 | 0 | 5 | 4 | 1 | 0 | 0 | 11 |
| 1979 | 1 | 0 | 0 | 2 | 1 | 3 | 0 | 2 | 1 | 0 | 0 | 0 | 10 |
| 1980 | 0 | 0 | 0 | 0 | 7 | 1 | 4 | 0 | 0 | 0 | 0 | 0 | 12 |
| 1981 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | | | | 2 |
| C - 1979 | | | | 2 | 13 | 9 | 2 | 4 | 2 | 1 | 2 | 0 | 34 |
| 1980 | 1 | 2 | 2 | 0 | 1 | 1 | 2 | 0 | 0 | 1 | 1 | 0 | 17 |
| 1981 | 1 | 2 | 2 | 0 | 1 | 1 | 2 | 0 | 0 | 1 | 1 | 0 | 11 |
| 1982 | 1 | 0 | 0 | 0 | | | | | | | | | 1 |
| D - 1979 | | | | | | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 3 |
| 1980 | 0 | 0 | 0 | 2 | 9 | 7 | 4 | 4 | 1 | 1 | 0 | 0 | 28 |
| E - 1980 | | | | 12 | 128 | 48 | 39 | 4 | 7 | 14 | 6 | 1 | 250 |
| 1981 | 1 | 5 | 7 | 4 | 16 | 13 | 4 | 2 | 4 | 1 | 0 | 0 | 57 |
| 1982 | 0 | 0 | 1 | | | | | | | | | | 1 |
| Total No. | 5 | 11 | 13 | 25 | 212 | 105 | 54 | 36 | 27 | 23 | 10 | 1 | 522 |
| Total % | 1.0 | 2.1 | 2.5 | 4.8 | 40.6 | 20.1 | 10.3 | 6.9 | 5.2 | 4.4 | 1.9 | 0.2 | |

A - Tagged at L&D 3, spring 1978.

B - Tagged in Lake Pepin, summer 1978.

C - Tagged at L&D 3, spring 1979.

D - Tagged in Lake Pepin, summer 1979.

E - Tagged at L&D 3, spring 1980.

ACKNOWLEDGMENTS

I wish to acknowledge the foresight of J. Skrypek and R. Sternberg who initiated the gill netting and spring creel census, respectively. Also, thanks to B. Hawkinson, G. Grunwald, D. Sogla, M. Ebberts and L. Watson for the many hours spent on creel census, fish sampling and data calculations. Creel census clerks were too numerous to mention. M. Ebberts also spent hours as the second scale reader. D. Schupp supplied much invaluable technical advice and editorial assistance and P. Wingate supplied final editorial assistance.

INVESTIGATIONAL REPORTS*

- No. 368 The Walleye Sport Fishery in Three Upper Mississippi Reservoir Lakes: Cass, Andrusia, and Big Wolf, 1971-75, by Bob Strand. March 1980.
- No. 369 Effects of Winterkill and Chemical Eradication of Fish on a Lake Ecosystem, by Farrell Bandow. April 1980.
- No. 370 The Summer Sportfishery in Voyageurs National Park and Surrounding Waters for 1977 and 1978, by Dennis Ernst and T.C. Osborn. April 1980.
- No. 371 Comparison of Marks on Scales and Dorsal Spine Sections as Indicators of Walleye Age, by Donald E. Olson. December 1980.
- No. 372 Evaluation of Madison, Donaldson and Kamloops Strains of Rainbow Trout (*Salmo gairdneri*) in Lake Superior, by Tracy Close and Richard Hassinger. March 1981.
- No. 373 Experimental Incubation of Fish Eggs in a Moist-Air Environment, by Donald E. Olson. July 1981.
- No. 374 The Effects of Water Levels and Other Factors on Walleye and Northern Pike Reproduction and Abundance in Rainy and Namakan Reservoirs, by Thomas C. Osborn, Dennis B. Ernst, and Dennis H. Schupp. July 1981.
- No. 375 Management of Ponds for Bait-Leeches in Minnesota, by Dirk L. Peterson. July 1982.
- No. 376 Factors Affecting Brown Trout Reproduction in Southeastern Minnesota Streams, by Dennis W. Anderson. May 1983.
- No. 377 Identification of Principal Spawning Areas and Seasonal Distribution and Movements of Muskellunge, by Robert F. Strand. August 1984.

*Complete list of all publications in the series available from Minnesota Department of Natural Resources, Division of Fisheries and Wildlife, Section of Fisheries, Box 12, 658 Cedar St., St. Paul, Minnesota 55155.