This document is made available electronically by the Minnesota Legislative Reference Library as part of an ongoing digital archiving project. http://www.leg.state.mn.us/lrl/lrl.asp

CN 076

IDENTIFICATION OF MOSSES

Introduction

Bryophyta (mosses and their alies) are inconspicuous but common species in many plant communities. Gier (1970) states that "There are about 660 recognized genera of mosses made up of nearly 14,009 species...". This same author states that there are "...many taxonomic problems among mosses, mostly depending upon morphological characters not yet studied and correlated".

We have initiated a study that will produce a species list of mosses on 22 sites (sites no. 1-22 included, Table TB- , Figure TB-). The resons for identifying moss species as a part of the regional characterization for this study in best explained by Steere (1970):

"Bryophytes (mosses and allies) are unusually sensitive among plants, to variations as the substratum upon which they live, not only in moisture, but in acidity, alkalinity, the presence of certain metallic ionic, and other factors." ... "The very close relationship of bryophytes to the soil, rocks, and backs upon which they grow, because of their lack of roots, combined with their sensitivity to the environment, cause them to be excellent indicators of the nature of the environment...".

P GIA

Methods

The complications with the taxonomy of mosses prompted us to contract-our this portion of the study. Miss Rae Barkely, currently employed by the Bell Museum of Natural History, University of Minnesota, Minneapolis Campus, was contracted to collect and identify moss species from 22 sites on the Minesite area. The entire contract (transportation, lodging, supplies, collecting, and identification) amounted to \$500.00.

Miss Barkely traveled to the study area twice during the 1976 season () and spent approximately 30 minutes on each site during each field trip. Mosses were collected from as many substrates as possible, placed in individual paper bags, numbered and dated. Samples were air dried and identified in the Twin Cities by using a combination of a binocular and a compound microscope².

Results

A species list of mosses for each of the 22 sites will be completed and in the possession of the terrestrial team by mid-January, 1977. Over 350 samples are being classified, representing approximately 100 Diffucurt species to date (per. com. with R. Barkely). Questionable speciments will be checked with A well KNOWN BRYCHOGIST, DR. HOWARD CRUM OF THE UNIVERSIT OF MICHAN, ANNE ARBOR & HIS GRADUATE STUDENT, MR. BRE BUCK, A PH. D. CANDIDATE.

Moss specimens will be stored at the University of Minnesota, St. Paul Campus, under the care of Dr. Clifford Wetmore. These speciments will be added to the existing collection and thus available for future studies.

Conclusions

No conclusions or comparisons between habitats can be made at this time.

REFERENCES CITED

Shacklette, H.T., 1965. Bryophytes associated with mineral deposits and solutions in Alaska. In Contributions to chemical prospections for minerals: U.S. Geological Survey Bull. 1198-C, p. Cl-Cl8.

Shacklette, H.T. 1967. Copper mosses as indicators of metal concentrations. In Contributions to geochemical prospecting for minerals: U.S. Geological Survey Bull. 1198-G, p. Gl-Gl&

BRYOPHYTA. THE ENCYCLOPEDIA OF THE BIOLOGICAL SCIENCE (SEC.ED.). STEERC, W.C. 1970. GD. BY PETER GRAY. PUCL. BY VAN NORSTEAND REINHOLD COMPANY, N.Y. : 132-134. GIER, L.J. 1970. Musei. EN THE ENCYCLOPEDIA OF THE BIRDSICAL SCIENCE/ (SEC. ED.). ED. BY PETER GRAT. PUBL. BY VAN NORSTRAND REINHOLD COMPANY, N.Y.: 582-83.